

8

OBSERVATIONS AND CASES

RELATING TO THE

OPERATION

FOR

ARTIFICIAL PUPIL;

IN A LETTER FROM

MR. MAUNOIR OF GENEVA,

TO

PROFESSOR SCARPA OF PAVIA,

WITH THE

PROFESSOR'S ANSWER.

COMMUNICATED BY

DR. MARCET.

Read May 28, 1816.

Letter from Mr. MAUNOIR, to PROFESSOR SCARPA.

SIR,

I HAVE delayed to convey to you the assurance of my gratitude for that confidence in my skill with which you have inspired the Marquis of Beaumanoir, until I should have some interesting information to give you relative to his case, and more especially until I should have in some degree justified your favourable opinion of me. I have been

indebted to you for one of the greatest enjoyments which can be experienced; that of creating, as it were, the faculty of sight, in a person whom time had not yet accustomed to the privation of a sense so precious, and who at the same time had lost nearly every hope of ever recovering it. I am indebted to you also for other, and still greater obligations. The charm of your letters, the friendship and esteem with which you honour me, and the excellence of your instructions have been to me the most powerful stimulants. You make me love my profession with greater ardour, and excite in my mind the liveliest desire of extending its limits. I transcribe for you in the words of the Marquis himself, the following account of the origin of his misfortune; the most trifling circumstances relative to this case cannot but be highly interesting.

“ In the severe winter of 1784, I was first attacked by an inflammation in both my eyes, which prevented me from performing my military duty: for five weeks I had recourse to the following remedies: blisters on the nape of the neck, foot baths, whites of eggs applied to the eyes, &c. In November, 1796, my sight was affected by a second inflammation. I had been wounded about this time and was copiously blooded to remove a difficulty of speech caused by the effects of the wound, which was received in the stomach, grazing the left ribs. Finding myself in the course of the winter affected with an incessant stammering, the bleedings were

repeated, during all which time there was a constant inflammation of the eyes. In the month of May they were restored to their natural state, and I recovered the entire use of my speech; but could neither give myself up to my professional service, nor to any thing which demanded fixed and serious attention.

“ During a journey in Russia towards the end of the same year, I suffered from several slighter attacks of inflammation, which subsided without the application of any remedies. Having been obliged to pass the winter of 1798 in Siberia, the brightness of the snow, the rigour of the climate, the violence of the whirlwinds, to which I was frequently exposed, the ardent reflection of the sun upon the snow and ice, the penetrating smoke which fills the greatest part of the habitations of this dreary country, all these causes contributed to keep my eyes, which were already weak and delicate, in a continual state of inflammation.

“ In 1800, I could neither write nor read. A little while after, I was still able to perceive the light, but less with the left eye than with the right. In the month of August of the same year, during my residence at Vienna, I consulted several eminent oculists, and among others Mr. Beer, who was of opinion that my sight could be restored only by an operation on the left eye, which he performed himself, but without success. Since that period I have

entirely lost the use of that eye ; and the inflammation which resulted from the operation, did not subside, notwithstanding the application of every remedy, before the month of May in the following year.

“ In 1806, I suffered again from another inflammation which lasted three weeks, after which time I entirely lost the power of conducting myself.

“ In 1810, having become totally blind, I placed myself under the direction of Dr. Frederick at Vienna, and in the month of September of the same year, recovered the use of the right eye sufficiently to conduct myself alone. This advantage was but of short duration ; intervals of more or less repose, and of severe inflammation, succeeded each other, until the year 1812, when I again totally lost my sight, preserving only the power of distinguishing, with my right eye, light from darkness.

“ In 1813, I resigned myself entirely to the will of Providence, my medical advisers having declared that they could do nothing more for me. However being at Pavia in the year 1814, I consulted the famous Scarpa, who recommended me to place myself under the direction of Professor Maunoir of Geneva.”

The state of his eyes when I saw them for the

first time at the beginning of April, 1815, was as follows.

They were large, prominent, the upper eyelid rather red than white, and slightly swelled; the conjunctiva covered with vessels a little too full; the iris of a bluish grey colour, and presenting, instead of the pupil in the centre, a white spot of the size of a small pin's head. This spot which could only be the corresponding portion of the capsule of the crystalline, seemed to be adherent to the iris in both the eyes, and was of a whitish grey colour in the left eye, and milk-white in the right. It was impossible to decide *a priori* whether the crystalline was transparent or opaque, since it was entirely hidden by the iris, and by a small portion of the capsule. The Marquis preserved the faculty of distinguishing very clearly the transition from darkness to light; but this power was much weaker in the left eye than in the right.

Saturday, 12th of August, 1815, I operated on the right eye in the following manner.

The patient being seated on a chair and having the head inclined upon a cushion, I placed myself behind him, and with the forefinger of the left hand confining the upper eyelid, whilst an assistant depressed the lower; I made with the right hand a semicircular incision in the lower and external part

of the cornea. This incision occupied a full third of the circumference of the membrane. A quick movement of the eye during the operation, obliged me to bring the edge of the instrument so low as to cut slightly the conjunctiva, which occasioned a slight hæmorrhage; but the application of a sponge dipped in cold water, and a moment's repose remedied this little accident. On reopening the eye, the iris was seen projecting a little from the wound in the cornea. I replaced it with the blunt point of my scissors. Introducing the two blades closed into the anterior chamber, and then opening them, I caused the pointed blade to penetrate the iris, leaving the blunt blade between that membrane and the cornea; then closing the scissors, a perpendicular incision of the iris resulted, describing a little more than half the chord of an arc of two-fifths of the circumference of the iris, traced on the side of the temple. This first incision not having occasioned the formation of a pupil of the necessary size; I introduced the scissors into the iris a second time a little obliquely; and immediately the pupil appeared of a satisfactory form and size, but exhibiting the crystalline entirely opake.

The second stroke of the scissors had divided the capsula: I therefore introduced a small curette, in order to endeavour to destroy what adhered of the crystalline to the shrunk and contracted circumference of the old pupil. It will be seen

presently that this attempt did not succeed. Lastly I effected the passage of a portion of the opake lens by means of a slight pressure with a large scoop exercised on the lower part of the globe of the eye. The crystalline, which was of a cheesy consistence, came out with the greatest ease, and though it was not entirely removed, yet a sufficient quantity was discharged to leave the artificial pupil of a most perfect black. This new pupil was on the side of the temple; and at the exterior and lower part of the iris. The old pupil which had neither changed in form or size, and remained always closed by a white opake body, was not comprised within the area of the new pupil, but was situated internally, and above it. The new pupil had the form of an irregular trapezium. The Marquis received immediately some confused idea of objects in the light.

The eye was reopened the fifth day after the operation. The wound in the cornea was well healed, and the conjunctiva scarcely red. The patient had suffered no pain, but when the eye was exposed to the light, he saw nothing; the new pupil which had preserved its form and size was obstructed by that portion of the crystalline which I had not been able to extract, and which was extremely swelled. I hoped however that this fragment of the crystalline would sooner or later be absorbed either entirely, or in part, and that the new pupil would in the end become black; but I

was much afraid of the formation of a gutta serena. During the first three weeks after the operation, the Marquis of Beaumanoir distinguished no one object, and his despair as well as my anxiety continued daily increasing.

Notwithstanding, I saw the lower edge of the crystalline gradually disappear, and the pupil in the corresponding part become black, and yet in spite of this favourable change, the sight did not return. All at once, early in the month of September, he distinguished first his hand, then his fingers, then the houses in the street. Since this fortunate moment his sight has gradually acquired strength in proportion as the crystalline was absorbed.

At present, December 1, he can read written characters of the size of a fourth of an inch; he distinguishes the hour by the clock; he perceives at the same time the two hands of his watch; he conducts himself with ease in the street or in the road; he avoids obstacles, distinguishing men, women, children, and animals; he does not confound a willow with a poplar, or a poplar with an oak. He evidently gains something every day, and makes a sufficiently apparent progress.

The Marquis wished earnestly that I should operate on the left eye. I did it with repugnance, and not till I had warned him that I expected from

it little or no advantage. September 11, I made an incision in the cornea of this eye, in the same manner as I had before done in that of the right; and at the very instant the iris came out of the wound swelled and enlarged by the crystalline which was spontaneously displaced, so that the lens was in fact without the eye, forming a hernia of which the iris was the bag, veiling it closely, and enveloping it on every side. I gave a stroke of the scissors to the tumor, and the crystalline came out with the greatest ease, to all appearance whole, but extremely small. The iris did not re-enter spontaneously, but it was easy to restore it to its place. The pupil then appearing very small, I gave it a second stroke with the scissors, when the pupil which was of an irregular form, appeared of sufficient size and of a perfect black.

When I examined the eye on the fifth day after the operation, I found the pupil partly obstructed by the opaque capsula, and partly contracted, but still sufficiently visible in the exterior and lower part of the iris, with a small round opening of a perfect black; it appears more than probable that if the retina had not been previously enfeebled by a violent disorder resulting from the first operation, the Marquis would have recovered the sight of his eye to a certain degree, without any difficulty. But the new pupil has remained during nearly a month, like that of an eye diseased with an amaurosis. It

is within a very few days only that this eye can be said to have gained something, and the Marquis can now distinguish the moon, a lighted candle, the fire, or any bright objects which pass before him. The only forms which he has yet been able to ascertain with precision, are the square frames of his windows.

I will not attempt to give you any idea of the form of the crystalline, which remains in the right eye, because it undergoes some change every week, and because it is probable that these changes will last until its entire absorption. I might assist this operation, in making a fresh incision with a needle upon the remainder of the crystalline, but I am unwilling to cause the slightest irritation to the eye, as long as the unassisted force of nature appears to act in an advantageous manner. It will not be till the natural force appears to be languid, or rather entirely inefficacious, that I shall expose by means of an incision a new crystalline surface to the action of the aqueous humour. With respect to the iris of the left eye, there is no crystalline to dissolve, and in consequence I propose to give it shortly a sufficient extent.

I proceed to lay before you some account of two other operations for artificial pupil, of which one took place a few days before that performed on the Marquis, and the other in the course of the last month. I copy simply my notes taken at the time.

“ Mr. Cherbuliez, bookseller, 46 or 48 years of age, has had since his infancy an albugo on the left eye, caused by the small-pox; it occupies a full third of the cornea on the side nearest the temple, and the rim of the pupil almost adheres to this disordered membrane, a very small portion of it is free, which makes the edge of a pupil of the size of a small grain of millet, with which Mr. Cherbuliez can only see indistinctly, with pain, and very near. The cornea of the right eye is irregularly and nearly always obscured by a species of mist which is the consequence of frequent ophthalmia. This same eye had been lately affected with amaurosis, from which it had very considerably recovered. It was but of little use to the patient. The crystalline of the left eye was evidently transparent, and it became necessary to take care of it. I operated on the third of August, 1815, in the following manner :

“ The patient was seated in a chair, his head resting on the breast of an assistant who at the same time raised the upper eye-lid. I placed myself behind him, and with the forefinger of the left hand, depressed the lower eye-lid. I then made with the right hand a semicircular incision of about three lines in length, and as much upon the opake part of the cornea as upon the transparent.

“ I then introduced into the wound my blunt scissors, and penetrating with one of the blades through

the small pupil which I have mentioned, I succeeded in making an incision without wounding the capsula or the crystalline, cutting the orbicular muscle in a transverse direction, and continuing nearly parallel between the fibres of the diverging or radiating muscle; this incision describes an entire radius of the disc of the iris. Its immediate result was a perfect triangular pupil of the most brilliant black.

“ Every thing went on well during the first 50 hours, when Mr. Cherbuliez having made a violent effort in going to stool, caused a small hernia of the iris. This accident did not change the form or the size of the pupil. I have three times effected the depression of this rupture of the iris, by introducing a needle for the cataract. At present the staphyloma is little or nothing.

“ Mr. Cherbuliez has resumed the ordinary course of his affairs, he writes and can read the finest written characters, nor is his eye fatigued even by a long series of occupation.”

I come now to the account of the last operation which I have performed.

“ A woman of the name of Saillard, 26 years of age, and mother of an infant of some months, received a thorn in her left eye, as she was cutting wood in a forest, about two years ago. She did

not suffer much at the time, but the consequences of the accident were fatal to the sight of the eye. Her state when I saw her for the first time was as follows :

“ The pupil which was shrunk and irregularly fringed, and entirely incapable of any movement of contraction or dilatation, was partly covered by a spot or shell, milk-white and apparently in conjunction with the iris. Where the pupil was not covered by this body, it appeared to me at first sight black, and the woman was still able to distinguish objects confusedly with this eye. The right eye had had a cataract for nearly a year, and the pupil was immoveable and fringed, the iris appeared to have been diseased, and the watery humour had not a healthy transparenence.

“ Being much occupied by a press of business, and not wishing to occasion a loss of time to the patient ; thinking besides that the crystalline of the left eye was transparent, and that I had nothing to do but to increase the pupil which was too small, I took with me only my scissors blunted at both points ; but at the moment of the operation, and favoured by a bright day, I perceived that the crystalline was decidedly opaque, though at the same time in such a manner, that I could easily have supposed it to be transparent the first time I observed it, and especially in a situation where the light was not favourable.

“ It was now evident, however, that it was necessary to extract the crystalline, and that if I enlarged the pupil with my blunted scissors, it could only be by cutting the iris in the upper part of its disc, an operation which would render the extraction of the crystalline difficult, and even perhaps impossible, since it was probable that some extraneous substance would adhere to it. In consequence of these considerations, I made my decision instantly, and performed the operation in the following manner :

“ I first made an incision in the lower part of the cornea ; then introducing the point of my instrument into the lower part of the iris, the puncture gave me an opening of the size of a pin’s head, sufficient to allow the passage of one of the blunted points of my scissors, with which I cut the narrow strip comprised between the hole caused by the knife and the natural pupil ; I then obtained immediately a pupil of a good size, and which exhibited a crystalline humour of a whitish grey colour. Then observing that the compression which I exercised upon the eye appeared in no degree to move the crystalline, I passed behind it a very small scoop, and drew it out of the eye by this means, with the greatest facility. The pupil was then of a perfect black, but the white spot had changed its situation and now obstructed the lip of the wound in the cornea. After having endeavoured in vain to remove it with the scoop, I wa

obliged to seize it with the forceps, by which means I drew it out, together with a transparent membrane which adhered to it, and seemed to detach itself from the iris: it proved on examination to be that of the crystalline. The white body was of the size of a lentil, and entirely of a cartilaginous nature. The patient saw immediately, and distinguished perfectly every object which I presented to the eye, but they all appeared coloured with a tinge of blue.

“ Six days after the operation, I shewed the woman to my society of medicine, where she was seen by Messrs. Odier, Peschier, Aubert, Colladon, De Roches and Morin. The cicatrix of the cornea was in a healthy state, the pupil of a good size and of a perfect black, the sclerotica was scarcely red, and the patient saw very small objects tolerably well. She returned to her house at Chable the next morning, seven days after the operation.”

I have the honour to be, &c. &c.

J. P. MAUNOIR.

Geneva, December, 1815.

Letter from PROFESSOR SCARPA to MR. MAUNOIR.

SIR,

I have received with the highest satisfaction the account with which you have favoured me of the success of your operation on the Marquis of Beaumanoir. It gives me great pleasure to have been in any degree instrumental to the service you have rendered him, and I request you to present to him my hearty congratulations.

Every lover of the science cannot but feel interested in the narration of the difficulty which attended this operation, as well as that performed on Mrs. Saillard, on account of the opake crystalline which in both cases adhered to the lower circumference of the iris, and the edge of the closed pupil. This has given me occasion to make some reflections on the subject, which I take the liberty of communicating to you with my usual freedom.

The detail of both the operations in the above mentioned cases, exposes a certain degree of difficulty and uncertainty in the proceeding, which I should wish to see removed or corrected; and have great hopes that you will be able to attain this desirable object.

In the first place, I am of opinion that it is not necessary to be scrupulous whether the crystalline

be partly or entirely opake, whenever the capsule is opake, and adheres to the iris behind the edge of the interior and inclosed pupil. In this case, only one remedy can be pointed out, namely, the removal of the opake adherent capsule, and consequently of the crystalline, whether it be transparent or opake.—In the second place, I think there is no reason to doubt that, in similar cases, it is adviseable to make an incision upon the iris proportioned to the size of the body to be extracted, rather than to make it small, which obliges the operator to divide the crystalline and the capsule, with the intention of extracting a part, and of abandoning the rest to the powers of absorption.—Thirdly, I would establish as a fundamental principle in similar cases, that after the complete extraction of the crystalline with its opake capsule, by means of the least possible introduction of instruments, the artificial pupil ought not to be too near the incision in the cornea, and, consequently, not too near the cicatrix occasioned by it.

The causes of the obstacles, to which you were exposed in the two cases above mentioned, may, I think, be perceived from a consideration of the principles I have just stated. In both the incision in the iris was too small in proportion to the size of the body to be extracted; and in both, the position of the artificial pupil was very disadvantageous; that is to say, on the side of the temple and close to the incision in the cornea.

After reflecting attentively upon this situation of the artificial pupil, and upon the obstacles which it presents to the operator, it appears to me, if I am not greatly deceived, that a method of operation compounded of that of Wenzel, and of your own, would perfectly answer the desired end. Wenzel, as you are aware, made an incision upon the cornea and the iris with a single stroke of his instrument, taking care that this transverse incision should pass through or underneath the centre of the inclosed pupil. He then took off, by means of the scissors, a portion of the edge of the iris, for the double purpose of extracting with facility the opaque crystalline with its capsule, and of leaving a permanent artificial pupil of sufficient size. In the method which I would suggest, after having made, in the manner of Wenzel, a transverse incision in the iris and in the cornea, I would introduce your scissors, blunted at both points, into the anterior chamber of the aqueous humour, with which I would make an incision in the iris, diverging from the cut made by the knife, so that your usual triangular edge might be the result, having a curvilinear side*. This aperture, which requires only a single stroke of the scissors, will be, I think, sufficiently large to allow easy egress to the crystalline and the capsule; and this so much the more easily, in proportion as the point of adherence of the capsule to the iris, is comprehended either entirely, or in a great mea-

* See Plate V. Fig. 1.

sure, within the two incisions. By this means the facility of making the crystalline and the capsule pass obliquely out from the iris will be increased on account of the enlarged space that will result from the cut with the scissors diverging from that made by the knife; and I should prefer this incision with the knife to the puncture made by you in the iris of Saillard, to afford a passage to the blade of the scissors. Besides this, the direction and the situation of the triangular edge of the iris will be calculated to leave a pupil not only permanent, and sufficiently large, but also placed opposite to the cut in the cornea, and accordingly more convenient for the purposes of vision; especially if it fall upon the side of the iris nearest the nose, which ought, if possible, always to be the case.

I have thus briefly given you my opinion on this subject. You will recollect that you permitted me to make objections and suggestions. I have done so; and it now belongs to you in the course of your practice to make trial of the method I propose, and either to confirm or reject it, or, which will not be difficult to you, to suggest some new and more practicable expedient. It is certain that this complicated case of the pupil, that is to say, where the crystalline is found opaque, together with the capsule adhering to the iris, requires an exertion of genius and skill, united to a more perfect method of operation, than that which has been hi-

therto practised. Your researches have already been so numerous and so successful in this branch of our profession, that we have a right to expect you to proceed in the completion of your work. Till within a very few years, our knowledge respecting the artificial pupil, and the method of conducting the operation, was involved in great obscurity, and practice was sometimes even in opposition to the known anatomy of the eye. It is by solid and fixed principles alone that we ought to be guided in all that variety of complicated cases, which frequently accompany and aggravate the inclosed pupil. In estimating the extent of the services which you have already rendered to the profession, I run no risk of errour in enumerating the following facts as the results of your most useful researches.

1. That no instrument is so proper as the scissors for making an incision in the iris.

2. That to do this, when there is no complication resulting from a cataract, a very small incision in the cornea is sufficient, about half the size of that which is made for the extraction of the crystalline,—a fact which on many accounts is of the highest importance.

3. That the formation of a triangular edge in the iris by means of a double incision with the

scissors, is the most easy and least painful of all the methods hitherto proposed for obtaining a *permanent* artificial pupil.

4. That the spots of the cornea present no obstacle, because it is possible to produce the artificial pupil in that part of the cornea remaining transparent, in the quarter opposite to that in which the incision is made,—a fact of the greatest importance.

5. That it is possible to obtain the artificial pupil without injury to the crystalline or its capsule, whenever these parts are preserved transparent, in spite of complete confusion in the iris.

If to these advantages, which your method of operation possesses over all those hitherto practised, you are able to add that of rendering as little laborious as possible the manner of making the artificial pupil, in those cases where it is necessary to remove at the same time the crystalline and its opaque capsule adherent to the lower surface of the iris, you will fulfil not only all my wishes, but those of all who are interested in the cause of humanity, and in the progress of science. The state of my health does not permit me to follow you in this useful career, and leaves me only the power of witnessing and of applauding your success. I have much pleasure in informing you that your scissors have begun to be used here, and Signor Morigi,

my successor in the chair of surgery, has lately availed himself of them with great success in an operation for the artificial pupil. He is a man of considerable ability, whom you will find mentioned with deserved praise in several places of my work.

I have the honour to be your most sincere friend and humble servant,


A. SCARPA.

Pavia, Jan. 1, 1816.

THE END.

9.

Treatise on Opthalmy.



9..

A
TREATISE
ON
OPHTHALMY;

AND THOSE
DISEASES WHICH ARE INDUCED BY INFLAMMATIONS
OF THE EYES.

WITH
NEW METHODS OF CURE.

BY
EDWARD MOORE NOBLE,
SURGEON.

PART THE FIRST.



Birmingham:

PRINTED BY SWINNEY & HAWKINS;

FOR
G. G. & J. ROBINSON, PATERNOSTER-ROW, LONDON;
CRZECH, EDINBURGH; AND M'KENZIE, DUBLIN.

1800.

1770

W. J. F. F. F.

W. J. F. F. F.

W. J. F. F. F.

1650231



PREFACE.

TO practice any particular branch of the medical profession, with success, requires a general knowledge of the whole science; at the same time, it must be acknowledged, that if, from any peculiar coincidence of circumstances, a person's attention is more immediately directed to one class of diseases, he acquires a greater power of discrimination, and, from the mind dwelling much on one subject, he frequently obtains new ideas, and strikes out into a new and preferable mode of practice.

It is of little importance to my readers, to be told, how my attention came to be directed more particularly to the treatment of diseases of the eyes; it is sufficient to inform them, that for some years, I have had a considerable practice in that department of my profession, and that the hopes of being able to improve that particular branch, has principally induced me to publish this Treatise.

It is not however on that account solely, that I have ventured to appear before the public. The art of curing diseases has, of late years, made rapid strides towards arriving at that *acmé* of perfection, beyond which it is not in the power of man to advance it. The light which has been let in upon us, by the doctrines of that great

genius, Dr. John Brown, and so ably seconded by a Darwin, and a Beddoes, has laid the foundation of a new era in the annals of Medicine, and has opened new views to the practitioner, in the theory and treatment of diseases.

Since that epoch no regular treatise, on the inflammation of the eye, has appeared, in which the mode of treatment has been founded on systematic principles; and when we consider the importance and delicacy of that organ, we shall be surprised to find, how little has been done, towards arriving at a rational plan of cure.

The regular routine of bleeding, blisters and cathartics, with a variety of external applications, are generally indiscrimin-

ately applied, with little attention to the nature, or the period of the disease; and there are some practitioners who act with still greater inconsistency, and maintain, that it is useless to inspect the eye; that it causes an unnecessary degree of irritation; and that we ought to trust entirely to internal medicines.

It is my intention to endeavour to point out the propriety of a certain mode of treatment, and to discriminate the proper time for the use of different applications.

On the first attack of an Opthalmy, the general circulation of the mass of fluids is not evidently increased, though it may be so in a short time, from the pain and irritation of the eye; I therefore consider

the disease as a topical affection, and that we ought chiefly to rely on local applications for a cure.

In speaking of the causes of Opthalmy, I have divided them into an increase of stimulus, or an increase of that something to which I have given the name of the irritable principle, but whether it ought to be considered as a quality of vitality, or as supplied by the brain, or from the atmosphere, I have not ventured to risk an opinion.

It probably may be thought, for a Treatise on the Opthalmy, I have entered into too long a digression upon the effects of *stimuli* applied to the body, and have been too prolix upon the laws relative to the accu-

mulation or exhaustion of this irritable principle. Much more might undoubtedly have been said, which would have better elucidated the doctrine, but I wished not to take up too much of my reader's time upon the discussion, and I knew not how, in saying less, to properly explain my ideas on those subjects.

It may, perhaps, not be improper in this place, to take a general view of what is intended to be given in the Second Part which will conclude this Treatise.

In the first place I shall enter upon the cure of the inflammation of the eye.

In the laws of the animal economy, there is scarce any fact more clear than that a

stimulus stronger than usual being applied to the moving fibre, makes it less easily excited into action, and that on the sudden subtraction of this increased *stimulus*, the motions of the part will be diminished.

Upon this law will depend my method of cure of the Ophthalmia. The treatment of the disease admits of a variety of modifications ; but my principal object will be the application of a *stimulus*, in a peculiar manner, as great as the eye can bear, without being thrown into convulsive motions, and when this *stimulus* loses its effect of causing pain, to suddenly remove it, and diminish all *stimuli*, or irritating causes, as much as possible.

By these means a diminished action of the vessels will be induced, the pain will be moderated, and an alleviation of the symptoms will take place.

By convulsive motions are meant, those motions which take place from the injudicious application of too powerful stimulants ; as the tincture of opium, which has been so much extolled, under the name of the *Tinctura Thebaica*, by Mr. Ware, from an old *formula* of the College of Physicians. The tincture of opium is the most efficacious application that has ever been recommended to the public for inflammations of the eye ; but, like all other powerful medicines, whose *modus operandi* we are not well acquainted with, it is daily employed improperly, to the great pain and distress

of the patient. It will be my endeavour, in the succeeding part, to lay down rules for its application, and to explain on what its salutary effect depends.

The better to explain my mode of treatment and its effect, I shall insert a few cases, by which I hope the rational practitioner, in prescribing for inflammations of the eyes, will be enabled to appropriate the applications to each individual subject, and may act with a rational probability of success.

I shall then proceed to the cure of the Ophthalmia which has been of long continuance, and has put on an atonic form. This will be followed by the cure of different diseased actions of the vessels of the

cornea and *conjunctiva*. To the whole will be subjoined, a concise inquiry into the powers and efficacy of those applications which are generally made use of in inflammations of the eyes.

I hope to be able to send the Second Part, which will conclude this Treatise to the press the beginning of next year.

Birmingham, September 1,
1800.

ERRATA—Page 128, line 21, for averted, read everted.
— 144, — 14, for infection, read affection.





TREATISE

ON THE

OPHTHALMY, &c.

INTRODUCTION.

THE Eye, from its peculiar organization and the delicacy of its structure, with the sickness and variety of its motions, besides being exposed to all those things which excite inflammation in general, is affected by a variety of noctious powers, which applied to another part would prove perfectly harmless: on these accounts, inflammations of the eyes are perhaps more frequent than of any other part, and, though not often dangerous

to life, are always troublesome, and, from the importance of the organ concerned, render a speedy removal of it extremely desirable.

Although total blindness from an Ophthalmia is not very frequent, yet a partial defect in the sight is not uncommon, and cases unfortunately sometimes occur in which the total loss of that important sense, with considerable pain and deformity, is the termination of the disease.

The hopes of being able to lay down a more rational plan of cure, and to point out the impropriety and danger of that indiscriminate manner followed in the treatment of Ophthalmia, which is too generally adopted, induced the Author to send into the world the following pages. After treating on the history, the causes, and cure of the disease, he will concisely inquire into the efficacy of a great number of applications to the eye which are in general and esteemed use.



THE DEFINITION AND HISTORY OF OPHTHALMY.

WHAT is usually meant by the term Ophthalmy, is a redness and inflammation, with pain, of the *tunica conjunctiva*, or that membrane which covers the anterior part of the eye, where it ceases to be transparent, and lines the inside of the lids ; with pain of the eye, on exposure to light ; for though an inflammation will sometimes arise in the coats and globe of the eye itself, as a primary disease, yet that is so very seldom the case, comparatively with the frequency of the inflammation of the *tunica conjunctiva*, is not, commonly, to be separately considered.

On the first attack of an Ophthalmy, the *tunica conjunctiva* becomes slightly red, with an increase of heat of the part ; the eye feels dry, and, at times, as if there were small particles of dust in it ; the lids do not move with their accustomed ease, and

the light causes some uneasiness. As the inflammation increases, the whole of the *adnata* becomes more red, and, to that degree, that, in a violent attack, there is not the least appearance of whiteness, neither can any vessel be distinctly seen, but an uniform fiery redness extending over the whole surface; the thickness of it is generally a little increased, and at times it is so much enlarged, as to make the *cornea* look as if it was depressed or sunk in the globe; which indicates a very violent disease, and has been called *chemosis* by authors: the sensation of heat becomes greater, and the patient feels as if pins or needles were piercing the eye, with a discharge of tears: the light becomes intolerable, even a small quantity produces great pain, with a copious flow of tears, and the lids are red and swelled. The pain is now not confined to the eye and lids only, but extends to the temple and forehead, and is sometimes diffused over the whole head. The irritation and pain produced are occasionally so

great, as to excite a considerable degree of fever, which, at times, though very seldom, is so great as to induce delirium.

Though the whole of the *conjunctiva* is generally inflamed, in nearly an equal degree, yet at times the redness is confined only to a small part, the other portion preserving its natural colour and appearance. When this happens, it will usually be found, that either some small extraneous body is adhering to the eye, or that there is a small pustule or ulcer on the *adnata*.

The light also, which, even in slight ophthalmies, commonly gives pain, sometimes causes very little irritation, though the appearance of the eye would strongly indicate the contrary; which must arise from the inflammation being confined to the external parts of the eye. On the other hand, in some cases, in which we suppose the *tunica boroides* and the internal parts mostly to be affected, though the appearance of the *con-*

junctiva shall be little changed, the smallest quantity of light causes great pain.

The redness or turgescence of the vessels on the first appearance of the ophthalmia never extends to the *cornea*, but is produced by the increased and diseased action, by which new vessels are formed, or the old ones elongated. It does not appear probable, that the vessels, which are seen on the *cornea transparens* are caused by the diameters of the old ones being enlarged, and allowing admission to the red particles of the blood, as they are never visible till the inflammation has continued some time, though it should be very violent and the action of the part great ; besides, were the vessels merely enlarged, the whole substance of the *cornea* would be red, whereas, unless in very severe and obstinate cases, it is always confined to the external surface, and frequently the vessels are very few in number, with generally either a thickening and opacity of the *cornea*, or an ulcer in the same part.

The discharge of tears is usually rather copious and the eye always moist; but cases are sometimes met with, in which the eye has been dry from the beginning, and continues so during the whole of the complaint: this arises from the irritation not being so great as to produce the sensation of pain, and on that account, the action of the lachrymal gland is not called into undue exertion.

It often happens, that either the discharge from the lachrymal gland, or from the small exhalent arteries of the *conjunctiva*, is so thickened as to form an adhesive matter, which firmly glews the lids together, more especially in a morning, after sleep. This, as it adds much to the irritation and pain in the part, should be prevented by the application of some bland soft ointment, to the edges of the lids, at bed time.

If the inflammation, from the beginning, has not been violent, or attended with much pain, and does not soon subside, from some

peculiar want of action in the part, or from the inflammation being occasionally increased, by want of attention to remove any exciting causes, or the use of improper applications, the eye will become so accustomed to its new motions, that the inflammation may be continued for a great length of time, without occasioning any opacity of the *cornea*, or being productive of much pain, and has generally been called the *atonic* or passive inflammation.

But when the inflammation has continued with violence for some time, the vessels assume various kinds of diseased action, producing different effects on the eye. Hence when the irritation and pain have been great, the arteries are thrown into a kind of action, by which they secrete a humor which is deposited in different parts of the eye. As it is only the minute extremities of the vessels, that have this secretory power, and those on the *conjunctiva*, that the looseness of the cellular membrane

that surrounds them, are easily expanded, any formation of matter is seldom found there. The matter is at times discharged into the anterior chamber of the aqueous humour, where, descending to the bottom, by its own weight, it has an appearance similar to the white speck at the root of the nails, and has been called *onyx*. When the opthalmia is more violent, it not unfrequently happens, that the quantity of matter deposited in the anterior chamber, is so much more considerable, as almost to fill it, and then it takes the name of *hypopion*. Indeed sometimes, the anterior chamber is filled with pus, which, going through the pupil, occupies also the posterior chamber, when, if not speedily absorbed, it becomes inspissated, and remains fixed, forming adhesions to the capsule of the chrystalline lens, or the *iris*, which prevents vision: or else, suppuration having previously commenced in the *cornea*, it gives way, and the matter is discharged through its aperture, which is afterwards succeeded by the

loss of the chrySTALLINE and vitreous humours causing the coats of the eye to collapse. But matter is more frequently deposited in the substance of the *cornea* itself, than in any other part of the eye. If the increased action and sensation are not very violent, the vessels in many parts of the *cornea* secrete a small quantity of *pus*, which is deposited at their extremities, and is not sufficient to produce a speck, but gives the eye a dull whitish muddy appearance which has been called *nebula*, or a cloud, the pupil is scarcely visible through it, and objects are not distinguished by the patient. The organization of the *cornea* is not much injured by it, as on the inflammation subsiding, the opacity is soon absorbed, and the sight returns. When this peculiar action of the vessels is confined to one part, and matter is formed between the *laminae* of the *cornea*, in such a quantity as to separate them for any space, a speck or film is produced; the degree of the ill effects of which will depend on its size and whether it

tuated near the external surface of the *cornea* or not. If the inflammation subsides before the matter discharges itself, it often is almost entirely absorbed, and, if on one side of the *cornea*, near the *sclerotica*, the sight is not in the least injured; but when it is on the centre, immediately before the pupil, as there is usually some slight opacity remaining, from its structure being injured, vision is not perfectly distinct. When the thickening of the *cornea* is towards the centre, and so great as to prevent the passage of the rays of light to the *retina*, the eye will frequently accommodate itself to its imperfections, by turning on one side, so that the transparent part of the *cornea* may be opposed to the object, and more perfect vision be obtained. I once saw a person, who had had an inflammation, succeeded by specks in both eyes in succession, some time intervening between each attack, and who had obtained such complete voluntary power over the muscles of the eyes, that, the opacity being so situated as on directing

the axis of both the eyes, to one object, figure was very imperfect, she could turn either or both the eyes to or from the object, or in contrary directions, as she pleased. When the collection of matter is superficial and breaks; the contents of the abscess are discharged and an ulcer is formed, which gives the appearance of a small hole or depression: if very small, it will sometimes fill up and disappear; but if more deep, it frequently leaves a flat opaque surface, which is a great impediment to the sight, causing objects to appear as if they were seen through glass, that had been made in water and whose surface is uneven. If the abscess should be very large and deeply situated in the substance of the *cornea*, and burst externally, such a considerable support frequently taken away from that part of the *cornea*, that the remaining portion is no longer able to support the action of the muscles, but gives way, and as, when the muscles act at the same time, they press the sides of the eye nearer together a

wards, a part of the *iris* is protruded through the wound in the *cornea*, forming a small black spot, which has been compared to the head of a common fly, and is spoken of by authors under the name of *pyocephalus*. If the protrusion is small, though the form of the pupil is changed to oblong, yet still the *iris* retains its motion and the sight is little injured: but if large, it loses its power of contraction, and dilation, and in some cases the edges, from being inflamed and remaining some time in contact, adhere, and the pupil is obliterated.


If the increased action of the vessels is considerable, yet not so great as to produce abscesses, which, on account of the structure of the part, cannot easily be formed, they go on to another kind of diseased secretion, and new vessels and parts are formed, or the old ones elongated. Hence, at times, we find an increase of cellular substance to the *connectiva*, with a congeries of new vessels,

which generally produce a prominent ridge running from one of the angles towards the centre of the *cornea*. At other times, when the inflammation has been violent and of some continuance, small vessels are to be seen on the external surface of the *cornea*; in general there are not many, and their course is from the edge of the *cornea* towards its centre, in right lines. Though commonly there are not more than from six to eight or ten of these vessels, in some instances the whole *cornea* is surrounded with the vessels which have the appearance of *radii*, converging to a centre; and, in some few cases when the inflammation has been rather severe, has several times a little subsided and has again returned in quick succession, the diseased action has gone so far as to permeate the whole body of the *cornea*, giving it the appearance of a highly vascular substance. To form these new vessels, it is requisite that the circulation be not too much hurried, otherwise, instead of an accretion of new parts, pus only is secreted. T

peculiar action of the vessels is not confined to the external surface of the eye alone. At times it affects the *iris*, and small spots or filaments are projected from its circular edge, which have somewhat the appearance of the *iris* in a horse. If they continue to be secreted, they extend themselves either till they unite with the projecting membrane on the other side, and thus form an opaque substance, which occupies the circular aperture of the *iris*, and prevents vision; or else, they unite to the capsule of the chrystalline lens, by which the motion of the *iris* is impeded, and the sight rendered very imperfect. It must be observed, however, that in this latter case, the vessels of the capsule of the chrystalline humour are previously affected with this kind of diseased action, which generally begins towards the centre, and if after the projecting fibres from the *iris* have continued some time, the eye is carefully examined, it will generally be found, there is a small opaque spot on the centre of the capsule.

In a few weeks, and sometimes even in few days, it will be seen, that, the *iris* has almost lost all motion, and that near edge on the capsule, there is a small circular clear ring, through which, what little sight the patient has, must be obtained, as the middle is occupied by the white opaque spot.

It sometimes though more rarely happens that the chrySTALLINE lens becomes opaque after a long and violent inflammation, more especially if it has been the consequence of a blow on the eye; and occasionally the action of the vessels in the internal part of the eye is so far diseased, as to change the structure of the vitreous humour or induce a paralysis of the *retina*.



The CAUSES of OPHTHALMY.

The same general causes, that excite inflammation in other parts of the body, will produce an ophthalmy; but, on account of the delicacy of the structure of the eye, and its exposed situation, it is liable to be affected by so many things which, applied to other parts, would be innoxious, that it is wonderful inflammations of the eyes are not more frequent than they are: add to which, as an organ of sense, it is stimulated in a manner peculiar to itself, by light and colours.

Every symptom and appearance of the disease, shew that the *impetus* of the blood to the vessels of the part is increased, and in course any thing that can bring on a longer and more frequent contraction of the fibres, may produce inflammation. This may be done, either by the application of those substances which stimulate the part

above the usual degree, causing an over exertion of the fibres ; or by diminishing for a time the usual *stimuli*, by which the irritable principle in the part will be accumulated, and the facility with which the fibres are excited into action increased; so that great exertions will be produced by the ordinary *stimuli*, or by those a little stronger than usual.

It is not however to be supposed, that a particular quantity of increased or diminished *stimulus* will produce an inflammation of the eyes : that will depend on the general state of the body, and the facility with which its irritable principle is brought into undue action, with the habitudes the fibres may have acquired of running into diseased associated motions, from a previous inflammation; as those who are in a convalescent state, from an intermittent fever, will experience a relapse, from the causes, which applied before the attack of the fever, would have been harmless:

those who have once been affected by *cynanche tonsillaris*, are very liable to a return of the same complaint.

But to know how, or in what manner inflammation is excited by the above means, it may perhaps be necessary to inquire into a few of the laws of life, which are connected with it.

All our motions, then, whether they are obedient to the will, or involuntary, together with the various orders of vessels, which perform the different functions of the animal body, as those of circulation, secretion, absorption, digestion, &c. whether irritative, or sensative, perform their motions from *stimuli* of various kinds, applied to the different parts; but to produce this action or motion, it is requisite that there be, as well as the *stimulus*, a certain quantity of the irritable principle, or of that something which constitutes life, in opposition to inert matter, in each fibre.

This principle has been generally supposed to depend on a something supplied by the brain, though some modern philosophers have attributed it to *oxygen*, obtained by respiration, from the atmosphere; and others have thought it might be explained on the supposition of the action of the electric fluid; but what or which ever may be, is in this place immaterial, I only wish to point out the effects of certain causes, indifferent circumstances, without venturing to inquire what these causes may be; and the facts being acknowledged whether the cause is considered, as the effect of the irritable principle, which may go under the name of vitality; or of irritability and sensibility, as called by Haller; or of excitability, as called by Brown; or the spirit of animation by Dr. Darwin; or of electricity by others, the facts, as abstracted from the cause, will be still the same.

As the health, and the regular performance of the different functions of the body depend upon a certain equalization of the *stimuli* applied and the quantity of the irritable principle, with the degree of tone existing in the moving fibre, when either of these are in a greater proportion than usual, the motions will become irregular and inordinate, and disease be the consequence. Thus inflammation may be induced, when the usual *stimuli* are increased or a new one superadded, the irritable principle being in the natural quantity, with proper tone of the fibre, as happens by the application of stimulant substances; or inflammation may be produced, when the *stimulus* is not greater than ordinary, but the irritable principle increased or accumulated, as takes place in going from a very dark to a light place, or from eating the usual quantity of food, after long abstinence.

Though it has been just remarked, that health consists in the equalization of the

stimulus and the irritable principle; y
 either or each of these may be increased
 diminished in a certain proportion, witho
 disease, commonly so called, being indu
 ed, but merely a tendency or predispositi
 to it, which is liable to show itself on a f
 farther increase of the *stimulus*, or of t
 irritable principle, which ever it happens
 be.

This may constitute the differences
 constitutions, or of what has been consider
 as temperaments. Thus in those peo
 whose supply of the irritable principle
 rather greater, than what health wo
 strictly require, great strength or tone
 the fibre is produced; the different functi
 are performed with great activity; the
 is hot and moist; the voluntary moti
 are executed with great facility and stren
 and their passions are strong; digestio
 carried on with vigour, and the powe
 assimilation is great, and the alvine eva
 tions are not frequent or large in quan

This is the most desirable state of existence, as the pleasurable sensations are strong, and the painful ones less frequent, than in any other: disease, also, though when it does come on, is generally violent and dangerous; yet it is difficult to be induced, as the system has, to some extent, the faculty of rendering harmless, those things that would prove noxious, to many other persons.

If a *stimulus* greater than natural is applied, unless it is attended with much pain, the parts are called into greater action, and the irritable principle is soon so much reduced as to put a stop to the undue exertions: or, being exposed to a *stimulus* less than natural, as cold; the great supply of the irritable principle, enables the body to prevent the accession of that torpid state of the vessels, which those of a contrary constitution experience; one feels comfortable and pleasant, whilst the other is shuddering with cold. We might carry the comparison still farther, and shew, that the

degree of cold that would destroy life in one, would be endured by the other with impunity.

On the contrary, in those whose irritability is barely sufficient to carry on the different functions of the body, pleasantly and regularly, as in those of a cool dry skin whose strength is exhausted by slight, short exertion; of moderate passions, and, the appetite is good, whose alvine evacuations are frequent, and the contrary state or costiveness, prejudicial, causing headaches and various unpleasant sensations; the system can but little accommodate itself to trifling changes of the usual *stimuli*; for the fibre, being in a state of atony or weakness and its mobility much increased, it is very susceptible of any change of the usual *stimuli*. If the body is exposed to a current air, which substracts the heat quicker than common, or to an atmosphere cooler than usual, a torpor is immediately induced, and with it, an accumulation of the irritability.

principle succeeded by irregular and diseased action: this by repetition is more easily brought on, and if the deficiency of *stimulus* should be greater than ordinary, and the torpor longer continued, inflammation will follow. Thus in those, who, in common language, are called scorbutic, whose hands, which are most exposed to heat of different degrees, are dry and cold, and whose skins are subject to crack, occasioned by a want of action, and atony in the cutaneous capillaries, from deficiency of the irritable principle; they regulate the external *stimulus* of heat, or so manage it, that the change shall not be very sudden, as, by wearing thick leather gloves, the skin of their hands will continue whole; but let them go from a warm room, into an air not so cold as to affect those of a contrary constitution, suppose at 45 or 50 degrees of Fahrenheit, the hands will become cold and pale, they will then look rough and dry, and will eventually crack. If one part of the hand more liable to torpor than the rest, which

often happens, and the cuticle is divided by putting a bit of oily adhesive plaister on it, which will confine the heat, and keep up the action of the vessels, by the next day the cracks will have disappeared and the skin be even softer and clearer, than that of the neighbouring parts.

With respect to the effects of a greater or less quantity of stimulus on the living fibre or the irritable principle, we cannot perhaps set the facts relating to it, in a clearer point of view, than by considering action, at different degrees of strength, a stimulus which is well known to excite one, and without whose presence, a certain proportion, life cannot exist: need I say, that I mean the *stimulus* caloric or heat.

When heat at a high temperature is applied to any part of the body, the *stimulus* is so very great, as to immediately extinguish the life of the part, after which

ganization is completely destroyed; for, it generally happens in atmospheric air, decomposition takes place, forming some elastic products, and a coaly or carbonaceous residuum. When the *stimulus* of heat has not been so great, or applied only for a short time, the life of the muscular extremities of the capillaries is destroyed, stronger and more frequent contractions of the text fibres in succession are produced, and these exciting pain, act as a new *stimulus*, which assists in increasing the action, and inflammation is the consequence,

If the quantity of heat is rather less, the organization of the capillaries is not destroyed, but the irritability is nearly exhausted, and the effects of the application of the caloric will depend on the method of treating the part. If nothing is done to prevent the ill consequence of the burn, the neighbouring fibres roused into exertion, not only by the stimulus of an extra quantity of heat, but also by the disagreeable

sensation arising from the injury done to the capillaries, will be impelled into more frequent contraction, which, producing pain, will increase of *stimulus*; and if not of the irritable principle, at least of the activity of it; and serum will be secreted, which will arise in the form of a blister. If, on the other hand, cold is applied, as by immersing the part in cold water, it will cause temporary ease, but the pain that follows the withdrawing of the part from the water will be more acute, for the coldness of the water prevents the fibres, already very much exhausted, from regaining their natural proportion of the irritable principle, whilst the neighbouring parts suffer an accumulation of it, and thus every symptom is aggravated. But, on the contrary, if immediately after a slight burn, warm alcohol or spirit of turpentine, (See Kentish on Burns,) be applied to the part, the first order of vessels, having been previously acted upon, by a much stronger *stimulus*, will be little affected, but the next order of vessels

will be excited into action, and their irritable principle diminished, when the pain will subside, and in a few hours afterwards, from the natural quantity of irritability being again accumulated, the healthy action of the part returns. When the quantity of heat applied is still less, the contraction of the vessels is increased, and the part becomes slightly red; but as the pain is not so great as to cause much sensation, the pain soon subsides, and the healthy motions return.

Our next degree of heat is what we are always exposed to, from the caloric in the atmosphere, under which, when moderate, all the functions of the body are pleasantly and regularly performed; but as it is liable to variations, being often in England, at one time of the year, sixty degrees hotter or colder than at another, according to the continuance of the greater, or less degree of heat than usual, or the rapidity of its changes, the body is liable to be affected in

different ways, and by diseases under different forms. Diseases too are more frequent when the daily vicissitudes of heat are considerable, as in hot days and cool evenings, or when the atmosphere is loaded with moisture, which, by causing the air to become a much better conductor of heat, substracts from our bodies quicker than ordinary, is clearly perceived in going from a well ventilated room, into which the sun has had free admission, or where there has lately been a fire, into one no colder, as indicated by the thermometer, but into which, the air, and the light have not been able to enter, and whose walls are damp, and the atmosphere of course, saturated with moisture.

Man is able to exist under very different degrees of heat; yet the parts of his body are so fashioned, as to render him unable to bear the fervid heat of the equatorial regions, with impunity: for heat, on his body being continually exposed to it, a

certain quantity, is inimical to the duration of life. It is difficult to judge what precise time, the omnipotent Director of the universe ordained, that man should remain in this little speck of the creation ; but certain it is, that life is protracted to the greatest length in the temperate zones.

It has already been said, that life consists in the application and presence of a certain quantity of *stimulus* and of the irritable principle, and it may here be added, that death will follow, when either of them are deficient in a certain degree. It may therefore be easily inferred, that the more the *stimulus*, or the irritable principle has been increased from birth, to carry on the animal functions, the sooner vitality will be extinguished. Thus between the tropics, the *stimulus* of the solar heat is the greatest, and the motions of the body are performed with more strength and celerity, in the early years of life; by the warmth of the climate liberty sooner arrives, succeeded by a pre-

mature and early old age, and in forty fifty years they die. In these, the fil having been previously much exerted fr the undue *stimulus*, more speedily lose power of being acted on by it, and anit tion becomes extinct.

This premature death is not induced caloric only, but by every other stimu The potation of much spirituous liqu the swallowing of large quantities opium, or great voluntary and contind exertions, will produce the same eff Hence the life of dram-drinkers is shor ed, of those of the Mahomedan faith, w being prohibited wine, make free opium, and of the hard working and l rious mechanic.

In this temperate climate, we all l re experienced the debilitating effects of a summer's day. The circulation is incre ed by it, both in frequency and force; the is hot and bathed in sweat, by the ; a

action of the capillaries, and, by noon, the whole body becomes languid and feeble, from the rapid expenditure of the irritable principle. On the contrary, the tonic power of cold, has been much spoken of, and highly commended by Physicians, but being only a negative quality, it acts, only, by its not allowing the irritable principle to be so quickly expended, and the moving fibre weakened.

The effects of a certain degree of cold will be different, on different constitutions : one, from a temporary diminution of the activity of the circulation, will feel invigorated; whilst another, from a torpor being induced, followed by an increase of the contraction of the fibres, stronger than their degree of vigour can endure, will be enfeebled and languid.

As we descend in the scale of heat and arrive at about the temperature at which water freezes, on the body being ex-

posed to its influence, a temporary quiescence of the capillaries of the skin brought on, and the pulsation of the arteries becomes slower; but when the muscular fibres have acquired a certain increase of the irritable principle, the diminished *stimulus* is sufficient to excite them into action, and if the quiescence has been great or long continued, strong contractions take place, as in the burning sensations we feel on coming into the house after riding horse back in frosty weather, or at handling snow, which causes what is called the hot-ache.

If on torpor being present, the usual quantity of *stimulus* or a greater one is applied, the exertions become so great, as to bring on inflammation, as in those whose fingers being benumbed with cold, they thrust them suddenly near the fire to warm them, which produces a *paronychia*: or in those who are first bitten, and have heat suddenly applied to their bodies. If

cold is still greater, such a degree of quiescence takes place, as to put a stop to all action, and a succeeding *stimulus* is applied in vain, and the death either of a part, or of the whole body follows.

We cannot carry our inquiries respecting the effects of a very great degree of cold, or rather of a very small quantity of heat, so far as we can of caloric at a high temperature: but we are told, that on some of the French Philosophers making experiments with ice and different mixtures to freeze Mercury, that, on one of them putting his finger for a few seconds, into the compound, which was some degrees below 0, it gave him an acute pain, as if pins, or sharp bodies were piercing his flesh.

I have been rather more prolix, in laying down the doctrine of the effect of the increase of *stimuli*, or of the irritable principle, than might perhaps, at the first view, appear necessary in a treatise on opthalmey:

but as my plan of cure will a good deal depend on what has been just said, I could not with propriety say less, to make my ideas on the subject sufficiently clear, and indeed, as we proceed, I shall have occasion to touch on it again.

Inflammations of the eyes then may be caused, either by an increase of the usual *stimuli*, or of a new one superadded;

Or, by an increase or accumulation of the irritable principle. But to be more particular:

First, Inflammation may be excited, by any thing which can stimulate the vessels of the eye into actions greater than nature as,

A. By violence done to the eye by mechanical means, which seem to produce their effect, from the form and solidity of the particles, destroying the continuity

the part, or from its motion being impeded, or by blows or wounds in the eye; warts, or encysted tumors, on the lids; an inversion of the ciliary edge of the eye lids; or by foreign bodies inserted between the lids and the globe.

B. By stimulant and acrid substances.

C. By those things which affect the eye, as an organ of sense.

D. By any thing that will increase the action of the lachrymal gland, which by causing a more copious secretion of tears, will add to the irritation, and particularly when they are confined, as in the measles, small pox, &c.

E. By the action of the vessels in the vicinity being increased, as by erysipelatous, and various inflammatory affections of the face; and wounds, or burns, of the same part.

2. By a *stimulus* less than natural, as cold, ‡ as by going from a warm room into cold air.

Having mentioned, in general terms, those things that will produce inflammation of the eyes, we shall proceed to speak of each in detail, and first:

1. Blows on the eye, according to the force with which they are given, or the greater or less mobility of the system, or of the part, may bring on very different degrees of inflammation. If slight, the effects are most commonly of short duration, but if violent, a confusion of the coats and inflammation takes place, which produces an incurable blindness.

‡ Though cold is mentioned here, as well as in other places, as a peculiar substance; the reader will remark that it is meant only to signify a smaller quantity of heat than is comfortable to the feelings, or the sudden subtraction of a portion of heat, when it is applied in a greater de-

After blows given by the hand closed, or by any obtuse body, though little inflammation would follow, and only slight injury seem externally to be done to the eye, yet in a few hours, perfect blindness may come on; but in general, it is after several days have elapsed. On inspecting the eye, it will usually be found that the *iris* is motionless, and dilated rather more than the sound eye, and that the chrySTALLINE humour is slightly opaque; yet in so small a degree as to render it little more than visible, and often pushed a little on one side. The lens generally continues getting more opaque, till it puts on the usual appearance of the cataract; and after some months, the motion of the *iris*, with the sensibility of the *retina* often returns. In these cases, blindness is caused in the first instance, by the insensibility of the optic nerve, as the slight muddiness of the lens, cannot sufficiently account for the want of vision: and afterwards by the opacity of the lens; for the violence of the blow, producing some

change in the small transparent vessels of the lens, they still carry on their diseased action, whilst the sensibility of the retina slowly returns.

A sudden blow, being given to the eye with a small blunt body, and probably in a peculiar direction, and with a certain force, will sometimes burst the capsule of the chrySTALLINE lens, and force the body of the lens, through the *iris*, into the anterior chamber. About five years since, a man was brought to me, who was struck on the eye, with a piece of steel, that flew off his tool, he was hammering. On inspection, I found, on the outer side of the eye, about midway between the circumference of the *cornea* and the outer angle, a small extravasation of blood on the *tunica conjunctiva*, which in other respects appeared to be a good deal injured by the blow, and the chrySTALLINE lens, was seen about half protruded through the aperture in the *iris* into the anterior chamber, of the aqueous humor.

hour; the *iris* also in one part was slightly lacerated. On the next day, the whole of the lens was discovered to be in the anterior chamber, and which became opaque three days after the accident. He recovered from the subsequent inflammation, as soon as could be reasonably expected, and though it is five years since, the size of the lens is not apparently diminished, and in that time, he has only had one attack of inflammation, so violent as to prevent him from attending to his usual employ, though it is frequently slightly red and irritable.

A few weeks since, also, a man wished my advice, for an inflammation, attended with a good deal of pain, which he had in one eye. On examining the eye, I saw, to my surprise, that the chrystalline lens, which appeared of the natural size, occupied almost the whole of the anterior chamber, yet not so much so, but that the *iris* could be perceived to have preserved its proper figure, and looking at the eye in

one direction, I thought I could perceive an opacity of the capsule of the lens. On my inquiring into the history of his complaint, he told me, that more than *two* years ago, as he was digging in a ditch, a twig struck him with considerable force on the eye; that he could see a little after the accident, but that by the next day the eye became very painful and swelled, and that on the inflammation subsiding he found he had totally lost the sight of the eye. In a few weeks, the eye became free from pain, and from that time to this last attack, which he attributes to cold, and which is a period of 22 years, he says it has not prevented him from attending to business; though he recollected, on my questioning him, that two or three times, the eye had been slightly red and troublesome, for a few days.

We learn from these cases, that the chrySTALLINE lens is not easily absorbed when deprived of its capsule; and that

may remain in the eye, and act as an ex-
traneous substance for many years, without
causing much irritation. Hence it is very
probable, that in depressing the cataract, the
lens is not absorbed, as is generally suppos-
ed, but lies buried in the vitreous humour,
and as it does not produce much irritation,
when situated in the anterior chamber of
the aqueous humour, where it must, in
some degree, press upon the irritable *iris*,
it is not likely to do so, when concealed in
the vitreous humour.

2. The danger attending wounds and
punctures, is still more formidable. A
sword or knife, dividing the coats of the
eye, causes an evacuation of the humours,
succeeded by total and irreparable blind-
ness. If the instrument is pushed with
much violence, it may pierce through the
globe, fracture the thin bones at the bottom
of the orbit, and penetrate the brain itself.
Such wounds are frequently followed, by
the most dreadful head-aches, large ab-

sceses following the inflammation, and the brain is compressed, immediate death

If the wound is inflicted by a needle or any small sharp instrument, the danger will depend on the depth and direction of the puncture. When the instrument merely pierced the *cornea*, without either touching the *iris* or the chrystalline lens, the wound usually heals in a few days with a moderate inflammation, leaving a small opacity, which when not towards the centre of the *cornea*, gives little inconvenience: but if, unfortunately, it penetrates, or even touches, the chrystalline lens, although at first, the sight should be much impaired, it is greatly to be feared that it will be affected in a week or two, and produce one species of cataract which will obstruct the rays of light in their passage to the retina, and prevent vision.

If, by the wound, the *iris* is divided or the *cornea* penetrated very near where it is

ned to the *tunica sclerotica*, part of the
is frequently protruded through the
erture, and forms one species of the
ophthalmoma. In that case, the figure of the
pupil is changed, from a circular to an oval,
or some other irregular form; but if the
protrusion is small, the part of the *iris*
which is not displaced preserves its power
of contracting and dilating, and the sight,
after the subsiding of the inflammation, is as
perfect as before. If the protrusion has been
greater, it forms a large irregular projecting
substance, which, by rubbing against the
cornea, adds much to the irritation, and in
consequence, the inflammation is much in-
creased, which frequently produces adhe-
sions of the *iris*, with loss of motion, a
thickening of the *cornea* or of the chrystalline
lens, that is succeeded by an incurable
blindness.

Wounds made through the *tunica sclero-*
tica and *choroides* must either enter the
chrystalline lens, causing an opacity of it,

or if it goes more inwards, will injure the *retina*, and produce a confirmed *gutta serena* with immobility and a dilation of the pupil. Wounds, therefore, that are made in the *sclerotica* are always hazardous, and the danger is increased, according to the magnitude of the solution of continuity, and the distance from the *cornea transparens*.

An extravasation of blood, under the *tunica conjunctiva*, is frequently produced by blows on the eye, as well as by punctures and small wounds, from some small artery or arteries being divided, which can pour out their contents externally. The quantity of blood thus extravasated is sometimes so considerable as to elevate the whole *conjunctiva* above the cornea in the same manner as in the *chemosis*, but in general the quantity is much smaller, only forming an irregular red spot or circle round the mouth of the vessel, and even if the eye has been otherwise injured, this red spot should be considerable, if

will be experienced, and the sight will not be at all affected. Absorption often goes on slowly, and the blood is several weeks in disappearing, to the great uneasiness of the patient; the *serum* is first absorbed, and the blood becomes darker coloured, and almost livid; it then gradually changes, and goes through all the intermediate shades of colour, from a very dark greenish yellow hue, and thence to its natural appearance.

Warts and encysted tumours of different kinds, situated on the lids, are frequently the cause of inflammation. Some people, more especially children, have repeated crops of warts on the lids, that are commonly situated towards the edge of them; and which, by their irritation, induce inflammation. The tumours that are met with on the lids are various in their size, and in the appearance of their contents, and on that account have had different names given to them by authors. They are at times of a

considerable size; and when they happen to be on the upper lid, by the impediment that arises to its free motion, and the pressure which they make on the eye, they are often a great source of inconvenience.

tumours for the most part are moveable, contain a matter of different degrees of consistence, from a glairy gelatinous likeness, somewhat similar to the white of an egg, to one as thick and as firm as full cream cheese.

Nearly allied to the encysted tumours above-mentioned are those small inflammatory tumours, which speedily terminate in suppuration: they are frequently formed towards the edge of the lids, and are commonly called Styes, Stians, or Pouks. These are often productive of a temporary inflammation; but as they usually suppurate in three or four days, it is seldom of any long continuance, unless the eye, from previous disease, is more inclined to run into disordered actions. When matter is fully formed

them, they may be opened with the point of a lancet ; or if the tumour should be situated on the edge of the lid, it will often be sufficient to extract two or three of the eyelashes, whose roots penetrating its body, on being removed, will afford an opportunity for the evacuation of the matter.

4. An inversion of the edges of the eyelids, called by authors *Trichiasis*, is at times another cause of Ophthalmia. This, from the continual irritation produced by the rubbing of the hairs against the eye, causes inflammation, which, if the inversion is not soon removed, will be productive of a good deal of mischief. The disorder generally comes on in a gradual manner, and unless, at its commencement, or joined with some other cause of inflammation, it gives no violent pain ; but an increasing irritation, with a constant flow of tears, on the motion of the lids. Many of the vessels of the *conjunctiva* are enlarged, and there is either an accretion of new

vessels on the *cornea*, or a general opacity it; and as both those diseased appearances are often absorbed, on the cause being removed, we sometimes see those that have been blind for months, on the inversion being cured, have in a short time recover their sight.

A præternatural or a double row of hairs sometimes formed, and again at other times either the hairs are placed irregularly to the ciliary edge, or there are only a supernumerary ones; these, by rubbing against the eye, cause inflammation in the same manner as the *trichiasis*.

5. Extraneous bodies inserted between the lids are not an unfrequent cause of opthelmy. The irritation they excite in the part in which they are fixed, and the hard and uneven surfaces, rubbing against the opposite part, on the least motion of the eye, or the lids, produce pain, with an inability to repeat the motion.

the lids, succeeded by a violent and dangerous inflammation, more especially if the extraneous bodies, besides the irritation from their bulk and hardness, possess any stimulant qualities, as lime.

6. There are a variety of substances, which stimulate the fibres into undue exertion exciting inflammation, whose effects cannot be accounted for, from the general laws of matter, as far as relate to their size, hardness or shape, but are owing to certain combinations, which produce such motions in the living system, that the knowledge of their component parts, would not lead us to suppose probable: these may be called chemical causes of inflammation, such are the application to the eye of essential oils, acids, or acid vapours, alkalies, and the different compounds, in which either an alkali or an acid is an ingredient, acrid juices, and that universal and diffused stimulant, caloric.

7. There are others which act on the eye as an organ of sense only, whose stimulant effects are peculiar to the *retina*; much light, particularly after the sensibility of the optic nerve is increased, from confinement in a dark place; exposing the eye for some time to snow, whose excitant power is much increased by a clear sky and sun shine; the eye being long fixed on vivid and bright colours.

8. Inflammation in many instances is brought on by the increase of the action of the lachrymal gland, causing a great flow of tears, whose saltness or acrimony seems to depend on the stronger exertion of the secreting vessels, as is seen in much weeping from mental affections.

The probability of inflammation being induced, is increased if, with the great flow of tears, the neighbouring parts are inflamed, as in the measles, when the redness of the eye-lids, and the parts adjoining,

ven the whole face are generally swelled
 and inflamed; and indeed inflammations
 of the eyes are very common, immediately
 previous to or during their eruption, which
 often continue for some time, causing small
 spots and opacities of the *cornea*. As the
 eruption of the measles disappear, the red-
 ness of the eyes generally subsides, though
 from the action of the vessels being increas-
 ed, during some days, and the mobility of
 the system being greater than usual, the
 vessels are extremely liable to run again into
 diseased motions on the application of a
 greater or less degree of heat, than common,
 and often, for months, the most trivial vari-
 ations of heat and cold, will excite a tem-
 porary inflammation, which at times is con-
 verted into a very obstinate and painful
 disorder.

The bad effects arising from an increased se-
 cretion of tears, will no doubt be much aggra-
 vated, if, with the inflammation of the lids,
 they should be so much swelled as to pre-

vent their being opened, which happens in the small pox. In that disease the eye-lids are frequently tumefied and firmly closed for several days, and from the confinement of the acrid tears, the most dreadful consequences follow. On the eye-lids being again separated, the *conjunctiva* is frequently found highly inflamed, and the *cornea transparens* ulcerated. If the *cornea* should be unaffected and the redness of the *conjunctiva* soon disappear, the eyes will some time be liable to a recurrence of inflammation, from trivial causes, the same as has been mentioned to happen, after measles.

It is generally supposed that those opacities of the *cornea*, which sometimes attend the small pox, are the consequence of a pustule or pustules being formed there; judging *à priori*, we might naturally conjecture, that when the face is entirely covered with the eruption, it would be probable the external surface of the eye would not escape.

very seldom, however, happens, that any pustule is found on the *tunica conjunctiva*; and though I have attended particularly to that subject, I have never been able to detect what I could call a small pock pustule on the *cornea*. It is true, on the opening of the eye there is seen, occasionally, on the *cornea*, a vesicle, filled with an opaque matter, which has a good deal the appearance of a small pock pustule, but instead of going off like the other pustules, it continues, and is absorbed the same as a common collection of matter from an ordinary inflammation; besides, it is never seen with the commencement of the eruption of the face, but only after the lids have been closed for several days, which makes it probable that it is the consequence of the confinement of the humid tears, with the consequent inflammation.

Whence is it that pustules are so seldom formed on the anterior surface of the eye? Do the vessels with difficulty run into

that kind of diseased action, from the peculiar structure of the part; or can the peculiar action of the vessels be prevented from the eye being always kept moist, and a considerable cooling process almost constantly going on, by the continual evaporation. We cannot easily inform ourselves of the heat of the eye, in its natural state, but in those animals who have a moist face, exposed to the air, as the noses of horses and dogs, on touching them, they generally give us the sensation of coldness. We know that the quantity of pustules produced depends in a great degree on the heat of the body, with respect to heat; hence the eruption is thickest on those parts which are covered with plasters, and the good effects of moderating the heat of the body is universally allowed.

Under the same head of an increased secretion of tears, or of their confinement to the eye as a cause of Ophthalmia, may be included the *fistula lachrymalis*. This

se, so named from a mistaken idea, that the ancient physicians had formed of its nature, cannot with propriety be called a *cul de lachrymalis*, unless in the very advanced stages of it. It is caused by an obstruction of the lachrymal canal, which conveys the tears from the eye to the nose, after having performed their office of moistening the eye, and by that means keeping the *cornea* transparent. On the passage of the tears into the nose being obstructed, the *cul de lachrymalis* becomes distended, and the tears, instead of being absorbed by the *lacrimal puncta*, remain in the eye, collected by the lower lid, till they are collected in such a quantity, as to run over the inferior edge, and down the cheek: these being a source of irritation by their long continuance in the eye, as well as by their stimulating effect on the lachrymal gland, causing a greater secretion of tears, excite inflammation.

9. Inflammation of the eyes may also be brought on by any thing that causes inflammation of the face, as various erysipelatous affections, particularly that species which is called St. Anthony's fire, or burns or wounds, in the vicinity of the

Having gone through the different causes which produce inflammation of the eye, by directly stimulating the vessels to a greater action, I shall proceed to the second division, in which, inflammation is induced by the subtraction of the usual *stimuli* for a time, and then from the sudden application of it again, in the natural quantity or perhaps rather greater than ordinary. This, by allowing an accumulation of the irritable principle, ultimately brings on inordinate and diseased action. The principle of irritability is generally diffused pretty equally through the fibres of our whole system, and, unless excited in some particular part, by great pleasure or pain, tends to an equilibrium; but in the

scles whose motions are involuntary, the supply of the irritable principle having been constant and perpetual from birth, their action being necessary to existence, though the usual *stimulus*, and, of course, their exertion is diminished, still they have their accustomed supply of this principle, which becomes accumulated, either, till the natural quantity of *stimulus* is re-applied, until the accumulation is so great as to render the part susceptible to be acted on by the smaller *stimulus*. (See Darwin's *zoonomia*).

The same laws with respect to the accumulation and exhaustion of the irritable principle, is observed in the inferior animals, and is extended also to the vegetable tribe. Allanzani remarks, that in the autumn bats bury themselves and become torpid; when the thermometer is at 54, and that, during the winter, they regain their irritability and re-appear in the spring, when the thermometer, is considerably below 54:

and, he observed, that when in spring exposed the vipers that were torpid to heat of 67 degrees only they died, though during summer, they bear a much greater degree of heat without the least injury. Vegetables sleep in winter, and are awakened by the vernal sun ; but die if a too powerful heat be suddenly applied, from their cumulation of irritability.

The voluntary muscles, or those of motion, have, from habit, acquired a facility by voluntariness, either of directing the irritable principle to or exciting it in certain fibres, in preference to others. Nevertheless, it is not to be supposed that it is ever passive state in them, as asserted by authors : though not supplied in that quantity as in the involuntary muscles, it is in constant action ; as is seen in the tremor of old people in the palsy of the head ; and from the traction of antagonist muscles, when which counterpoise them are cut through or lose their action, (called by Halle

insita), from a paralytic affection. It is probable that this voluntariness is a good deal acquired by habit, from the extreme awkwardness of an infant learning to walk, and muscles, from frequent use, acquiring strength, or more properly speaking, from obtaining the power of directing or exerting a large quantity of the irritable principle in a given time to certain muscular parts. It would appear, too, from the almost incessant motions of young animals as of children, who, unless asleep, are in almost constant action, or in the play of puppies and kittens, that in the earliest period of existence it was more constantly and uniformly diffused over the whole system.

Though an excess or defect of the accustomed *stimuli* produces a temporary quiescence of the part, yet as the causes are wholly different, so are their effects. In the first case the torpor is induced by the *stimulus* exciting the muscular fibres into greater action than natural, which expends the

irritable principle faster than it can be generated, and by that means diminishes the power of being excited, and renders the part incapable of carrying on its usual motions, unless by the application of a stronger excitation cause. On the withdrawing of the strong *stimulus*, the ordinary ones do not produce their accustomed effects till the irritable principle is again collected in its usual quantity; when the motions of the part proceed as before.

On the contrary, in the quiescence produced by the deficiency of *stimulus*, there is an accumulation of the irritable principle, which is called into exertion by a *stimulus* less than natural, which, once brought into action, the fibres are excited into such strong contractions as cause pain, which acts as a powerful *stimulus*, and if violent, or long continued, exhausts the irritable fibre, and induces torpor: the irritable principle is then gradually collected, the parts are ob-

their ordinary *stimuli*, and their motions performed as usual and health returns.

This desirable termination however does not always happen, for on the sudden application of the natural *stimulus*, when the irritable principle is much accumulated; as in those who have been long without food, or who have been exposed to much cold, inflammation will succeed: this will arise more easily, if, from previous disease, the fibres are easily excited into inordinate and irregular exertions from associated actions.

From what has been said, it may be inferred, that the irritable principle with respect to the moving fibre may be in three different states:

1st. When the expenditure by the action of *stimuli* is equal to its supply, it may then be said to be in a state of *Tone*.

2d. When the expenditure exceeds supply of this principle to the fibre, it may be said to be in a state of *Exhaustion*.

3d. But if the supply of the irritant principle given to the moving fibre, exceeds the expenditure, it may then be said to be in a state of *Accumulation*.

A deficiency of *stimulus*, is by far the most common cause of opthelmy. On the application of any stimulant substance, we perceive that it induces immediately a swelling of the danger, and we hasten to remove it; which being done, the very cause which induced the increased action, and which would continue it, is taken away, and the contractions directly begin to diminish.

If the heat of the atmosphere is increased, which is the *stimulus*, we are most exposed to, on its being again brought to its ordinary standard, no inflammation follows, but the force of the contractions gradually

ase and the healthy action returns. No additional thing can be applied to the eye, dependent of its relative degree of heat and cold, but what will act as a *stimulus*, and on that account, in considering those uses that will diminish action, we must totally exclude all substances, to which the eyes are not usually exposed.

The eyes are exposed to the *stimulus* of the ears, of the blood, of the air, of light, and heat. An alteration, in the usual secretion of the tears, may increase, but cannot diminish their stimulating effect; the blood, according to its quantity or quality, may have its exciting effects, either increased or diminished, but as that will act as a general use on the system, it cannot properly be introduced here; the atmospheric air also, it is supposed, from the experiments of Beddoes, to owe its stimulating power to the *oxygen*, that is one of its constituent parts, the proportion of which, is subject to very slight variations, cannot be thought

to differ much in its effects on the eye at different times, and the absence of light during the healthy state of the eye, does not appear to diminish the action of the vessels though in an inflamed or irritable state produces a considerable effect on the eye; there is only one cause then, to which we can attribute the diminution of the *stimulus* and the decrease of action, which is the loric.

The quantity of heat to which we are exposed is varying every minute, and we often change in a few hours, more than thirty degrees; this changeableness is greater, from the inventions of civil life, from hot rooms, and partial clothing, and the exposed and uncovered state of the eyes, whilst other parts of the body are defended by substances, which transmit heat slowly, make them more liable to be affected.

Whether heat is applied to the eye above
 under the usual degree, previous to in-
 flammation, a torpor will be induced which
 will be more likely to happen, when the
 change is sudden; as in going from a warm
 room into the night air, but as, at the time
 of freedom occasions any unpleasant symptom,
 it is disregarded till the subsequent reac-
 tion takes place.

It is in most instances there is a predis-
 position to the disease, and though the
 torpor at first may be small, and the re-
 action also not great, yet it may be sufficient
 to produce another state of quiescence,
 or more considerable than the first,
 which again produces an accumulation of
 the irritable principle, and the accumula-
 tion and exhaustion, continue alternately
 to come on, till a violent disease is formed.

In treating on the causes, which gene-
 rally produce Ophthalmies, it may perhaps
 be thought necessary, that something should

be said respecting *scrophula*, and *sy*, as they are considered, by medical writers on the subject, as often the causes of inflammation of the eyes. It is very common to see those, who have evident signs of a scrophulous habit, affected with Opthalmy; and when once it comes on, it is in general, not only a long time in subsiding, but very liable to return from small changes in the quantity of the organic *stimuli*, and particularly from variations in the atmosphere. I conceive that *scrophula* cannot be said with propriety to be the cause of Opthalmy, but merely renders a part of the system more susceptible to the action of any exciting causes, to which it is exposed, from its inability of resisting noxious powers, and its not being able to accommodate itself to small changes.

The *scrophula*, according to Dr. Cullen, depends on a *peculiar* constitution of the lymphatic system; that, however, it advances but little towards arriving at

urate idea of the disease. May we not conclude that this peculiar constitution is owing to a want of energy or activity in the whole system, but particularly of the absorbents? That it depends a good deal on a deficiency of vigour of the whole body, appears, from those causes that tend to produce it, having a debilitating effect; as lying in low, moist, and confined situations, inflammatory diseases, small pox, poor diet, want of pure air and exercise; and that general weakness and want of energy are present, is probable from the cold extremities, the frequency in the winter of chilblains on the hands, and kided heels; from injuries threatening the body being feebly resisted, and from the slow recovery of the constitution from disease. That there is a deficiency of irritability, or want of energy in the absorbents, in those that are predisposed to *scrophula*, appears from the moistness, from the difficulty with which ulcers heal, on account of the want of power of the lymphatics to absorb the fluid secreted

by the capillaries, causing the discharge of an acrimonious fluid.

Concluding, that in scrophulous H there is a deficiency of activity in the system in general, and of the absorbent in particular, it will easily be understood that if there should be an accumulation of an irritable principle, from defect of *stimulus* as from cold; on the increase of excitement succeeding to the torpor, the cutaneous absorbents may take up a greater quantity of fluid than usual, and this increased action not extending to the next order of absorbent vessels, beginning at the gland, they may be supplied with more than they can absorb, and an accumulation of fluid may be produced, causing an enlargement of the gland, and in the end, inflammation from the want of energy, too, the system rendered more susceptible of being affected by any irregularity in the usual *stimulus*; thus they are subject to *phthisis pulmonum*, *cynanche tonsillaris*, catarrh, and ophthalmia.

d in those disorders it will be recollected, that the parts affected are not only easily excited into action, but are also particularly exposed to the sudden changes in the heat of the medium that surrounds us.

When the ordinary *stimuli*, as heat and cold, food, &c. are uniformly and regularly applied, the different functions of the body are performed properly; and the secretions and absorptions are equal; but as from the deficiency of strength of the muscular fibres, a torpor of the extreme vessels is easily induced, their actions must frequently be irregular, and this often recurring, it is not wonderful that so delicate a part as the eye, and which is so much exposed, should in our earlier years be affected with inflammation, which, when once brought on, is, from association, much more likely to return on the same irregular action, again taking place. It will in general be found, that the more delicate and fair the skin is, the more susceptible they are of diseased motions.

On examining the inflamed eyes of those who have signs of *scrophula*, as fair skin swelled over-lips, and some of the glands of the neck enlarged, there is seldom a general increase of action of the vessels of the *conjunctiva*: there are vessels to be seen larger than natural over the whole surface, but the white colour of the *tunica albuginea*, is visible between them, near to its usual appearance; much light gives pain, causing a plentiful effusion of tears, and when the disease has continued for some time, the ciliary edges of the eye-lids are often affected; some of the hairs come out, and the small sebaceous glands, *glandulae Meibomii*, are inflamed, and at times ulcerated, causing the Psorophthalmy of Mr. Ware.

It has been supposed that these small glands are diseased on account of that tendency of enlargement, and inflammation of the glands in *scrophula*; but I think it may be better accounted for, from the great

bility of scrophulous habits, and the
 frequent and almost incessant discharge of
 acrid tears, over the edge of the eye-
 lids, which are often so acrimonious as to
 excoriate the cheek. When these glands
 have once acquired this diseased action, it
 is more easily repeated, and eventually
 slight causes will bring it on, and they
 may be a good deal inflamed when little
 increased action is to be seen in the eye.
 It would be paying an ill compliment to a
 fine complexion, to say, that all those who
 have fair skins, have a tendency to *scrophula*;
 I may, however, venture to assert, that the
 more delicate the skin of my patient, the
 more tedious is the cure of the Opthalmy,
 and the more liable it is to return from
 slight causes.

This rule, I need not say, will not hold
 good inversely, for there is another disease
 to which people with dark complexions are
 subject to, which will now be slightly
 touched on.

The disease alluded to, as far as I can recollect, has not been accurately defined by medical writers. In common and familiar language, it is called scorbute, though it does not seem to have any connection with the *scorbutus* of nosologists. It appears to be a debility, and want of action in the cutaneous vessels, and is marked by a dry and harsh skin, little or no colour in the cheeks, the cuticle by places coming off in powder, in scales; when the skin is abraded it takes a long time in healing, eruptions in different parts of the body are common, especially in those that are most exposed, as the face and hands, sometimes in small red pimples which suppurate, at others, small vesications filled with a transparent fluid arise, attended with an itching and burning; they break, and are succeeded by others, and in this manner it will occasionally continue for months, causing much pain and uneasiness. People of this constitution slowly recover from inflammations of the eyes, which is seldom attended with much pain, unless

for a few days at the commencement of the disorder; and though the vessels of the *conjunctiva* shall be a good deal enlarged, the light causes little inconvenience. A discharge of adhesive matter from the lids also, is frequent, part of the eye lashes fall off, and the glands ulcerate; indeed sometimes, after the inflammation has repeatedly recurred; it is entirely confined to the lids.

From the cutaneous vessels having less energy of action than usual, some degree of torpor will take place from a *stimulus* very little under the ordinary standard; and it is not uncommon for the patient, though he cannot recollect being exposed to either an unpleasant quantity of heat, or cold, to go to bed with the eyes and eyelids apparently well, and to awake in the morning with the lids inflamed, and firmly closed with an adhesive matter. On carefully cleaning the lids with a little warm water, by evening the inflammation frequently subsides, and the part affected

nearly recovers its natural appearance. In those cases, it will generally be found, that the patient, just before going to bed, has been either in a hot room or near the fire and has gone out of doors, and exposed himself to the cool night air, which has brought on torpor, and a succeeding reaction.

Opthalmies of the most violent kind have been said to arise from a *metastasis* of the gonorrhœal poison from the *urethra* to the *tunica conjunctiva*, or from carelessness and want of cleanliness, by the application of matter from the part affected to the eye, by means of the finger, or the handkerchief: latterly, however, the fact has been a good deal doubted, and I think with great reason.

None, perhaps, will deny the probability of the direct application of syphilitic, or gonorrhœal matter to the eye, not being able to produce a disease, which, from the

length of time that its removal would require, would most likely terminate in the loss of the sight; but what I wish here to inquire into, is, the possibility, or the degree of probability of the matter of *gonorrhœa* being translated from the penis to the eye, or of the capability of a person to affect himself, by matter secreted from his own person.

A *metastasis* in the *gonorrhœa*, often takes place, when the change is made; either to the *testis* producing *hernia humoralis*; to the neck of the bladder, where it is attended with what has been called the *Algedo*; between the *præputium* and *glans penis* causing a spurious kind of *gonorrhœa*; or to the groins exciting buboes. In these instances, the *metastasis* may happen, from a premature or sudden cessation of the primary discharge; but on that stopping, how the poison can so suddenly be conveyed to the eye, a part so distant, and whose action appears not to be con-

connected with the *urethra*, by any of the known laws of the animal economy, is what cannot easily be conceived.

It is true, Monsieur St. Yves says, that the venereal Opthalmy is very rare, but that he has seen several attacked with it, and that in most of them, the disease began two days after the commencement of virulent *gonorrhœa*, that the matter, not running off by the usual passages, was removed to the eye, through which it flowed, and stained the linen in the same manner as gonorrhœal matter usually does.

Astruc also gives a long and minute description of the venereal Opthalmy; but attending to his description, it will be found, that he had never seen the disease except in *one or two* of the patients, of the younger Saint Yves, who, he says, had the goodness to show them him, and he adds that the elder St. Yves was, he believes, the first that described the complaint. He does

t give so terrible an account of it, as
 veral later writers have done, neither does
 reckon it so certainly fatal. His de-
 scription of it is most minute; some parts of
 which are fanciful, and given apparently
 from some presupposed ideas he had form-
 ed of the disease: others answer tolerably
 well to the purulent eye. He says, that
 though the venereal Ophthalmia succeeds a
 suppressed gonorrhœa, it is *only* in those
 persons whose eyes are naturally weak and
 tender; or who have had them hurt, either
 by a blow or some extraneous matter, by
 which he thinks a free entrance is *opened*
 to the redundant venereal poison. The
 discharge from the eye, he likewise adds,
soon in a moment subsides, on the return of
 the original complaint.

Mr. Benjamin Bell, says but little about
 Ophthalmia, from a suppressed gonorrhœa,
 but acknowledges that it sometimes hap-
 pens, and mentions the case of a young
 man, "in whom the purulent discharge

“ from the eyes appeared eight or ten da
“ after the commencement of a mild
“ *norrhæa*, which came on after being he
“ ed with drinking port wine, when t
“ discharge from the *urethra*, which h
“ previously been copious, disappear
“ almost entirely. By a quantity of ble
“ being taken from the temporal artery
“ one side; from the division of such vesse
“ on the eye balls as were turgid, a
“ the scarification of the inflamed parts
“ the eye-lids; by the application over
“ eyes of poultices in which opium, a
“ *cerussa acetata* were dissolved, and gen
“ laxatives, the pain, inflammation, a
“ discharge of matter soon lessened, a
“ in the course of a fortnight, no symp
“ of the disease remained, but a degree
“ irritability on exposure to much lig
“ with which both eyes continued to
“ distressed for five or six weeks at
“ wards.”

When a person has a *gonorrhœa*, on the application of a greater quantity of *stimulus*, as the drinking of much wine, as in the case of the young man mentioned by Mr. Bell, the inflammation would be increased, but no suppression of the discharge would take place, unless on exposure to cold, which at the same time would be liable to bring on torpor, and a succeeding inflammation of the eyes, more especially if they were already weak; which makes me suppose that the same imprudent exposure to cold, that caused the diminution of the discharge from the *urethra*, for it was suppressed, produced the inflammation and discharge from the eye. This supposition is farther strengthened, from Mr. Bell proceeding to say, “ that in the course of the following year, on the same person being attacked with a *gonorrhœa*, but of a more violent nature than the former, he was again seized *after exposure to much cold*, and riding on horse back, to a similar affection of his eyes.” In

this instance too, blood letting, and the remedies formerly prescribed, proved successful.

Mr. Ware, in his “ Remarks on the Ophthalmia, &c.” takes notice of the observation of Mr. St. Yves, upon which he makes the following strong remark ; “ this account “ is the more surprising, because such “ effect as is here described, has never been “ observed by other writers on the subject “ or any one of the faculty, with whom “ I am acquainted ; though some of them “ have had a long and extensive practice “ both in the Ophthalmia and gonorrhœa, and the same, I believe, will be confessed by every medical man of experience : observation ; and I trust, we shall not be going too far, in declaring, that the cases which have been said to be owing to a suppressed gonorrhœa, have been caused by cold ; Astruc, we find, is of opinion that it only happens in those persons who have naturally weak eyes, or in

ws, and foreign bodies inserted between lids, and Mr. Foote, in his treatise on venereal disease, says, that he does not live in this doctrine of *metastatis*.

There is another method by which it is said, the eye may be affected with a *rrhæa*, and that is, from the patient conveying, by his fingers, or any other means, the matter from the *penis* to the eye, which it is asserted, will bring on an ophthalmia, accompanied with a discharge, similar to that which had previously taken place from the *urethra*: but however probable that may appear in theory, it would be extremely difficult to prove, that it has been proved in one instance, by experience. If it is possible for the eye to be affected in that manner, other parts would alike be exposed to its baneful action, and what a dreadful source of misery would be let in upon mankind?

Fortunately, however for us, it appears that we are incapable of infecting ourselves by our own diseased secretions, which are caused by the direct application of poison. On the application of any specific poison, as the gonorrhœal, to our body, a certain diseased action is brought on, a *gonorrhœa* produced; the matter which, is capable of exciting a fatal disease in another person, though unable to contaminate the body that produced it. It was not for that law of the animal economy that every infectious disease must be fatal. Matter from a chancre, would cause other chancres, and those again, other till the whole body was covered, and death closed the scene. The same may be said of *gonorrhœa*; and other diseases which owe their action to a peculiar poison. The poison of the viper, Mr. Foot says, does not affect the viper that secretes it, though it will another viper, or any other animal into which it is effused. It has been before remarked, that syphilis

gonorrhœal matter being applied to the eye, would produce its baneful effects on it; that however is a disease which is not here alluded to, but the secondary one, which is the same that has been meant by writers on the subject.

Mr. Foote in denying the possibility of matter from the same person, infecting him at a different part, gives two cases in which gonorrhœal Ophthalmia appeared about the same time as the diseased action shewed itself in the *urethra*. In both of which, it was very probable, that the matter from the infected person, was applied to the eyes. In one of the cases, the patient, who had weak eyes, was in the habit of washing them with his own urine, and which he did, half an hour, after he had had connection with the woman who gave him the gonorrhœa.

Mr. Ware seems to think it very doubtful, if a person can be affected with a dis-

case of the eyes, from the application of own gonorrhœal matter, and though should not be authorised to put the matter to the test of experiment, yet I appeal to every one who has had much opportunity of observation in these matters, whether he has not seen many patients who, to a want of common cleanliness, have added a greater degree of inattention, to prevent any effects that might arise from the application of the poison to other parts of the body, when the discharge has been copious, he not only handled the parts a good deal but even wiped off the matter with his pocket handkerchiefs, and yet the eyes as well as every other part, have not been affected. Whatever part is in pain, attracts most our attention, and we naturally lay our hand to it, and this hand is often applied to the eyes, the nose, and the mouth; for then, if it were possible to infect the eye, that kind of Ophthalmia would be very common, and yet I have never met with it. I confess, I have repeatedly attended

who, at the same time, laboured under *gonorrhæa*, and had a profuse discharge of matter, similar to *pus*, from the eyes, and which the patients have attributed to the *gonorrhæa*, but which clearly was the common purulent eye, which if it happens to come on when the *gonorrhæa* is present, the discharge being similar, has been supposed, without much examination, to proceed from the same cause, and has probably given rise to the idea of the gonorrhœal ophthalmia.

Though the purulent discharge from the eye is most common to children, yet it is not unfrequently is met with in adults, where there has not been the least suspicion of a venereal taint; and I have seen it extremely copious, and as far as the external appearance of colour, consistence, and staining the linen, exactly similar to gonorrhœal matter, and sometimes attended with great deal of pain. Had this happened with a *gonorrhæa*, what would have been the general conclusion?

It has been supposed too, that inflammations of the eyes may be brought on the system being universally infected with *syphilis*, though it does not appear, writers on the venereal disease, that it has any particular action on the eyes; and the Ophthalmia has been observed to take place when mercury has been given rather largely, to correct or remove the original syphilitic taint, and which of course would render the constitution peculiarly liable to inflammation, from the slightest concurrence of exciting causes. If it is necessary to give this disease any name, and allowable to seek one from a predisposing cause, it would perhaps, accord more with its nature to call it the Mercurial Ophthalmia.

It has been also said, that Ophthalmies are at times epidemical and contagious, and that they depend on some peculiar property in the surrounding atmosphere. If by some peculiar property in the atmosphere is meant its relative qualities with respect to heat

and cold, moisture and dryness, &c. which may bring the body into such a state as to make it much more susceptible of taking on diseased action, from any irregularity in the application of the usual *stimuli* inducing inflammation, I have no objection to the term; but if, by it, we are to understand that, at times it is probable the air may contain certain opthalmic particles, if it may be allowed the expression, which, when applied to the body, will produce a specific disease, as variolous effluvia causes smallpox, I must confess it appears to me highly improbable.


Inflammations of the eyes sometimes come very suddenly, and apparently without any sufficient preceding cause; this, by common people, is called a *blast*, and is by them been referred to something noxious in the air, and connected confusedly, in their minds, with what they have observed to happen to trees. It is, likewise, no unusual thing to see several people in one

family, affected at the same time, with Ophthalmies: this happens generally among the poorer classes, and commonly either in the spring or autumn, when the state of the atmosphere is particularly variable, and changes from heat to cold are daily, often sudden.

When the Ophthalmia is more frequent than usual in any particular district, it is generally attended, more or less, with a purulent discharge similar to *pus*, from the eye, which probably has some effect in spreading the disease. Though nearly the whole of a family may be affected at the same time, it should be recollected, that they are all exposed to the same exciting causes, and that among the poorer classes especially they are not sufficiently careful to dry themselves after washing, with different cloths, and cases have come under my observation in which it was very probable, that the purulent discharge, which is at times very acrid as to excoriate the skin in the vic-


of the eye, applied to another person's eye, produced inflammation, probably from its irritating quality: for it was not always succeeded with a similar discharge.

It has been asserted, by some able physiologists, that the vessels being thrown into such a diseased action, as to produce various or rubiulous matter, or *even common* *sores*; that this matter applied for the first time, to any part denuded of its cuticle, or where it is very thin and sensible, a similar action will be induced in those vessels, and the same kind of matter secreted: but that, when they have already gone into that kind of action, they are not again susceptible of it: when the matter acts as an irritating substance, and a slight inflammation only is induced. Granting that to be the fact, will it not count for the application of matter from one person's eye to another, sometimes producing a discharge of purulent matter, and another only a slight inflammation?





THE CURE.



HAVING now, rather in detail, mentioned the occasional remote causes of Ophthalmy, it may be requisite to say a few words, on the proximate cause, or that peculiar action of the vessels, which constitutes inflammation. It is not my intention, to discuss the improbability of the multifarious causes, which have been supposed to be the immediate occasion of inflammation, or to point out the strong objections, that might be made to many of them; that belonging more properly to a treatise, written professedly on the subject of general inflammation.

It will be sufficient for our purpose, to say, what, on taking a view of the various causes, which have been said to bring on

inflammation of the eyes, appears the most probable, and which, I trust, will lead us to a more rational plan of cure.

Were I to examine the whole body, I should discover that part, which was the most proper, from the facility with which a variety of extraneous bodies act on it, and the quickness with which any change, in its circulation, is shewn, I could not fix on one more adapted to show the nature of inflammation, than that to which these pages are devoted, the eyes; from which I will take an example of the effects, and action of a body, whose power is owing to qualities, about the nature of which there is little doubt.

A particle of sand being inserted between the lids, and the globe of the eye, by its mechanical irritation, stimulates the vessels, or the fibres against which it is immediately applied, into more frequent and stronger contractions, and a slight

ness is immediately visible; by association, the lids are forcibly closed; the contents of the excretory duct of the lachrymal gland, is pressed out; and the eye suffused with tears.

If by the flowing of the tears, the sand removed, the irritable fibre being stimulated not more than usual, the increase of action of the vessels subside, and the circulation is carried on as before.

But, if the irritating substance still remains, the contractions of the fibres increase in frequency, and force, the vessels, joining to those first affected, are brought to action, by association, and these in cause those that are nearest to them, to contract, till the parts are affected for the distance from the fibres, first stimulated.

Now, supposing one of the circular fibres or an artery to contract, with a greater

energy than usual, which will suddenly bring the sides of the vessel nearer to each other, and diminish its diameter, it will follow, that there will be a stronger *impetus*, and a greater quantity of blood propelled against the next circular fibre than before, which, at the same time it enlarges its diameter, will stimulate it to contract more violently: the same reasoning will hold good with respect to the adjoining fibre, by which the resistance of the minute vessels, which commonly admit only the transparent part of the blood, will be overcome, and red globules admitted: these and the increased distention will act as strong stimulants and produce heat, which will still add to the irritation.

If the increased action and pain continue for some time, though the original cause of the disease be removed, the inflammation will remain; for the activity of the irritating principle in the part being greater, the vessels acquire new habitudes and new actions which keep up the diseased motions.

When the inflammation is only several days in subsiding, after the removal of the exciting cause, as it is continued by the pain which arises from the enlarged diameters of the vessels, and the subsequent force of the contractions, on the pain remitting, the inflammation diminishes, and the action of the vessels partially return to their ordinary state, and, after a few remissions and exacerbations, health is restored.

That the enlargement of the vessels, in what is called active inflammation, depends on the resistance from the irritable principle present in the arterial fibres being overcome by the *impetus* of the blood, appears from the experiments of Dr. Buvina, who found that, on injecting fresh blood into the blood-vessels of living animals, the blood only penetrated into those vessels in which red blood circulates during life, but having suddenly killed the animal, by dividing the spinal marrow, the injections immediately penetrated to the most delicate vessels of

the *periosteum*, and other parts, giving them a red colour, which they have not in the living animal.

Since a variety of substances produce the same effect on the eye, as the grain of sand just described; but which, from their form or appearance, do not indicate that quality we are, perhaps, authorised in concluding they owe their power also to their stimulating nature. The first we may call mechanical *stimulus*, the latter a chemical one.

The chemical *stimulus* from the full manner in which it operates, and its being more diffused, and applied to many fibres at the same time, is more apt to produce violent disease.

As the increased action of the vessels is the very essence of inflammation; though the pain, and the greater heat when present, add much to the irritation.

they are only the consequence of, and
 abundant on, the increased action; to re-
 ve which, our principle attention ought
 e directed.

It has already been shown, that a
 ter action than ordinary in a part, must
 end either on the usual *stimuli*, or the
 able principle being increased; it will
 before necessarily follow, that to di-
 sh the action, the *stimuli* or the irrit-
 principle must be lessened.

The common external *stimuli* may be
 eased, or a new one added; or the
 ons of the part, may, by the pain,
 the distention produced, act as
stimuli, or the new motions may in-
 se the quantity of the supply, or the
 ity of the irritable principle.

It is universally agreed, that, when the
 mon *stimuli* are increased, they should
 moderated; or if any other is added,

that its removal should first be effected before any thing can successfully be applied, to the eye, to take away the inflammation. I shall proceed, then, to point out the most eligible means of removing or obviating, the ill effects of the exciting causes.

1. **BLOWS.** On a violent blow being given to the eye, it is to be feared that derangement of the internal parts will be produced, which may terminate in the loss of sight; our care, however, in the first instance, is to moderate the inflammation, to do which, it will perhaps be more necessary to remark, what ought to be done, than to point out what should be done, as, I fear, our extreme officiousness, has often been the cause of more mischief, than the original injury it was intended to cure.

By a blow, the healthy actions of a part are deranged, to restore which, is out of the power of art; it must be the work of

re, and on the first infliction of the injury, she is generally, more inclined to do too much, than too little. The common method of applying bandages, and compresses on the eye, with the intention of keeping out the light, or preventing the motion of the lids, ought not to be adopted: we also to prevent, or moderate the inflammation, use poultices of various kinds, cloths wet with different articles, and laid on the eye, all of which are highly improper and injudicious; for by confining the tears, and increasing the heat of the part, as well as by their pressure, they add much to the inflammation.

Practitioners very frequently order linen cloths, wet with different kinds of lotions, to be kept constantly applied to the eye, both after blows and wounds, as well as in common inflammations; and this they do, not so much, perhaps, on the supposition that their lotions have any specific power in abating inflammation, as to keep

the eye more cool, upon the principle that water is a better conducting *medium* for heat, than air, and that the evaporation that takes place from the wet cloth, assists in subtracting the heat.

This mode of reasoning is fallacious; although it is well known, that a certain quantity of water will transmit heat quicker, than a given quantity of air, and that the evaporation which is always going on from water, makes it in summer several degrees cooler than the air, and that even a bottle, containing liquor, and wrapped in a wet cloth, becomes cooler, as is well known in hot climates, where they cool their wine by that process; yet the analogy will not hold good when applied to an animal body.

In one case there is a certain quantity of heat contained in a body, which is at a temperature nearly equal to its own, which is diminished by the evaporation

the water; in the other, the wet cloth is applied to a body, at a much higher temperature, and whose heat is being perpetually renewed.

A wet cloth being placed in contact with an inflamed part, in two or three minutes, becomes the heat of the skin, and the external part of the cloth, which will suppose to be four or six folds, will be found, on trying it with a thermometer, both before, and after the application of the cloth, to have sunk only *two degrees*, and even this small diminution of heat, must not be attributed to the skin being cooler, but to the heat substracted from the external surface of the cloth by evaporation, as we find that in warm weather, water is that much cooler than the air.

The application, then, of a wet cloth, does not diminish the heat of the skin, but the part of the cloth in contact with the body soon becomes dry and hard, and acts

as a stimulating substance, independent of its weight, and every one who is at conversant with diseases of the eyes, may have often observed how small a pressure on the lids, is productive of evident irritation, and an increase of the inflammation.

If a thin piece of linen cloth, was always to be kept uniformly wet, by a regular supply of moisture, which would be attended with a great deal of trouble and inconvenience it probably might be practicable; but the application of a cloth to the eye, which is renewed every two or three hours, which is the usual mode, is undoubtedly prejudicial.

Little more can be done directly against a blow on the eye, than keeping the patient cool, putting him on an antiphlogistic plan, proportioned to his habit of body, the nature of the symptoms, and desiring irritating causes to be avoided. The eye should be bathed six or eight times a day, and

keewarm † water, and kept shaded from the light, particularly if it gives pain.

I mention warm water, to be made use of, to wash any irritating substance, whether it is a foreign body, matter secreted by the eye, which is often found in small vessels, and occasions a good deal of pain; and when I come to speak of the cure of the Opthalmy, in the second part, I shall make it appear, that the application of warm water is much more beneficial than cold. This assertion, however, is only to be considered as applicable to the intermediate stage of an Opthalmy; for after it has put on the atonic stage, cold water is often most proper. How difficult is it to get better of preconceived opinions, particularly when supported by something like reason! To superficial observers, inconsistency appears so evident. If a part is hotter than usual, say they, should we not endeavour to cool it, by applying cold water, and how absurd it would be, to use warm water when the part is already too hot!

Although they should find, when the sensation of coldness from the water was gone off, that the part was more comfortable than before; or if they should be prevailed upon to make trial of warm water under a disguised name, and call it *fomentation*, and should find ease; that the *stimulus* of the heat from the water could be serviceable to them, would be farthest from their thoughts.

I claim not this as a new practice, though I hope to extend its application, by pointing out in a more decided manner than has hitherto been done, the utility of it. A Benedict Duddell, who styled, himself surgeon and

In almost all cases of inflamed eyes, for advantage will be received from the use of shade, which will be found to do as much good, as bandages and poultices often harm. With respect to the kind of shade that is most proper, I would recommend a paste-board hood, to be worn, at a greater or less distance from the eyes, as the particular case may require. The hood should be made of a size, not larger than will keep off the necessary degree of light, and will be better if lined with black, instead of green, silk, as is commonly done.

An oculist, and who published in 1729, "A treatise on the horny-coat of the eye, &c." almost universally recommends the application of warm water, in diseases of the eye, which he orders to be used *incessantly*, for twelve hours together, and, as he says, with great success. Mr. Ware, in "Additional Remarks on the Ophthalmy," in his "Remarks on the *fistula lachrymalis*," says, that in some cases of Ophthalmy on applying "hot water to the eye, the eye was highly grateful, both *cooling* and *strengthening* the eye to which it was applied." He attributes its good effect unphilosophically, to the cooling process that takes place in the evaporation of the water;—but I must defer entering farther into this question, till I come to speak of its operation in abating inflammation.

It should be recollected, that this hood only intended as a shade, and should not press the head too much; to overheat it, or come very far over the face, to exclude the cool air. If a small bit of cane or whalebone, of such a length, as to go almost round the head, is firmly sewed to the inside of the upper part of the shade, and then tied behind, with a ribbon, to the side of the head, the lower part will be thrown out from the eye, and from the slightness of its pressure; the facility with which it is raised higher or lower; and the steadiness with which it preserves its situation, admit most any quantity of light may be admitted, according to circumstances.

The eye-lids and the adjoining parts are frequently discoloured after blows, and as the appearance is very unpleasant, its speedy removal is very much desired.

A great many things are thought, by the common people, to have the power of re-

moving this appearance in a few hours; a raw meat bruised and laid on the part Solomon's seal, conserve of roses, &c. but the powers of animal life are not so easily or quickly excited to do what is wished of them; the change must be gradual, and time given for the absorption of the blood though it may be expedited, by the frequent application of any gentle stimulant, as equal parts of alcohol and water, with a little camphor, tincture of opium, or æther diluted.

2. WOUNDS:—On any instrument substance penetrating the eye, nearly the same mode of treatment should be adopted as in blows. If the *tunica cornea*, or *sclerica* is divided to any extent, the sight will be very much endangered, and as the preservation of the eye, if it is possible to preserve it, will depend on the wound healing speedily, the greatest care should be taken that the curative process of nature be not interrupted by the circulation in the part being hurried.

The patient must be requested to keep the eye-lids closed, and should be confined to a cool, dark room, with nothing at all over or upon his eyes ; but if the lids are inclined to adhere together, and confine the tears, their edges may be kept constantly moist, by the application of any mild ointment ; though it generally answers the purpose better, if the ointment is made solely of animal fat : costiveness should be obviated ; febrile symptoms moderated ; the antiphlogistic plan attended to ; and any matter that may be formed gently and carefully removed, by bathing the eye with a little warm water.

If bandages, &c. are prejudicial after blows, they are much more so after wounds of the eye, though their application to guard the eye from the light, are, in general, so much more convenient to the friends of the patient, than confinement in a dark room, that they are too often had recourse to.

When the wound is healed, and an inflammation remains, it must be considered as an inflammation arising from any other cause, whose original exciting cause has been removed, and treated accordingly.*

For the extravasation of blood, under the *tunica conjunctiva*, which is the consequence of the division of a blood-vessel, the application of the vapour of æther, on account of its penetrating quality, has been much extolled. Probably this penetrating quality of æther is merely imaginary, it not being able to pass the external membrane, and that the good effects, resulting from its use, ought to

* How much more rational is the present mode of treating wounds of the eye, to what it was formerly! when, after depression or the extraction of the cataract, the unfortunate patient was confined to his bed, his eyes bathed with a variety of *powerful specifics*, and covered with a load of bandages three or four weeks! When the bandages were removed that his best friend, cool air, might be admitted to him; to moderate the inflammation, alas! it was then often too late, and the miserable victim, sacrificed to mistaken principles, was doomed to darkness.

attributed to its stimulant powers ; as I
have found, the dropping a little spirit and
water into the eye answer equally as well.

In speaking of the subsequent ill effects
wounds or blows upon the eye ; an opa-
city of the chrystalline lens, was mentioned
an occurrence not infrequent : these kind
cataracts, as well as many others, are
generally thought incurable, and the whole
dependence of both the patient and the prac-
titioner, has been either the depression or
extraction of the cataract. In many cases,
I have been so fortunate, as as to impede or
stop the progress of the disorder, and in
a few, to remove it, when fully formed.

In the succeeding part, when I come to
speak on the consequences of inflammation
of the eyes, I intend laying before the pub-
lic, my mode of treatment, in opacities of
the chrystalline lens, with some cases in
which it has been attended with success :
it suffices at present to say, that I chiefly

depend on the application of a variety of stimulants to the eye, of such a strength and repeated as often, as may be sufficient to keep up a moderate, but perpetual, irritation.

3. Warts and tumours on the lids are mentioned as another cause of inflammation. Warts, after arriving at a certain size, generally suppurate and disappear, but if they are productive of pain, they should be removed.

If the warts are attached to the lids by a small pedicle, they may be cut off with a thread, or more expeditiously by a pair of scissors, or a scalpel.

It is those, however, in general, with a large broad base, that we are consulted about. The best way to remove them, is to cut them off even with the skin, with a fine scalpel, and then touch the part with caustic; a small slough will be formed, and

=====

operation that succeeds will take away
the remainder of the wart.

In removing tumours of the encysted
kind, from the lids, authors commonly
commend the skin to be divided, and the
tumour to be dissected out entire: this, if it
is situated on the upper lid, is often a trou-
blesome and painful operation, and I think
is attended with unnecessary violence;
at least in a great many that I have remov-
ed, I have always found it sufficient, merely
to lay open the cyst, by cutting into it
through its whole extent, and after evacuat-
ing its contents, to bring the edges of the
wound together by small flips of adhesive
plaster, when in a day or two it generally
heals by the first intention.

4. The mode of treatment, with the rea-
soning on the nature and cure of the Op-
thalmia, arising from an inversion of the
lids, is so accurately given by the inge-
nious Mr. Ware, in his Remarks on the

Opthalmey, and so perfectly accords with n
own experience, that I shall beg leave
give it in his own words.

“ For an Opthalmey, produced by an in
“ version of the lids, a palliative cure m
“ be effected, or to speak more conformal
“ to the fact, a present and temporary rel
“ may be given to the patient, by taki
“ out the lashes, with a forceps, a pair
“ knippers, or any instrument of the li
“ kind. But while the lids retain this
“ verted state, no sooner do the hairs gr
“ again, than the disorder will again retu
“ nor can the patient be ever properly f
“ to be cured of the complaint, till
“ edges of the lids are restored to their
“ tural position, and can be kept in it.”

“ It is, however, necessary, that a
“ tinction be made between an inversio
“ the upper and lower lid, for though
“ inversion of either will produce the s
“ effect, yet in the different lids it app

to arise from different causes, and consequently to require different methods of cure."

"The upper lid, and its ciliary edge, both in motion and at rest, are preserved in their natural situation, by the equal, though contrary actions of the *musculus orbicularis*, and *Levator palpebræ superioris*. But the lower lid, whose motion is very small in comparison with that of the former, has no muscle correspondent to the elevator of the upper; and is preserved in its natural state, by the equal action of the orbicular fibres spread over it, and the counteraction of the skin which covers it; in which last respect it differs materially from the upper lid, the skin of which, on the contrary, being always very thin and flaccid, is incapable of any such counteracting power."

"From the above account it is manifest, that when the *trichiasis* affects the upper

“ lid, it must be owing to a relaxation
 “ the *elevator palpebræ superioris*, and
 “ contraction of the superior part of the
 “ *orbicularis*; whereas in the case of
 “ *trichiasis* affecting the lower lid, it can
 “ only arise from a relaxation of the skin
 “ and a contraction of part of the *orbic*
 “ *laris*. And as in these two cases, the
 “ causes of the disorder are very different
 “ so they will of consequence, require
 “ very different treatment. In the *trichiasis*
 “ of the lower lid, it will be necessary
 “ to increase the counteracting power
 “ the skin, which covers that lid, so as to
 “ prevent the undue contraction of the *m*
 “ *culus orbicularis*: whereas in the *trichiasis*
 “ of the upper lid, it is plain that
 “ sole object of attention, must be to give
 “ an additional *stimulus*, to the *Lev*
 “ *palpebræ superioris*, for the purpose
 “ exciting it to action.”

The *trichiasis* of the upper lid happens
 but seldom. But in an instance of

and, that did occur, an entire cure was produced by a friend of Mr. Ware's, on a young gentleman 18 years of age, by making an incision through the integuments of the upper lid, from the inner angle of the eye to the outer, and then separating the fibres of the *orbicularis*, so as to denude the expanded fibres of the elevator muscle, as near to their termination in the edge of the lid as possible, and applying a small cauterizing iron, adapted to the convexity of the globe of the eye, and made pretty warm, to the tendino-carnous fibres.

The thought was certainly ingenious, and answered the purpose of curing the eversion of the ciliary edge; at the same time, producing a slight eversion of the lid, which though its appearance is unpleasant, causes none of the disagreeable effects of the other. The gentleman says, the iron was made pretty warm, and that he passed it two or three times over the fibres. I

conclude, therefore, that the iron did not as was supposed act as a *stimulus*, exciting the muscle into greater action, but was so hot as to produce a subsequent inflammation, if not an eschar, sufficiently great to cause the fibres of the *Levator palpebræ superioris*, either to adhere to each other, to the neighbouring parts, and by that means prevent their natural action; as happens in the adhesions that follow burns.

Though it has been said, that the weakness of the contraction of the *Levator palpebræ superioris* is the cause of the *trichiasis* of the upper lid, yet an inversion of the ciliary edge may be produced, not so much by that muscle being weaker than usual, as from its having a greater resistance than is natural to overcome, by the skin being thicker and more relaxed, or the cellular membrane enlarged, by the deposition of a small quantity of sebaceous matter, which will require a greater force to move and elevate it.

In that case undoubtedly, and probably some others, advantage might be obtained, by applying small slips of good adhesive plaster to the skin, between the upper lid, and the edge of the orbital process of the frontal bone, passing it over the eye-brow to the forehead, and drawing the skin in folds under it. An effect something similar to what is intended by the above contrivance, is observed in common life, as people when they are drowsy, and make great voluntary efforts to keep their eyes open, or to separate their lids under than usual, corrugate the skin of the forehead, by the contraction of the *corrugator supercilii*, and the *occipito-frontalis* muscles.

Relief too, in some instances would probably follow, as suggested by Mr. Mathen, from the application of an iron ring, so adapted to the form of the orbit, that part of it may press on the cartilage of the lid, and prevent its inversion; and

frequently a very trifling pressure will answer the purpose. The ring should be covered with silk, and made to fasten to the head with a ribband, by a projecting part from each side of the ring, to which it may be attached.

The *trichiasis* of the lower lid, is a much more common complaint, and when recent, can generally be cured, by a proper application of adhesive plasters. Often nothing more will be requisite, than to make a fold in the skin below the inverted lid, by drawing its edge from the eye, and fixing on a-piece of adhesive plaster to keep the fold in its situation.

But it sometimes happens that there is almost a constant flow of tears from the eye, particularly on touching the plaster, which moistens the plaster, that it either cannot be fixed firmly on, to preserve the necessary fold of the skin, or if it does adhere at first, the constant moisture of

osens it, on these accounts the use of adhesive plasters, has been often laid aside as ineffectual, when a little more attention to their application would have performed a cure, with ease.

I have frequently succeeded in such cases, by placing the patient's head in such a situation, that the tears, instead of running down the cheek, shall pass out, at the external angle, and by these means prevent the part, on which the plasters are to be placed, from being moistened, and when wiping it perfectly dry, have fixed the end of the plaster as near to the edge of the lid, as I conveniently could, and pulling the other end, have turned out the inverted edge as much as requisite, before the end, that was held between the fingers was applied to the cheek. At times two or three thin slips of adhesive plaster must be fixed on, though one is generally sufficient.

It is not absolutely necessary, that the plasters should be applied to the edge of the lid: indeed in some cases, if they are placed at a little distance, they will succeed as well, if not better, from much larger pieces of plaster being made use of, and the fold of the skin increased. The elevation proposed can occasionally be obtained with great facility, by placing a piece of adhesive plaster half an inch broad, and two inches long, nearly an inch below the edge of the lid, and having fixed one half of it, a probe being placed under the plaster, near that part which adheres, and pushed upwards, a very considerable elevation may be made, which will prevent inversion of the lid, and by the contraction of the corrugated skin, remove the cause of the complaint.

Most medical writers on disorders of the eyes, have described and recommended different forms of instruments, whereby taking hold of a small portion of

in, beneath the lid, and hanging by it, may by its weight, keep the lid from turning inwards, which, they say, will at the end cure the disease.

I do not doubt the application of an instrument, frequently answering the end proposed; at the same time, I must observe, I have met with no case in which such an instrument would be applicable, that might not be cured by a proper use of adhesive plasters, which ought certainly to have the preference, as they give much less pain and inconvenience, not to mention the unsightly appearance of a piece of skin depending from the cheek.

When the *trichiasis* is slight and recent, the application of adhesive plasters, will commonly be sufficient to restore the skin to its natural state, but in more obstinate cases, when the inversion is great, and the skin about the eyes very much relaxed, it may be necessary to cut off a small por-

tion of the loose skin, below the edge of the lid, and afterwards confine the lips of the wound together, by means of two or three sutures. The wound heals in a few days, and the lid is generally restored to its proper place.

In those cases, however, in which the ciliary edges are not only inverted, but also contracted in length, neither of the above methods will succeed; under such circumstances, relief is only to be obtained, by enlarging the circumference of the ciliary edges, either by making a fine incision in a right line from the outer angle to allow of sufficient motion to the lid, or by a compleat division of the cartilage of the lid, through the middle, which will cause each edge to recede towards the angle, leaving a division between them, which will not only remove the complaint for the present, but prevent the possibility of its return in future.

[125]

When the Ophthalmy proceeds from a supernatural row of hairs, growing out of the inner termination of the edge of the lids, which is generally attended with much trouble and distress, no means have hitherto been devised, better than the extraction of the supernumerary hairs, which only gives temporary relief; for though other attempts have sometimes been made, they have proved so very unsuccessful, as not to deserve further notice.

5. The fortuitous insertion of extraneous bodies between the lids, is another frequent cause of inflammation. These, during their continuance in the eye, produce great pain, and an inability to move the lids, with an increased secretion of tears. The flow of tears is, in general, sufficient to wash away the offending substances; which, on their failing to do, they must be removed by means appropriated to the nature of the adhering bodies, or the manner in which they are fixed in the eye.

The foreign bodies being either sand or dirt, the lids may be separated by the fingers, and on the patient being desired to look towards the opposite side to that in which the extraneous substances lie, they may often be removed with wet lint on the point of a probe, or with a small pair of forceps.

When there is reason to suppose there are many particles in the eye, it may be necessary to throw a stream of warm water over it, by means of a syringe, or to immerse the eye in a wine glass filled with water.

An eye cup, whose shape is adapted to the form of the eye, is generally recommended to be used in these cases, as well as for the application of lotions in Ophthalmy, which is very often attended with an evident increase of the complaint.

If a common wine glass is filled with water, and the head being bent forward

The edge of it is gently pressed between the lower lid and the cheek bone, the whole of the anterior part of the globe may be easily covered with water, and the upper lid allowed free motion: the depending situation of the head also, will permit more easily the removal of the offending particles.

But in applying a common eye cup, the pressure on the lids is considerable, and the heat from the eye rarefies the air in the cup, so much as to cause a considerable pressure and irritation; the quantity of water made use of is much smaller than with the glass, and the head being held back to suffer a more free admission of the water, the probability of the extraneous body being washed away is diminished.

When the adhesion of the extraneous body is so strong as to resist these attempts to remove it, it may be endeavoured to be separated by a thin blunt pointed scoop, something larger than a common probe, or

lastly, on that proving ineffectual, with the point of a lancet, taking great care that neither the *cornea* or the *tunica conjunctiva* is wounded, which, in the first case, might leave a scar that would afterwards injure vision; or in the latter cause so much irritation, as to render the subsequent inflammation dangerous.

If the extraneous particles adhere closely to the inside of the upper lid, though a free application of water is not sufficient to remove them, it will be necessary to turn the inside of the lid outward, which may be done without much difficulty: the lower edge of the lid, or part of the eyelashes, is taken hold of between the thumb and finger of one hand, whilst a probe, or any similar instrument, held in the other hand, is pressed on the outside of the lid, a little above the edge of its cartilage. On the lid being averted, the foreign particle is brought directly into view, where it may be removed, as was before directed.

ther by a little lint on a probe, or, in case of necessity, by the point of a lancet.

Lime, from its general consumption, and being often in exposed situations in considerable quantities, frequently by the carelessness of workmen, or the playfulness of children, gains admission to the eye, and is the cause of violent inflammations, and the most dreadful head-aches, followed by a large formation of matter, in the chambers of the eye, or extensive suppurations of the *cornea*, ending in incurable blindness.

Lime, applied to the living fibre, is a strong stimulant, and acts as a violent caustic, from its tendency to unite with the sebaceous matter of the body: in its unslacked state, it is very greedy of moisture, and in suddenly absorbing water gives out much heat, which considerably facilitates this combination.

Any large particle of lime should be immediately removed, by the means above recommended, for extracting sand, or dirt, and as the most serious consequences are to be apprehended, from a small quantity only remaining in the eye, it should be well bathed with some liquid.

Lime will unite with water in a very small quantity, and with the expressed oil more easily; but as it will combine with great facility, with all acids, they, in the diluted state, seem to claim the preference as an application in these cases. I have frequently found common vinegar, diluted with three or four times its bulk of water, applied with a wine-glass, as before directed, answer very well, and, as in case of emergency, it can generally be procured, it makes it the more convenient.

It is true, the diluted acid acts as a stimulant, yet in the degree of strength mentioned, it does not give much pain.

and although it produces a temporary increase of the inflammation, the action of the part soon abates, particularly if it is immersed in a little warm water, and the removal of the whole of the lime, is so desirable, that any momentary pain ought to be disregarded.

Let me again urge the necessity of the most care being taken, to wash away all the lime that may adhere to the eye, having frequently seen the worst effects arise from inattention on that head, though it is not at first attended with much pain, or apparent injury.

On examining the eye, after lime, in any quantity, has been in it, part of the cornea, or *conjunctiva* will be found white, and will have the same appearance as is seen, when a strong caustic is applied to other parts of the body, and in fact a combination is taken between the lime, and the corneal fibre; and as that must be removed:

ed, and new parts formed, by a peculiar animal process; care should be taken at least that sensitive action be interrupted by the injudicious application of stimulant medicines.

The eye should be defended from light by a paste-board shade, such as directed, Page 106, and the eye frequently bathed in lukewarm water, to wash away any matter that may be formed by the eye more immediately affected, and to remove the discharge from the eye, which is often adhesive and irritating: the lids also, when they are often a good deal swelled, stimulated by the acrimony of the tears, may be kept moist, by some mild ointment.

When the part, that has been burnt, is thrown off by the efforts of the vessels, and the pain, and inflammation have subsided, and an opacity of the *cornea* remains, it must be treated in the same manner as opacities after common inflammations.

I hope I shall be excused for once more repeating, that when the eye has been burnt with lime, lukewarm water only ought to be applied, as none of the common applications made use of, to subdue inflammations of the eye, are proper; and if it should be deemed necessary to do something more than patiently trust to nature, and frequent ablution, it may surely be considered a very pardonable imposture, to prescribe water, to which colour may be given, by a few drops of *tinctura croci*, or something of a similar kind; that the patient may be cheated, as it were, for his own advantage.

In the town of Birmingham, where I reside, whose inhabitants are chiefly employed in the working of metals into a variety of articles, pieces of steel are often struck off their tools into the eye, and from the defence afforded the eyes by the lids, the steel generally pierces the *cornea*, where it is either buried between its *laminae*, or

sticks in it, presenting a sharp surface to the irritable membrane which lines the lids.

The metal, in general, is very small in quantity, forming only a minute dark speck, and as, from the violence of the blow, it flies off in a heated state, the *cornea* is burnt, yet from the smallness of the particle on its being speedily removed, no perceptible opacity remains. It is always desirable to have it immediately extracted, for though if left in the eye, nature would remove it by suppuration, yet on that account the danger of its inducing an opacity, which might hereafter prevent perfect vision, is very much increased.

The method I take to extract particles of steel from the eye, is with a small straight bladed knife, made rather thick in the back, and not brought to a very fine point. Taking this in my right hand, whilst with the left I draw down the lower lid, the patient's head being supported by an assistant.

who stands behind, and elevates the upper
 end, on the point of the knife being placed
 under the particle of steel, it generally, on
 the first or second attempt, is disturbed, and
 the point of the knife, having been pre-
 viously well rubbed on a loadstone, attracts
 the steel, which is immediately removed.
 This knife I also find very convenient for
 extracting extraneous bodies of different
 kinds, and from its form and strength much
 preferable to a lancet. The lancet is
 awkward to turn about in the hand, and
 having a double edge, often wounds the
 edges of the lids; it is also brought to so fine
 a point, as frequently to do considerable
 injury to the eye.

When the steel has buried itself in the
cornea, it becomes much more difficult to
 extract, as it will be requisite to insert the
 point of the knife into the small aperture
 made by the passing of the metal, and lay
 down the *cornea* down to it, when it will be
 nearly in the same state as if situated on the

surface, and must be extracted in the same manner.

If the steel is situated beyond the centre of the *laminæ* of the *cornea*, and more interiorly, towards the anterior chamber of the aqueous humour, great nicety is required in extracting it, least an opening should be made into the anterior chamber, the metal forced into it, and the patient lose his eye. In such cases it may be best to trust to nature, who will often succeed; for the practitioner would certainly be much reflected on, the patient not knowing the peculiarities of his case, if, in attempting to extract a bit of steel, the sight should be destroyed.

In instances of this kind, as the *cornea* in its natural state is insensible, and no rough part of the steel rubs against the lids, no much irritation is excited, suppuration very slowly proceeds, and as it inclines to the surface of the body, the metal by degrees comes out with the matter.

It might, probably, be supposed, that were the particle of metal to gain admission into the anterior chamber, the inflammation produced would be so great as to cause a suppuration of the internal parts of the eye; and an evacuation of the humours, and that every method ought to be tried to extract the article in the first instance.

It has already been shewn, in speaking of the effects of blows, that the *chrystalline* lens has been confined in the anterior chamber for years, without producing much pain, and a justly celebrated lecturer, of London, informed me, that once, in making the section of the *cornea*, to extract a cataract, the point of the knife broke, and he believes remained in the eye, though the patient recovered without any unfavourable symptom. The point of the instrument, however, might never have been in the eye; it might speedily have been washed away by the flowing of the aqueous humour; I will therefore relate a case, exactly in point,

wherein will be seen the effects of a piece of steel remaining in the aqueous humour.

A few years since, a young man came to me from the country, to have a something taken from his eye, which, he said, flew into it a few days before, as he was hoeing turnips. The eye was very slightly inflamed, and on one side of the lower part of the *cornea* was seen, what appeared to me a piece of iron, a good deal larger than usually found in similar cases; the *cornea* was completely healed over it, and as he did not complain of pain, I declined attempting to extract it; gave him a mild lotion, and desired him to come again in a week.

It was nearly a month before I saw him again when the inflammation of the eye was nearly the same as on my first seeing him, at times giving him a little uneasiness; if he looked at any thing earnestly, the eye in a short time was filled with water: I now found the extraneous substance, which

had before observed in the *cornea*, to be in the anterior chamber, a very short distance from where I had first seen it, and hanging by a small filament of matter, in the aqueous humour, and waving, on the eye being moved.

On bringing a large magnet near his eye, and moving it in different directions, the substance obeyed its motions, shewing it to be iron; and though I attempted to remove it, by placing the magnet in various situations, and giving the particle of iron both a quick and slow motion, I was not able to detach it from the *cornea*.

A few weeks afterwards he called on me again, and said the eye was better, but rather weak, and thought if he could have something to *strengthen it*, he should soon be well. The eye still looked slightly inflamed, or as if it was exposed to some small irritation. I particularly requested he would call again, but since that time

have never seen him; I conclude, from the circumstances of the case, he has not experienced much inconvenience from it.

6. Essential oils, alkalies, acids, acrid juices, &c. are next mentioned, as exciting causes of inflammations of the eyes. For their removal, recourse should be had to frequent ablution with warm water, and on an inflammation succeeding, it ought to be considered, whether there has been any part destroyed, by a combination between the animal fibre and the foreign substance, or whether it has acted merely as a stimulant, and increased and diseased actions of the vessels been induced in consequence.

In the first instance the same directions are applicable, as are given after the eye has been burnt with lime; but in the latter case, it must be treated the same as an Ophthalmia, after the application of cold, which will be considered in the second part.

7. When an Opthalmy arises from exposure to too much light, it generally proceeds from a predisposition of the part to diseased action. The exciting cause must be avoided, and the case treated the same as a common inflammation.

8. The eighth cause mentioned, as producing Opthalmy, is the increased secretion of tears, especially with their confinement, to the eye.

With respect to the Opthalmy occasioned by much weeping, from mental affections, or the almost constant flow of tears, which attends the rubiulous eruption, I have no observations to make which are applicable to this place. It is not so with the inflammation of the eyes, which, more or less, is brought on by the eruption of the small pox, when the pustules on the face are in any quantity.

A small pock pustule has its margin inflamed, and swelled, and when the pustules are numerous, the whole external surface of the body is enlarged. The pustules having the effect of inflaming, and enlarging the part, on which they arise, if situated on the eye lids, they become inflamed and swelled; they are closed, and in a short time very firmly so, by an adhesive matter, discharged from the eyes. The question, then, is, whether we ought to endeavour to prevent this closing of the lids, or not.

Some, from the idea that the lids, being closed, are a great preservation to the eyes, will on no account allow them to be touched; but suffer the discharge to collect in considerable quantities, about the edges of the lids; whilst others, from an over officiousness, are almost constantly endeavouring to separate them, to remove the matter as it is secreted. Between these two practices, a due medium should be

preserved. The swelling of the lids being
 the uniform effect of the pustules, cannot
 be abated, but by the natural course of the
 disease, and what we ought to attempt is,
 that the eyes should be irritated as little
 as possible, and the confinement of the
 tears prevented. The lids should be
 touched with the greatest gentleness, any
 matter that collects, removed with a little
 warm water, and the edges of the lids
 kept *very moist* by the application of some
 mild ointment.

The lids should never be free from the
 ointment, that it may prevent the confine-
 ment of the acrid tears, which is generally
 the cause of the succeeding inflammation:
 Besides, if they are kept constantly moist,
 there will be no danger of *strabismus*, which
 sometimes is brought on, by the eye-lids
 being partially opened, on the the decline
 of the disease, and the patient endeavour-
 ing to see, through the small part which
 admits light.

For the inflammation that is an attendant on the *fistula lachrymalis*, some relief may be experienced, by frequently emptying the lachrymal *sac*, by making a pressure on it with the finger; yet its removal must not be expected without the cure of the original disease, but how, or by what means that is to be effected, would lead into by far too long a discussion to enter upon here.

9. Erysipelatous and inflammatory affections of the parts adjoining the eye, the last causes mentioned, as producing Opthalmy by a direct increase of *stimulus*. The original inflammatory infection which requires our first attention, and all that is to be done for the inflammation of the eye, is to keep it cool, and as free as possible from all irritating causes.

END OF PART FIRST.



ON THAT

DISORDER

OF THE

Pekell

CORNER of the EYE,

COMMONLY CALLED

Fistula Lachrymalis.

By PERCIVALL POTT,

Senior SURGEON to St. Bartholomew's-Hospital.

*Neque enim credunt posse eum scire quomodo morbos curare
veniat qui unde hi sint ignoret; eum vero recte curaturum,
in prima origo causæ non fefellerit.*

A. CORN. CELSUS.

The SECOND EDITION, improved.

L O N D O N :

Printed for L. HAWES and Co. at the *Red-Lion* in *Pater-*
noster-Row. M. DCC. LXIII.

1650234

T H E

P R E F A C E.

***B**^R frequently conversing with some of that part of the profession who come to London to attend the Hospitals, and to improve themselves in the Art of Surgery, it has appeared to me that the FISTULA LACHRYMALIS, tho' a very common disease, is one with which many of them are very little acquainted, either with regard to its cause, seat, or method of cure. Some are totally ignorant of every thing relating to it, others who have an imperfect idea of its nature, and seat, are yet much at a loss how to vary the method of treating it according to its different states and circumstances;*

upon which distinction the probability of a cure does often in great measure depend; for if those means which are only proper in one state of the disease are used in another, the patient will be fatigued to no purpose, and the surgeon by being frequently disappointed will be inclined to think those cases incurable, which have only failed through his own mismanagement.

THERE is hardly any chirurgical disorder which requires a more close regard to all its appearances and variations than this does; and whoever expects to conduct it successfully, must attend to it constantly: this is, perhaps, the great reason why it is so little understood; the object is too minute, and the process often too long to engage the attention; besides which, it hardly comes under the name of an operation, the great and almost only object which they who come hither from the distant countries have in view: the operative part of surgery is what they

have seen the least of, and therefore they are the more desirous of becoming acquainted with it; this desire is a very laudable one, and ought certainly to be encouraged, but still the operative part of surgery is far from being the whole of it; and I cannot help thinking, that by attending a little more to what is called common or practical surgery, our art might still be considerably improved, practitioners rendered more expert, and mankind much benefited.

THE merely curing diseases is not all; that was done (sooner or later) while surgery and anatomy were in their most imperfect state, and while every branch of medicine laboured under many inconveniences which are now happily removed; but the different methods in which chiral disorders are treated, or their cures attempted, will make so considerable a difference in the confinement and sufferings of the patient, as to be very well worth attending to.

IT

IT may possibly be thought foreign to my present purpose, but I cannot omit this opportunity of adding a few words on a subject which appears to me highly deserving of some notice, as its influence may be very extensive and very prejudicial; it is the false idea which the by-standers at an operation generally have of chirurgic dexterity; to which word they annex no other idea than that of quickness. This has produced a most absurd custom of measuring the motion of a surgeon's hand, as jockeys do that of the feet of a horse, viz. by a stop-watch; a practice which though it may perhaps have been encouraged by operators themselves, must have been productive of most mischievous consequences. Tute et celeriter are both very proper characteristics of a good chirurgic operation; but tute stands as it should do, in the first place, and the patient who suffers the smallest injury from the hurry of his operator has no recompence from the reputation which the latter obtains from

capital operations unforeseen circumstances will sometimes occur, and must be attended to; and he, who without giving unnecessary pain from delay, finishes what he has to do in the most perfect manner, and the most likely to conduce to his patient's safety, is the best operator.

I have endeavoured to make the following tract as plain and as intelligible as I can; and if it should appear prolix to those who are already acquainted with the subject, I must beg leave to observe, that it was not written for their information; but if any of those who were unacquainted with it before should from hence gain any useful knowledge, my end will be answered, and I shall be very much pleased.





O F T H E

Fistula Lachrymalis.

S E C T. I.

HE antient writers were in general so little acquainted with the anatomical structure of the parts concerned in this disease, that both its cause, and seat have been very erroneously represented by most of them; other disorders, very different both from this and from each other, have been confounded under the same general appellation, and the means made use of toward obtaining a cure, being adapted to such misconceptions, were rough, painful, and most commonly ineffectual.

THE fluid which perpetually moistens the eye, was supposed to be secreted by
B that

now called the caruncle, and to flow from thence upward thro' the puncta lachrymalia *. The caruncle was by many thought to be the seat of the disease in question, which was said to be produced, either by a defluxion from the brain † on this part, or by an abscess formed within the body of it; or by a lodgment of the tears, become acrid and corrosive in consequence of such stagnation ‡; while others looked

* FALLOPIUS, who has very accurately described the puncta lachrymalia, sacculus, and duct, as well as the disease, has yet fallen into this common error. “Ad oculos ipsos ex faucibus egrediens venio, in quibus primum præter misere anatomici duo foramina parva in angulo interno posita, quarum unum est in palpebrâ superiori, alterum in inferiori, in viventibus adhuc hominibus, si quis inspicere voluerit apparentia, quæ foramina habent meatus quos sub carunculâ encanthidos vel epicanthidos dicta uniuntur in quendam communem sinum in narium cavitatem definitentem per canalem proprium in osse squamoso, quod internum angulum occupat insculptum.

“PER hos meatus major lachrymarum pars ut ego in flexibus mulierum observavi, ad oculos emanat.”

FALLOPIUS.

Non enim os solummodo cariosum, verum etiam glandula ita erosa erat, ut quotiescunque puer ploraret lachrymæ per ipsam fistulam copiosè extillarent. HILDANUS.

† FISTULA lachrymalis fit ex humorum decursu, qui currunt ad lachrymalis angulum juxta nasum, nec propter eorum multitudinem, et grossitiem possunt exire, &c. hi autem morantes ibi diutius corrumpuntur, et locum ulcerant,

LANFRANC.

ÆGYLOPS est tumor abcessorius inter majorem angulum, et nares proveniens. PAULUS.

‡ AT the great corner of the eye there is a glandule made for receiving and containing the moisture which serves for lubricating

looked upon it as a kind of encysted tumor. The swelling in the inner corner of the eye, the frequently attendant ophthalmia, the involuntary flux of serum down the cheek, the excoriation of the eye-lid, and the discoloured discharge upon pressure, strengthened their opinions, and confirmed their prejudices.

THEY who supposed it to be caused originally by a defluxion of the inflammatory kind, tending to produce an abscess, had recourse at first to those general methods and means which were thought most likely to prevent such consequence: these not answering, they proceeded to open the supposed abscess, and to endeavour the digestion of it; on the other hand, they who supposed it to be an encysted tumor attempted the eradication of it, either by knife, caustic, or cautery; and all of them taking it for granted when the discharge

B 2

was

lubricating the eye; this glandule, sometimes by a sanguine or pituitous defluxion falling violently from the brain, swells and imposthumates, and ulcerates, &c. AME. PAREY.

Hæc caruncula ab acrim humorum affluxu turgescit nonnunquam intumescit, et abscedit ulceraturque, ulcere non raro in fistulam abeunte adeo ut subiectum os corrumpatur:

MUNNICKS.

PER pusillum utriusque palpebræ foramen lachrymæ naturaliter effluunt. FAB. AB AQUAPENDENTE.

LACHRYMÆ veniunt per lachrymalia a foramine quodam arvo, et quasi insensibili in fine pilorum. GUIDO.

was apparently purulent, or much discoloured, that the bone was rotten, advising the use of escharotic applications, or the use of iron to destroy the callosity, and to dry and exfoliate the caries; and these methods failing, as in the nature of things they very frequently must, they pronounced the disease to be incurable.

A MORE minute and careful examination into the anatomy of the parts has given us a more true idea of the disorder, and furnished us with a more rational, as well as a more successful method of treating it. We now know, that the caruncle is not the organ which secretes the tears, but that this office is performed by a gland, situated near the outer corner of the eye; that the lachrymal fluid is in its nature perfectly innoxious; that an obstruction in the nasal duct is most frequently the primary and original cause of the complaint; and that its seat is in the *sacculus lachrymalis*.

UPON these principles the modern practitioners have, with great industry and ingenuity, endeavoured to find out some means, whereby this obstruction may be removed, and the parts restored to their natural and healthy state, without such pain and destruction.

destruction, and deformity, as the antient methods occasioned; or, these failing, to establish a new artificial passage, which may in some measure supply the place of the natural one.

ALL these means have the merit of being founded on the natural structure of the parts concerned: when the more easy, and mild ones succeed, the patient gains a considerable advantage; and when they do not little time is lost, nor is any more efficacious method rendered thereby less practicable: in this, as in every other part of surgery, the more simple means ought to be first tried, pain should be avoided as much as possible, except when absolutely necessary, and then it must be submitted to.

S E C T. II.

THAT the motions of the eye-lids may be performed with the utmost ease, that the tunica cornea may be kept constantly clean, bright, and fit for the transmission of the rays of light, and that dust, and other hurtful particles, may be immediately

immediately washed away, the surface of the eye is continually moistened by a fine limpid fluid.

THIS fluid is derived principally from a large gland, situated under the upper edge of the orbit, near the outward corner of the eye, which gland is of the conglomerate kind, and lies in a small depression of the os frontis; its excretory ducts, or those by which it discharges the secreted fluid, piercing the tunica conjunctiva, just above the cartilaginous borders of the upper eyelids.

WHILE the caruncle was thought to be the secretory organ of the tears, this gland bore the title of *glandula innominata*; but now that its use and office are known, it is called *glandula lachrymalis*.

By irritation from any sharp or poignant particles, a large quantity of this fluid is immediately secreted, and by the motion of the eye-lids is as immediately derived over the surface of the eye, by which means such particles are washed and wiped off. Sometimes also the passions of the mind produce an immediate increase of this

perly, called tears ; a constant secretion of too large a quantity causes a disease, called epiphora ; and a deficiency of it makes the motions of the lid difficult and painful.

ALTHOUGH the fluid secreted by the lachrymal gland is considerable in quantity, yet, when it is not suddenly produced by irritation from without, or passion within, it is so constantly and gradually carried off, as to create neither trouble, uneasiness, nor blemish.

THE edge, or border of each eye-lid, is formed by a thin cartilage, the figure and consistence of which keep the lids properly expanded ; these cartilages are covered by a fine membrane, and are called cilia, their internal edges, do, upon every motion, sweep over every point of the surface of the cornea ; this motion, tho' almost imperceptible, unless attended to, is very frequently performed, and as the secretion of the fluid is also constant, the eye is by this means kept always moist, clean, and bright.

AT

AT the extremity of each of these cartilaginous borders of the eye-lids, on the side next the nose, is a small papilla, or eminence; and in the middle of each of these is a small hole, or perforation, which being made in the cartilage is not liable to collapse while the parts are in a sound state, but remains always open; they are called the *puncta lachrymalia*, and their office is to receive the lachrymal fluid, as it runs off the cornea along the edges of the eye-lids, thereby preventing it from trickling down the cheek; and that there may be no impediment to the constant execution of this office, during the time of sleep, as well as that of being awake, the internal edges of the cilia do not come into immediate contact with each other in that point where these orifices are.

FROM each of these *puncta lachrymalia* proceeds a small membranous tube, which tubes soon enter into, or form a pouch or bag, situated near the inner angle of the eye, just below the union of the two lids, under the *musculus orbicularis palpebrarum*; the bag is called the *sacculus lachrymalis*, and its office is to receive all the lymph

the upper part of this sacculus lies in an excavation, formed partly by the nasal process of the os maxillare superius, and partly by the os unguis; the lower part of it is confined in a long channel, and forms a tube, or duct, which descending obliquely backward, communicates with the cavity of the nose, behind the os spongiosum superius, by an opening whose size is somewhat different in different subjects.

THIS passage is called the ductus ad nares, or the ductus nasalis, and thro' it, whatever is received by the sacculus from the puncta does, in a healthy and sound state of these parts, pass into the nose.

THE membrane which lines this sacculus and duct, is in its structure much like to the membrana pituitaria narium, from the surface of which a clear viscid mucus is secreted, and by which the sacculus and passages are constantly moistened and kept pervious.

WHILE the parts are in a healthy, sound state, the fluid secreted by the lachrymal gland passes off thro' the puncta, sacculus,

C

and

ble, but when they are in a diseased state the case is otherwise ; this membrane, like all other vascular parts, is liable to inflammation, by which means it often happens, that it is so thickened as to obstruct the nasal duct, and thereby much impedes or totally hinder the passage of any thing thro' it, in consequence of which obstruction the sacculus is filled by its natural mucus, and the derivation of the serum from the lachrymal gland thro' it being thus prevented, it runs off from the eyelid down the cheek ; this obstruction continuing, and the mucus still lodging, the sacculus is dilated, and produces that tumor in the inner corner of the eye, and that discharge, upon pressure, which characterize the first state of the disease in question, and, in conjunction with several other attending symptoms, prove its seat to be in the lacrymal sac, and nasal duct.

S E C T. III.

ALTHO' the seat of this disease is the same in almost every subject, yet its appearance is very different in different persons, and under different circumstances; these variations depend principally on—

1. THE degree of obstruction in the nasal duct.
2. THE state of the cellular membrane covering the sac.
3. THE state of the sacculus itself.
4. THAT of the bone underneath.
5. THE general state and habit of the patient *.

SOMETIMES a serous kind of defluxion, which the lining of the sac and duct is so thickened as to obstruct, or prevent the passage of the fluid thro' them into

C 2

the

As the state and circumstances of this disease are really various, and differ very essentially from each other, the general custom of calling them all by the one name of *fistula lachrymalis* is absurd.

the nose, makes the whole complaint, and the cellular membrane on the outside not being diseased, there is no appearance of inflammation; in this case the duct is stopped, and the sacculus dilated, but without any alteration in the colour of the skin, a fulness appears in the corner of the eye next to the nose, and upon the application of a finger to this tumor, a clear viscid mucus is discharged thro' the punctum lachrymalia, the patient feels no pain, nor finds any inconvenience, except what is produced by the discharge of this mucus and by the trickling of the lymph down the cheek.

IN some cases the mucus is not perfectly and always clear, but is sometimes cloudy and looks as if it had a mixture of milk or cream in it, at first waking some of it is generally found in the corner of the eye and the eye-lashes being smeared over with it during sleep, most commonly adhere together in the morning.

THIS is the most simple state of the disease, what the French have called the *hernia*, or *hydrops sacculi lachrymalis*; it is frequently met with in children who have been

been tickety, or are subject to glandular obstructions, and in this state it sometimes remains for some years, subject to little alterations, as the health or habit shall happen to vary, the sacculus being sometimes more, sometimes less full, and troublesome, the mucus which is pressed out sometimes more, sometimes less cloudy, and now and then attended with a slight ophthalmy, or an inflammation of the eyelids, but which, by common care, is easily removed.

IF the sacculus is not much dilated, the discharge small, and produced only by pressure, the chief inconveniences are the weeping eye, and the gumming together of the lids, after sleeping; but these, by being attended to, may be kept from being very troublesome, and if the disease makes no farther progress, may be so regulated as to render any more painful process totally unnecessary.

IF the dilatation is considerable, the swelling is more visible, and the quantity of mucus is larger; it is also in this state more frequently mixt and cloudy, and
more

more troublesome, from the more frequent necessity of emptying the bag ; but if the patient be adult, it may, even in this more dilated state of it, be kept from being very inconvenient.

If an inflammation comes on, the tumor is thereby considerably increased, the discharge is larger, as well during sleep, as upon pressure, the skin covering it, loses its natural whiteness and softness, becomes hard, and acquires an inflamed redness, and with the mucus a mixture of something, which in colour resembles matter, is discharged, especially if the pressure be made with any force, or continued for any time : this circumstance, added to the painful sensation, and inflamed appearance of the parts, has been productive of a supposition, that in this state there is either an ulcer, or an abscess within the sacculus or duct.

As this is an opinion, which tho' it may possibly sometimes have some foundation in truth, yet is in general entertained much too hastily, and is also the principal source whence most of the mistakes concerning this disease have sprung, I would beg
leave

leave to be managed in a few words on this subject.

It has already been observed, that from the surface of the membrane which lines these parts, a thin mucus is secreted, by which its surface is smeared over, in the same manner as is that of all the membrane which covers or lines the fauces, larynx, and internal parts of the nose, the antra of the jaws, and the sinus's of the sphenoid and ethmoid bones, &c. While the lachrymal sac is free from disease, and the ductus ad nares open, this mucus is nearly limpid in colour, small in quantity, and passes insensibly into the nose with the fluid from the lachrymal gland; but when, by the obstruction of the nasal duct, that passage is denied, it necessarily lodges in the sacculus, by distending and irritating its containing bag, it is increased in quantity, altered in colour, and discharged at the puncta lachrymalia, as it either becomes too much for the sac to contain, or as it is forced out by pressure. This is a short and succinct account of the true nature of the disease, and such as will fairly and truly account for all its symptoms and appearances, without having recourse to

3 either

very seldom, if ever, attend it.

THAT which is mixed with the clearer part of the mucus, and which from its pale yellow hue is taken for matter, is not matter, but mucus, which in this part, as well as several others in the body, does, either by being confined beyond the necessary time, or by inflammation, or irritation of the gland or membrane which secretes, or contains it, or even from general affection of the habit, put on a yellow, purulent colour, when there is neither abscess, nor ulcer in the part whence it comes.

So many instances of this are producible as to put the matter beyond all doubt, the urethra, vagina, and all the sinus's of the head which communicate with the nose, furnish us with them daily; the linings of all these are constantly imbued with a mucus naturally clear, and no more in quantity, than is necessary to keep the membranes moist, but either inflammation or irritation does immediately so add to its quantity, and so alter its colour, that in the two former, the same mistake has often been made as in the subject in question, that
is,

purulent, and produced by ulceration of the parts.

These two fluids pus and mucus, which have been so frequently confounded together, do really differ so widely from each other in their nature, constitution, sources, purposes, and effects, that to distinguish them properly, and to point out the true character of each, seems to be a matter of much importance; it would carry me too wide from my present purpose to attempt it in this place, and therefore I shall only just mention what may serve merely to illustrate that.

If I conceive rightly of this affair, mucus, considered in a general sense, is the effect of a natural secretion made by glands, membranes, or other bodies appointed for that purpose, and is so far from being originally the consequence of disease, that, in a due quantity, it is absolutely necessary for several very important purposes in the animal œconomy, which purposes, when this fluid is deficient, must be ill-executed, and some kind of disease, or defect follow: whoever will reflect on the uses of it

D

in

fulæ, of the tendons, in the sinus's of the skull serving the purposes of speech, in the cavity of the nose where the olfactory nerves do their duty, in the prostate gland, larynx, trachea, urethra, and vagina, will be easily convinced of the truth of this assertion, both with regard to its natural uses in a healthy state, and proper quantity, and the share it frequently has in the production of diseases, when it is either vitiated or redundant.

Pus, or matter, is certainly no natural secretion; suppuration, tho' it is an act of nature when some parts of the body have been forcibly divided from each other, is nevertheless to be regarded as the effect of violence and destruction, at least of division; for, without entering minutely into the origin or nature of it, I believe, I may venture to affirm, that the dissolution of some of the solid particles of broken capillary vessels, and a mixture of some part of the juices which should circulate thro' them, make a necessary part of its production; however constant its appearance may be in the progress toward healing a wound, or sore, yet it

never

never is produced, even in the smallest quantity, without some degree of erosion, some breach in the natural structure of the parts, and when such breach is healed the discharge necessarily ceases.

ON the contrary, mucus may by irritation, relaxation, or defluxion, on its secreting or containing parts or organs, be increased to a quantity far beyond what is necessary or useful, and produce thereby a disease in parts where there is not the least degree solution of continuity, as in the cases of tenesmus, stone in the bladder, fluor albus, and simple gleet from the urethra; as also in that kind of defluxion on the nose and fauces, producing a catarrh, and in the immediate effect of all sternutatories.

OTHER differences between the nature and properties of the two fluids might be mentioned, but if these already cited are just, they will be sufficient to evince the impropriety of confounding them together, either with regard to theory or practice.

NOR is this mistake of discoloured mucus for matter confined to the lacrymal sac

only ; the two circumstances of pain, and yellow colour, having, in almost all times, produced the same misconception in the virulent gonorrhea of both sexes ; this has been called pus, and being said to proceed from ulcerations in the urethra and vagina, tho' the repeated testimony of those who have, immediately after death, examined the parts of persons so diseased, has often been produced to the contrary ; and tho' the discharge itself, when properly examined, will always prove the contrary ; inflammation and irritation of the membranous linings of the urethra, and vagina, will fully account for all the appearances in this disease, in which there is neither matter, nor ulcer, nor abscess : whoever will attend to the discharge made from a purulent ulcer, will find it widely different from that which issues from either of the above parts in a gonorrhea.

AGAIN, in case of strictures in the male urethra, the discharge occasioned by a bougie, properly and judiciously used, is a discoloured mucus, and not matter, tho' it is generally so called ; it is from the discharge of this mucus, and the dilatation of the passage, that the relief is obtained, not from

from any destruction or division of parts ; the bougie, which produces true matter, does much more harm than good, makes a fore where there was none, and where there ought to be none ; how often do catarrhus defluxions on the trachea, and larynx, wear toward the close a deep purulent colour, so as to deceive the unknowing into an opinion, that it is matter from the lungs ? But no judge of these things ever had recourse to abscesses or ulcers for a solution of such appearance. The argument drawn from the quantity of these discharges is as erroneous as those taken from its colour ; as an inflammatory defluxion on the part does generally occasion the latter, so meer irritation will produce the former, which does also generally cease when the irritating cause is removed or appeased : how immediately is a most troublesome tenesmus cured by a glyster of starch and opium ? What large foetid discharges are made from behind the prepuce of many persons, not only free from all venereal teint, but without any ulceration of the parts, by a kind of exsudation ? To what length of time will they not continue, if neglected, and how immediately do they cease by the use of a spirituous

ous or vitriolic wain? How often is the fluor albus, even in some of its worst circumstances, moderated, not to say cured, merely by washing away the acrid mucus, which lodging in the rugæ of the vagina, continually irritated the parts to a fresh discharge, and perpetuated the disease? What quantity of slime is there in the urine of those who have a stone in the bladder? And how totally does it cease, upon that stone being discharged, or taken away? Whereas neither cleansing of parts, nor removal of irritating bodies, does, or ever can procure an immediate cessation of a discharge of true matter, which being occasioned by a solution of continuity, an erosion or division of the parts whence it proceeds, must decrease gradually, and at last can only cease by such part becoming whole again.

IN short, the two fluids are so absolutely different and distinct, that the blending them together in our ideas of disease, proceeding from, or producing either of them, cannot be too industriously avoided; it is a subject on which a great deal more might be said, as it would comprehend, or have relation to many disorders, which perhaps
are

to; but being beside my present purpose, I shall say no more about it, only desire, that I may not be misunderstood, as if I meant to assert, that there never is abscess or ulcer in the lachrymal sac, and duct: No, I only mean to signify, that it is my opinion, that the yellow, or purulent colour of the discharge, which is generally received as a proof of such, is no proof at all; that this colour may be, and most frequently is, dependent on other causes; that tho' by the suppuration of the cellular membrane covering the sac, the upper part of it sometimes becomes sloughy, and bursts, yet the lower part of it, and the nasal duct, are often at the same time perfectly sound; and that there never is abscess or ulcer within, while the skin is entire and preserves its natural hue, and softness, let the colour of the discharge be ever so yellow; circumstances of no small consequence in the treatment of this disease.

THE inflammation of the cellular membrane covering the sac, is a circumstance which makes a considerable difference, both in the appearance of the disease, and
in

is confined merely to the surface of the tumor in the corner of the eye; in others it spreads still farther, affecting the eye-lids, cheek, and side of the nose.

WHEN the parts are in this state, the mucus within the bag has generally the appearance of being matter, that is, it wears a deep yellow colour, and is of a more thin consistence; if the puncta lachrymalia are naturally large and open, and the inflammation confined to the surface of the sac, its contents will pass off pretty freely, and the skin will remain intire; this is what the antients called the simple, or imperfect, or anchylops.

BUT when the skin covering the lachrymal bag has been for some time inflamed, or subject to frequently returning inflammations, it most commonly happens, that the puncta lachrymalia are affected by it, and the fluid not having an opportunity of passing off thro' them, distends the inflamed skin, so that at last it becomes sloughy, and bursts externally. This is that state of the disease which is called perfect, Aigylops, or Ægylops;

thro' the puncta lachrymalia, while the skin was intire, is now made thro' the new opening, and by excoriating the eye-lids and cheek increas'es the inflammation, and gives the disease a much more disagreeable appearance. In some the matter bursts thro' a small hole, and after it has discharged itself, the tumor subsides, the neighbouring parts become cool, and tho' the skin covering the surface of the sacculus is sloughy and foul, yet there is no reason to believe that the sac itself is much diseased below; in others the breach is large, the skin remains hard and inflamed, and from the appearance of the sore, there is reason to suppose the whole inside of the bag to be in a diseased state; and in some cases, which have been much neglected or irritated by ill-treatment, the cavity of the sacculus seems to be filled with a loose ill-natured fungus, which gleans largely, and produces inflammation and excoriation of all the parts about.

THERE is also another circumstance which sometimes is found to attend this disorder, viz. a carious state of the bones; this was by our forefathers supposed to be

E

a fre-

son for their so free use of caustic, cautery and scalpra, in the treatment of it; but since the disease has been more minutely examined into, this circumstance has been found to be a very rare one. When the fistula lachrymalis is a symptom of the lues venerea, as it sometimes is, the bones are indeed often carious, but then, the fistula is not the original complaint, but produced secondarily, and is a consequence of the diseased state of the os ethmoides, and ossa spongiosa of the nose, and is not curable by any local means or applications, but depends intirely on the cure of the disease, of which it is a symptom.

I HAVE also seen an abscess after the small-pox, which by falling on the lachrymal bag, has made it all slough away, and leave the bones bare, which circumstance I have also seen attend the free use of strong escharotics applied to destroy what is called the cyst, but without the accession of some other disorder producing it, or the most absurd method of treating the complaint; I believe that a caries of the bones will very seldom be met with. Indeed the combination of other diseases, either of the general habit, or affecting the
same,

same, or the neighbouring parts, does often make a very material difference, both in the appearance of the disorder, in the prognostic, and in the proper method of treating it, which therefore should always be enquired into; for instance, the patient is sometimes subject to an habitual ophthalmia, or a lippitudo, which will add to the deformity, and give a good deal of additional trouble during the cure; an opzæna, or some other disease of the membrane, and cells of the ethmoid bone, or a polypose excrescence within the nose, are now and then combined with it; the habit is sometimes, (as I have before observed) infected with the lues venerea, of which this disease may be a symptom; strumous glandular obstructions are its too frequent companions, and what is worst of all, it is sometimes cancerous.

S E C T. IV.

FROM what has been said, I think it will appear that this disease, in its primary and most simple state, consists in a retention or lodgment of mucus in the sacculus lachrymalis, in consequence of an

E 2 obstruction

obstruction of the natural passage from the bag into the nose; that by means of this lodgment the sacculus is distended, irritated, and sometimes inflamed; that the fluid which passes from the lachrymal gland over the eye to the puncta lachrymalia, being prevented by the fulness of the sac from getting into it, runs down the cheek, and therefore that the characteristic marks of the disorder, when recent, are a small tumor in the inner corner of the eye, an involuntary flux of serum down that side of the face, and a discharge of mucus thro' the puncta lachrymalia upon pressure.

THIS lodgment, being originally produced by the stoppage of the natural duct, it follows, that the first curative intention, is the removal of that obstruction; which is sometimes practicable, but more often not; the degree of obstruction, its date, the state of the adjacent parts, and some other circumstances rendering it more or less so in different subjects.

THAT the inexperienced practitioner may be guarded against giving a hasty prognosis; or making attempts, which
however

however fatiguing to the patient, must in the end prove fruitless; and that he may be enabled to understand the disease more perfectly, I shall take the liberty to divide it into four general heads, or states, under which all its lesser distinctions may be comprehended.

THE first consists in a simple dilatation of the sacculus, and obstruction of the nasal duct, discharging, upon pressure, a mucus either quite clear, or a little cloudy, the skin covering the bag being intire, and perfectly free from inflammation.

IN the second, the tumor is somewhat larger, the skin which covers it, is in an inflamed state, but intire, and the discharge made thro' the puncta lachrymalia, is of a pale, yellow, or purulent colour.

IN the third, the skin covering the sacculus is become sloughy, and burst, by which means the swelling is in some measure lessened; but the mucus, which while the skin was entire, used to be pressed out thro' the puncta lachrymalia, now discharges itself thro' the new aperture; the ductus ad nares, both in this and the preceding

By the improper use of medicines of the latter kind, it frequently happened that the skin became inflamed, and burst, the discharge which necessarily followed this accident, together with the heated appearance of the parts about, confirmed their opinion of a collection of matter within, and according to such supposition, they attempted to obtain a cure by dilating the orifice, and endeavouring to make an incarnation from the bottom of the hollow; not being acquainted with the situation, or use of the nasal duct, they took no care to free it from the obstruction under which it laboured, but dressing the sore like a common impostumation, permitted it either to be filled up with a loose fungus, or to contract itself to a narrow fistulous orifice, which daily discharging a discoloured kind of fluid, and not healing by such means as they made use of, they concluded the bone underneath was carious, and made way
down

I believe no one will venture to say, that the nature and seat of the disease is more or better explained by what the Doctor has said, than by the Serjeant; and I think it is perfectly clear, that neither of them had any true idea of it at all, they both mistook the caruncle for the lachrymal gland, and the disease for an encysted, or a scrophulous tumor, which ought to be brought to suppuration; the lachrymal sac, the ductus ad nares, their use, and the disorder of them creating the complaint in question, they were totally unacquainted with.

with a cutting instrument, or by destroying them with caustic, and cautery, intending to procure an exfoliation, and thereby a firmer basis to heal on *.

BUT since the use of the ductus nasalis has been known, since it has been discovered that an obstruction in this, is the primary and principal cause of the disorder, and that what passed for the cavity of an abscess is really the sacculus lachrymalis, both the intention of cure, and the means, have been considerably altered.

IN the first and most simple state of the disease, viz. that of mere obstruction, without inflammation, much pains have been taken to restore the parts to their natural state, and use, without making any wound or division at all; the introduction of a probe, the injection of a fluid, and a

F constant

* HUMULO summum ejus foraminis excipiendum, et totum id cavum sicut in fistulis dixi, usque ad os excidendum.

CELSUS.

CORPUS id quod inter angulum usque ad abcessum est excolumus et carnes e profundo educimus; quod si igitur per summa ruptus fuerit abscessus, totum id quod eminet usque ad os excidendum.

PAULUS.

SI vero per hæc medicamenta non curetur, aut recedivaret pollea, signum est quod os est corruptum de subtus quare, tunc oportet locum detegi et os corruptum removeri.

LANFRANC.

of the sacculus in the corner of the eye, are the principal means by which this has been attempted.

SOME few years ago M. Anel made a probe of so small a size as to be capable of passing from the eye-lid into the nose, being introduced at one of the puncta lachrymalia, and passing thro' the sacculus and duct; with which probe he proposed to break thro' any small obstruction, which might be found in its passage.

HE also invented a syringe whose pipe is small enough to enter one of the puncta, and by that means to furnish an opportunity of injecting a liquor into the sacculus, and duct; and with these two instruments he pretended to be able to cure the disease whenever it consisted in obstruction merely, and the discharge was not much discoloured. The first of these, viz. the passage of a small probe thro' the puncta, has a plausible appearance, but will, upon trial, be found very unequal to the task assigned; the very small size of it, its necessary flexibility, and the very little resistance it is capable of making, are manifest deficiencies in

in the instrument; the quick sensation in the lining of the sac, and duct, and its diseased state, are great objections on the side of the parts, supposing that it was capable of answering any valuable end, which it most certainly is not.

THAT the passing a fine probe from one of the puncta lachrymalia into the nose is very practicable, I know from experience, but I also know from the same experience, that the pain it gives, and the inflammation it often excites, are much greater than any benefit which does or can arise from it.

It is said that the principal use of this probe is to clear the little ducts leading from the puncta into the sacculus, and the obstruction of those ducts is often mentioned as a part of this disease, by which one would be led to suppose that it was a circumstance which frequently occurred, whereas it is seldom if ever met with, and when it does happen, can never produce the disease in question, the principal characteristic of which, is a discharge into the inner corner of the eye upon pressure made in the angle; this discharge is made from the sacculus, thro' the puncta, and proves

F 2

that

that the latter are open ; the passing a probe therefore thro' these seems to be perfectly unnecessary, since a stoppage of them would never give rise to that disease, which consists in an obstruction to the passage of any thing from the sac into the nose, and not from the eye into the sac.

THE syringe, if used judiciously while the disease is recent, the sac very little dilated, and the mucus perfectly clear, will sometimes be found serviceable ; I have used it where, I think, it has been much so ; I have by means of it injected a fluid thro' the sacculus into the nose, and in two or three instances have effected cures by it, but I have also often used it ineffectually ; it gives no pain, and a few trials render the use of it very little troublesome.

FABRICIUS ab aquapendente invented an instrument, which was so contrived, as by means of a screw to make a pressure externally on the lachrymal bag, from the use of which, he says, his patients received much benefit ; this instrument has been considerably improved by late practitioners, and is still recommended as very useful.

ALL

ALL the good that can be obtained by compress, and bandage, this screw is capable of procuring; but it is also subject to all the same inconveniences, arising from the impossibility of determining exactly the due degree of pressure, for if it be so great as to bring the sides of the upper part of sac into contact, all communication between it and the puncta will be thereby stopped; if it be but slight, the accumulation will not be prevented, nor does it in either case contribute to the removal of the obstruction in the nasal duct, the primary and original cause of the disease.

IF the curative intention was to procure an union of the sides of the sacculus, as in the case of parts separated from each other by the formation of matter or sloughs, and the pressure could be made uniformly and constantly, possibly it might be so managed as to answer a valuable purpose; but as that is not the intention, the pressure, whether made by an instrument, or by a common roller, and compress, contributes little or nothing toward a cure, nor did I ever see one effected by it, altho' I have several times tried both.

THAT

THAT some slight obstructions of the nasal duct have gone off while the compression has been used, I do not deny, but am in great doubt concerning the share which it had in removing them, having seen more than one instance of a cure being obtained by the use of a proper regimen and medicines, in slight and recent cases, where nothing was used externally but a vitriolic collyrium; and having been always disappointed in my attempts by mere bandage of any kind.

BESIDE these means of attempting a cure without incision, the gentlemen of the French Academy have favoured us with some others, such as, the introduction of a probe into the lower part of the nasal duct within the nose, the injection of a fluid by the same orifice, the passing a seton from the punctum lachrymale superius thro' the sacculus and duct and out at the nostril, there to remain 'till the cure is compleated; and for those purposes they have invented and given figures of a number of probes, syringes, and many other instruments which, they say, have been very successfully used; far be it from me to say that
they

trying those, or any other means by which mankind may be cured of diseases with the least possible fatigue and pain; but from the experiments which I have made of most of these processes, I must beg leave to suspend my assent to their general utility, or even to their frequent practicability.

REPEATED trials upon dead subjects will undoubtedly enable a man to pass the probe, or perhaps now and then the seton, but he will also find it often absolutely impracticable, and in the few instances in which he may chance to succeed as to this attempt, what will in general be the consequence? not what the writers on these subjects have taught him to believe, a cure, without a sense of pain, and degree of inflammation, which the patient, before such attempts were made, was free from, an exacerbation of the disease, and a loss of much time, as I have more than once experienced. To which consideration may be added, that infants and young children are very often afflicted with this disorder, and that such processes as these, are absolutely impracticable upon them.

I SHOULD

I SHOULD be very sorry to be misunderstood in what I now say, to have it suspected, that I mean to derogate from the character of those gentlemen who have been the inventors of these operations, or, that I speak slightly of them, either because they are not my own, or because I have not been able to succeed in the use of them: it would give me great concern if I thought it would be believed that I acted upon so mean, so narrow a principle; no man is or would be more pleased with any real improvement in our art than myself, but having taken all the pains in my power to apply the discoveries of which I am now speaking to practice (the only test of good surgery) and having found them most frequently impracticable, always ineffectual, I think myself obliged to say so.

ANEL's syringe I have used successfully and think it may now and then be very well worth trying, in recent cases more especially, as it may always be used without giving any pain, or running the risk of raising an inflammation; but I must also beg leave to observe, that if the bag is

skin and cellular membrane uninflamed, and the parts about soft and easy, if the patient will take care not to suffer too great an accumulation, will, by the frequent use of a vitriolic collyrium, keep the eye-lids clean and cool, and carefully avoid such things as irritate the membrana narium, or occasion a sudden flux of lymph from the lachrymal gland, the disease may for many years, nay often for life, be kept from being very troublesome, or inconvenient, without any surgery at all.—

S E C T. V.

WHEN the disease is got beyond the simple state just described, that is, when the parts round about are much, or constantly inflamed, or the skin covering the tumor is burst, there is something more to be done, if a cure is intended.

IN this state an opening in the upper part of the sacculus lachrymalis becomes in general absolutely necessary; and as a wound made by a knife leaves a much less disagreeable scar than that which necessari-

G

ly

being a mere simple division, the other
loss of substance; it will always be found
best to anticipate the accident of bursting
by making the opening, as soon as the in-
tegments are in such a state as to threaten
it.

FOR the making this incision authors
have been very particular in their direction
with regard to its place, manner, and form;
they have ordered it to be semilunar, hav-
ing its concave part toward the eye, and
that the point of union of the lids should
be exactly opposite to the center of the in-
cision; this lunated figure was calculated
to correspond with the course of the fibres
of the orbicular muscle, upon a supposition
that a transverse section of them, would
produce an inversion of the lower lid, an
effect which never follows; all that the
surgeon need observe, is to take care to
keep the knife at a proper distance from
the juncture of the palpebræ, to begin the
incision a very little above a line drawn
from that juncture toward the nose, and to
continue it downward; its form may full
as well be straight as any other, and the best
instru-

looked history.

IF the facculus is already burst, the place of opening is determined, and the orifice may be enlarged with a knife, or dilated.

THE incision made, the facculus should be moderately distended, either with dry lint, or a bit of prepared sponge; by which means, an opportunity will be gained in two or three days of knowing the state of the inside of the sac, and of the ductus nasal; if the former is neither roughy nor otherwise diseased, and the obstruction in the latter but slight, it sometimes happens, that after a free discharge has been made for some days, and the inflammation occasioned by the first operation is gone off, that the sac contracts itself, the superficial dressing, with moderate pressure, heals the fore, the lachrymal fluid resumes its wonted course, and the disease disappears.

OF this I have seen more than one instance, and perhaps it would happen oftener, if the very absurd manner in which

ing the bag, did not prevent it; in this state success is to be expected from the most gentle treatment only; whatever irritates inflames, or destroys, will infallibly prevent it.

If this simple method does not succeed, or from the state of the parts seems unlikely to do so, another must be tried, which the opening already made will enable us to put in practice: the point to be aimed at is, if possible, to render the nasal duct pervious to the lachrymal fluid; and we must endeavour to obtain this end by such means as give the least pain, excite the least inflammation, and leave the parts as near as possible in their natural state, that is, we are to endeavour to dilate the passage from the sac to the nose, by some means which will gradually distend it without destroying its texture, in the same manner as the dilatation of the urethra ought to be effected in the case of strictures, by passing either a probe, or a piece of cat-gut, or a bougie, gently into it, as far as it will easily go, and repeating it
occasi-

occasionally, until it is got quite through, and the passage is free *.

EVERY man will determine for himself, by what means he will endeavour to accomplish this end, nor is it of very material consequence which he prefers, provided it be done gradually, and without giving pain; a proper dilatation of the upper part of the sacculus by dry lint, or a bit of prepared sponge, will be found useful previous to the attempt toward passing any thing into, or thro' the duct; and it will also be necessary that the surgeon be possessed of a just idea of the size and direction of it, both in a natural, and a diseased state; for whoever has formed one only from viewing its bony channel in a dry skull, will upon experiment find himself much deceived with regard to its diameter in a living subject; the membrane which lines it is not extremely thin in a healthy state, and when it is inflamed or thickened by obstruction, the passage thro' the duct is thereby

* THIS caution is very necessary to be observed in the cure of strictures of the urethra, in which case the proper intention is gradually to dilate the passage, and to procure an increased discharge of mucus from the lacunæ; this should always be done gently, and by means which give as little pain as possible, whatever irritates or gives pain will certainly do mischief, will add to the obstruction, and increase the dysury.

quiere rendered very painful, if it is not quite shut up.

THEY of our ancestors who mistook this disease for an abscess, and found (as indeed they always must) extreme difficulty in filling it up with sound flesh, generally had recourse to escharotic medicines for the destruction of that fungus which seemed to hinder them from accomplishing their end; by which conduct they irritated all the neighbouring parts, increased the inflammation, and were most frequently frustrated in their expectation of a cure at last. The same kind of medicines were also used by those who supposed the disorder to be an encysted tumor, with intention to eradicate the cyst, which, they thought, prevented a cure by remaining behind; and both these methods of practice were vindicable, supposing their idea of the disease had been a true one, which it most undoubtedly was not; their reasoning was right, but their principles were wrong; they were in general very little acquainted with the structure and use of the parts, and totally mistook the nature of the disease.

BUT

BUT now, that we are thoroughly acquainted with both, this kind of practice ought surely to cease, as the preservation of the sacculus and duct, and not their destruction, are, or ought to be, intended: all cathæretic medicines must be wrong and prejudicial, at least while the intention is such; an intention at all times rational, and sometimes capable of being fulfilled.

NOTWITHSTANDING the destruction of the bag is allowed to be wrong by most surgeons of the present time, yet there are many, who, by their manner of dressing it, after they have opened it, do really, tho' not intentionally, produce the same effect as our forefathers aimed at: it is still a custom with many, as soon as it is opened, to distend the cavity of it with a hard tent, or with dossils of lint charged with escharotic medicines, such as mercurius precipitatus ruber, &c. by which means the inflammation is increased, the skin and edges of the incision hardened, and the inside of the sacculus put under a necessity of casting off a slough. This is one of several instances still remaining of our adhering to old methods of practice, after the principles

many formed have been allowed even by ourselves to be erroneous ; for this manner of dressing the fore is effectively the same as the antients made use of, while they supposed the disease to be an abscess of the caruncle, an encysted tumor, or a callous ulcer with carious bone ; and was by them intended very properly for the destruction of such callosity, to assist the exfoliation of the supposed caries, and to procure a firm basis to incarn upon.

ON the contrary, the point which ought first to be aimed at, immediately after having made an opening into the sac, is to endeavour to remove the obstruction of the natural passage from thence into the nose, by the means already mentioned, which design this method of cramming in escharotic dressings must necessarily frustrate, must frequently render a simple case complex, and at least retard that cure it is designed to expedite.

THE only excuse that can be now made for such method of dressing, is that the surgeon is satisfied that the ductus ad nares cannot be restored to its use, and therefore
by

as may fill up its cavity, and hinder the accumulation, or lodgment there in future.

IF this was feasible, perhaps it might be a vindication of such treatment, but unfortunately it neither is, nor can be so in general; and whoever will attentively examine the natural situation and structure of the parts concerned, will immediately see why it cannot. All, or the greatest part, of the diseased and obstructed duct, lying in its bony channel out of the reach of what is applied to the inside of the sacculus, must prevent the generation of a firm basis at its bottom, and produce a fresh collection of mucus, which in a short space of time lifts up the cicatrix, into a new tumor, and requires the same treatment as if nothing at all had been done.

ON the other hand, it must not be denied, that now and then a cure has by this means been effected, but it has been so rarely, that it can hardly be admitted as an authority or vindication of so irrational an attempt.

H

THE

them of very quick sensation, and easily irritated; and dressings are in fact extraneous bodies, and therefore when applied to such parts cannot be too soft and light: suppuration is an act of nature, not of art, and is always best executed, when she is least disturbed: this is a general truth, and will hold good in all parts of the body, even where suppuration may be most wanted; but in the present case, in which the lower part of the sac, and all the duct, are often in such state as not to require any suppuration at all, escharotic dressings of any kind, by producing inflammation both of the eye and caruncle, by rendering the edges of the fore hard, or sloughy, and by destroying the communication between the puncta lachrymalia and sacculus, must necessarily counteract the only proper intention of cure.

I would not in this place be thought to mean that a mere superficial pledgit is all the dressing that is required; no; a moderate dilatation of the upper part of the sacculus is at first absolutely necessary, in order to get easily at the duct below; but
this

corroliive applications of any kind, and is best accomplished by prepared sponge, which will distend to almost any degree, without destroying.

WHEN a passage has been once obtained, it should be carefully kept open, either by a piece of cat-gut, a small bougie, a leaden probe, or something of that sort ; and when it is thoroughly established, the fore may be permitted to contract, until it becomes no more than what serves for the introduction of the bougie into the duct ; in this state I would advise, that it be kept open for some time, injecting now and then a little aqua calcis, softened with mell. ros. thro' from above into the nose ; and when it appears, that the passage is so free, and so well established, that there is good probability of its preserving itself, the orifice in the angle of the eye, by being covered only by a superficial bit of plaster, pledgit, will contract and close ; and if, during its closing, moderate pressure be laid on the sacculus, to prevent a fresh accumulation of mucus, it will assist the cure.

WHETHER the sacculus in a healthy and undilated state, is endued with any degree of contractile power, which it loses by being distended, or to what other cause it may be owing, I know not; but have more than once been foiled in my attempts toward this method of curing the disease by a fresh collection of mucus, notwithstanding the nasal duct has remained open as appeared by the discharge made into the nose, upon pressure on the tumor, the immediate subsidence of the said tumor, and the passage of an injection, or small probe after having again opened the sac. Some of these have, upon being again healed, remained good cures, and others not; the uncertainty which attends these cases is great, and the event never to be known but by experiment. Whoever says, that none of them are to be cured by the foregoing method, errs as much as he would, who should expect it to succeed in all; where the disease is in such state, as to admit its being tried, it is very well worth while, as it is not painful, nor tedious; and where it does not answer our expectations, it is no hindrance to any other more efficacious one, being made use of afterward;

stances in the patient, or in the state of the diseased parts, must produce a variation in the necessary treatment, both in general, and particular; a bad habit will require the use of internal remedies; the combination of other diseases of the neighbouring parts will add to the difficulty and trouble; and even the fairest, and such as seem most likely to succeed, do sometimes resist this, and indeed every other attempt.

FROM the necessity of keeping the eye bound while dressings are applied for the dilatation of the sacculus, an inflammation is frequently raised; this added to the necessary discharge of serum, mucus, &c. is apt to heat and excoriate the parts about; therefore, warm fomentations, cooling collyria, epulotic cerates, and renewing the dressings as often as shall be necessary, with whatever else can contribute toward keeping the skin clean and cool, must be found serviceable, as well as pleasant, and should never be neglected.

S E C T.

THE last state which I mentioned of this disorder, is that in which the natural passage from the sacculus to the nose is so diseased as to be quite obliterated, or in which the bones are sometimes found to be carious.

THE methods hitherto described have all been calculated to preserve the natural passage, and to derive the lachrymal fluid again thro' it; in this attempt they are sometimes successful, but when they are not, there is no chirurgical means left, but to attempt the formation of an artificial one in its stead.

THE upper and hinder part of the sacculus lachrymalis is firmly attached to the os unguis, a small and very thin bone just within the orbit of the eye; which bone is so situated, that if it be by any means broke thro', or removed, the two cavities of the nose, and of the orbit, communicate with each other, consequently the os unguis forms the partition between the hinder part of the lachrymal bag, and the upper part
of

making a breach in this partition that we attempt the formation of an artificial passage for the lachrymal fluid.

THIS operation, if considered merely as a perforation, is no invention of the moderns, the ancients undoubtedly performed it; but tho' it was executed much in the same manner, as it is now, yet it was not done with the same intention.

FROM the accounts which our ancestors have left us of the disease in question, it is plain, that they supposed it to be always attended with a degree of callosity, and often with caries, and that the surest way to obtain a cure was to lay the bone bare; this they effected, either by caustic or cautery, according to the humour of the surgeon, or the fears of the patient; if caustic applications were used, they waited the separation of the eschar; and if they found, or believed the bone to be altered, they applied an actual cautery to it; if the bone which the iron was applied was the os maxillaris, it was too thin to bear much heat, much pressure, consequently was easily burnt, or broke thro', and by that means an opening

was also sometimes made use of instead of cautery, and the same effect produced thereby *.

By each of these methods, a passage being made from the *sacculus lachrymalis* into the nose, a cure was sometimes accidentally obtained; but the cautery was applied, either to destroy the supposed callosity, or to desquamate a caries; and the *terebra*, either for the same reason, or to make a passage for the discharge of matter, which lodged, and as they thought, hindered the healing of the sore; for as they were not acquainted with the natural passage of the lachrymal fluid, it would be absurd to suppose,

* *Oculo et cæteris junctis partibus bene obtectis os fermento adurendum est vehementius, quod si jam carie vexatum est, quo crassior huic squama abscedat quidam adurentia imponunt.*

CELSUS.

Cum isto pulvere in veritate fere mortificabam omnes fistulas curabiles, et cum cauterio ferreo, aut æneo—— factâ mortificatione tali totius carnis usque ad os, cum pulvere aut unguento superdictis superpone, mortificato butyrum et escharâ aspice et si fuerit os corruptum cauteriza ipsum usque ad ejus profundum.

GUL. DE SALICETO.

Postea si homo fuerit delicatus, per istud foramen mittatur *Canellus ferreus* vel æneus subtilis usque ad profundum si poteris, et per ipsum canellum ferrum candens immitte et fistulæ radices decoque: at si timuerit ignem immittatur pillula de unguento ruptorio.

ROLANDUS.

Osse detecto ferrum imprime calidum supra ipsum et ipsum cauterium mediocriter comprimendo postea imple totum vulnus cum oleo rosarum misto cum vitello ovi.

one. Callosity and caries were their two characteristics of the disease; the dissolution of one, and the exfoliation of the other, were all they had in view from the use of either caustic or cautery, and the perforation of the os unguis was either accidental, or made merely for the discharge of matter *.

I

INDEED,

* FABRICIUS AB AQUAPENDENTE who in general copies Paulus, speaks of the perforation as meant only to make a depending orifice for matter, "post carunculæ et loci excisionem, terebra humorem aut pus in nares derivarint."

FAB. AB AQUAPENDENTE.

GUL. DE SALICETO, and indeed many other of the ancient writers, speak of using both cautery and terebra to the purpose of deriving the matter and sanies which lodge in the nose; and, by making a depending orifice, procure a firm basis to heal on. "Aspice os, et si fuerit corruptum cauteriza ipsum usque ad ejus profundum, et concavitatem cum cauterio punctuali, et perfora ipsum ad aliam partem, ejus ut sanies per nasum fluat, deinde incarnetur et consolidetur."

GUL. DE SALICETO.

Indeed the formation of an artificial passage for the laryngeal fluid could make no part of the intention of those, who were not rightly acquainted with the natural one.

PAULUS mentions perforation with the terebra as the practice of some in his time, but from what he says, it is plain he did not practice it himself, or think it necessary; and that he regarded it only as a method of making a depending orifice; his words are, "Quod si jam carie vexatum est ferro candenti, acuto, ac in cuspidem abeunte adsumimus spongiâ frigidâ madente oculo imposita."

SUNT qui post carunculæ excisionem terebra usi humorem aut pus in nares derivarint; nos autem satis habuimus eousque solum fermentis ad Ægyptopem accomodatis adurere ut squama abscederet."

PAULUS AGINETA.

SEE also FAB. AB AQUAPENDENTE.

the old writers have left us on this subject it will appear, that tho' they knew that : passage into the nose was sometimes a consequence of their use of the terebra and cautery, yet the operators had no very accurate knowledge of the parts they made so free with ; no precise idea of the bone on which their instruments were applied, or thro' which they passed ; nor of the place most immediately proper, for such application of them ; sometimes they perforated the os unguis very properly sometimes the cautery or terebra was thrust into the bony channel of the natural nasal duct, and sometimes they were applied to the nasal process of the maxilla superior : the direction given by most of them to rasp the bone (*scalpris abradere*) and to impress the cautery with some force, that the bone may be sooner exfoliated, (*ut citius squama abscedat*) plainly prove that either they were not aware of the tender structure of the os unguis, or that they did not intend to apply their instruments to it ; if the former was the case, the perforation was accidental ; if the latter, they must have often done much more harm than good ;

destroyed unnecessarily parts which have little or nothing to do with the disease; and by such treatment of them must have much oftener prevented, than accomplished a cure *.

THE intention of the present practitioners in making this perforation is different from that of their ancestors; but it is more rational, and founded upon the nature and use of the parts concerned in the disease; it is to form and maintain a new artificial passage from the lachrymal bag into the nose, when the natural one can no more be rendered useful, and without any view to any thing else; this, I say, is the aim of them all; but tho' they are perfectly agreed in their intention, yet they are not so with regard to the instrument which they use, some still continuing the actual cautery,

* PETRUS DE MARCHETTI, tho' perfectly sensible that the os unguis was often broke thro' by the cautery, yet insists upon it, that it served no other purpose than to hasten exfoliation, "*Præterquam quod hujus perforationis non alius sit usus quam ut os perforatum aut inustum citius abscedat. Observandum tamen non esse perforandum os nisi præsentē maxima ipsius corruptione, sola siquidem ejus superficie corrupta aut alterata sit fuerit partem læsam abradere.*"

PETR. DE MARCHETTI.

AND Mr. Verdue, a very modern writer, is also of the same opinion, "*Le meilleur remede pour amorter l'acide qui cause la carie, c'est de passer légèrement un cautère actuel sur l'os sans le percer.*"

THE ancients preferred the cautery, for reasons which have already been assigned ; but since the symptoms of callosity and caries have been found to be very infrequent, and the os unguis has been perforated solely with a view to make an artificial passage into the nose, the cautery has with many lost much of its ancient credit, and other instruments have been substituted in its place, which give less pain at the time of using, and leave less deformity afterward.

BUT tho' many have laid aside the hot iron, yet it still has its advocates, who prefer it to every other instrument, and who have therefore endeavoured to obviate its inconveniences ; they have directed that the cannula thro' which it passes, be made of a conical form, and so large at its lower end, as that they shall not touch each other ; they have ordered this cannula to be wrapped round with wet rag, at the time of using it ; they have placed a check upon the top of the iron to prevent its point from going too far, and have been particular

soon as it is got thro'.

But notwithstanding these and every other caution, the cautery gives great pain at the time of using, lengthens the attendance, and most commonly produces unnecessary deformity, even in the hands of the most dextrous, not to mention the horror occasioned by thrusting a hot iron into the corner of the eye.

WHEN the inconveniences arising from the use of this instrument, even in the best hands, are important, it may be easily guessed what they must be in those of the clumsy and ignorant; and therefore unless some real advantage attends it, it ought certainly to be so discouraged, that no one may attempt to revive it; let us then see with what intent it has been used by those who have appeared most fond of it, and who may fairly be supposed to have best known how to manage it.

THE defence made by the wet rag against the heat of the iron, the disproportioned size, and the figure of the cannula, very plainly shew, that its effect is designed

the check at the upper as clearly shews, that that point is designed to pass no farther than just thro' the bone, while all the ill effects are occasioned by the upper part of the cautery on the eye-lids and angle of the eye; now, if it is not designed to produce any effect on any of the parts thro' which it passes down to the bone, but merely to burn thro' that and the membrana narium, and thereby make an opening into the nose, I do not see how it differs from any other perforator of equal size, except in the mischief it does to the parts above, to which it should do nothing.

It does indeed burn the bone and membrane, thro' which it pierces, and thereby prevents the orifice from closing again immediately, and this is certainly the principal end of perforation, by whatever instrument it is performed; but it is also as certain, that the same end is obtainable by means less mischievous and less horrible.

Our ancestors had a very plausible reason for using it, their ideas of callosity and caries always accompanied this disease, and authorised them to make use of such applications

these are symptoms which very rarely occur, or even if they do, that they are removable in a much easier manner, we are no longer vindicated in continuing an alarming and a painful process, when we can obtain the same end by much gentler means; for whether the membrana narium be burnt thro', or divided in any other manner, it is the future method of dressing that opening that must maintain it, let it be made by whatever instrument, or in whatever manner it may.

THE late Mr. Cheffelden was a warm patron of the cautery, took a good deal of pains to prevent it from doing mischief, and has said in its defence, that—"other methods of curing this disease have been much recommended, though often unsuccessful, but this well performed is infallible." After so positive an assertion, I am sorry to be obliged to say, that it is contradicted by manifold experience, that there have been many instances of perfect cures performed without the use of a cautery; and that some of those who have been cauterized by Mr. Cheffelden himself have
been

took, prevent the effect of the heat of the iron, or leave his patient without a weeping eye.

THE intention is merely to make an opening thro' the os unguis and membrana narium into the cavity of the nose, and to treat that perforation in such a manner as that it shall most probably remain open, and give passage to the lachrymal fluid from the puncta, after the external sore is healed.

The extreme thinness of the bone renders the passage of the instrument very easy, and if the breach which is made, be of any tolerable size, I am inclined to think that it never is filled up again by bone, but that when it is closed, it is by the membrane; and therefore it is the surgeon's business to make a pretty large opening in the bone, and to prevent its being closed again, by rendering the edges of the membrane on each side of it callous.

To make this opening, many different instruments have been devised, and used,
a large

each of which, if dextrously and properly applied, will do the business very well; the one necessary caution is, so to apply whatever instrument is used, that it may pierce thro' that part of the bone which lies immediately behind the sacculus lachrymalis, and not to push it too far into the nose for fear of injuring the os spongiosum behind, while it breaks its way.

For my own part I have always used the curved trochar, which has served my purpose well, and from which I have never experienced any inconvenience; in using it, the point should be turned obliquely downward, from the angle of the eye toward the inside of the nose; the accomplishment of the breach will be known by the discharge of blood from the nostril, and of air from the wound upon blowing the nose. The most precise direction in this part of the operation will be of but little use to him who has no idea of the natural structure and disposition of the parts concerned, and who ought therefore to get such information as soon as he can; but whoever is at all acquainted with this matter, or will

K

attend

ridge, that the lachrymal sac is connected to all that part which is anterior to this ridge, and that the posterior part of the bone contributes to form the orbit of the eye, and has little or no connexion with the lachrymal sac; the trocar must be applied therefore to that part of the bone which is anterior to the ridge, and consequently behind the lachrymal bag; by the passage of the instrument, all this part of the bone will in all probability be broken, but from which no mischief will ensue.

AN attention to the natural situation of these parts will also show the practitioner, that if the point of his instrument be passed in a transverse direction with regard to the nose, the os spongiosum superius will be unnecessarily wounded or broken, and if it goes in too perpendicular a direction, it may get into the channel of the natural duct, and its point will be stopped by bearing against that part of the maxilla superior which contributes to the formation of that channel.

it may break the os unguis to some distance from the place where its immediate point is fixed; to which I can only answer, that I have performed the operation a great number of times, and never yet have seen any inconvenience to arise from it; indeed a total removal of a small piece of the bone would be a thing rather to be wished for than avoided; if we may reason by analogy, it seems to be a necessary requisite toward preserving a future passage, for we very well know in a caries of the bones forming the roof of the mouth that they are sometimes bare for a large compass, and by casting off leave a considerable aperture into the nose, yet in many cases, when the virus is removed, and the habit recruited, that opening will so contract as not to suffer a small quill to pass where you might have introduced your finger, nay often will quite close, and therefore tho' the opening made in the os unguis may possibly in spite of all endeavours be again closed up, yet a free breach in it seems to be the most likely means to prevent it, and upon this principle I have always turned the perforator round very freely whenever I have used it, have never seen any mischief from it,

with it, in some measure to this method of using it.

As soon as the perforation is made, a tent of lint should be introduced, of such size as to fill the aperture, and so long as to pass thro' it into the cavity of the nose; this should be permitted to remain in 2, 3, or 4 days, till the suppuration of the parts renders its extraction easy, and after that a fresh one should be passed every day, until the clean granulating appearance of the sore makes it probable that the edges of the divided membrane are in the same state; the business now is to prevent the incarnation from closing the orifice, for which purpose the end of the tent may be moistened with spir. vitriol. ten. or a piece of lunar caustic so included in a quill, as to leave little more than the extremity naked, may at each dressing, or every other, or every third day be introduced, by which the granulation will be repressed, and the opening maintained, and when this has been done for some little time, a piece of bougie of proper size, or a leaden cannula may be introduced instead of the tent, and leaving off all other dressing, the sore may be suffered,

will permit, which should be of such length that one extremity of it may lie level with the skin in the corner of the eye, and the other be within the nose.

THE longer time the patient can be prevailed upon to wear the bougie, the more likely will be the continuance of the opening, and when it is withdrawn the external orifice should be covered only by a superficial pledgit, or plaister, and suffered to heal under moderate pressure.

THERE is another method which has been much recommended by some French writers to prevent the closing of the opening in the os unguis, which is to introduce a cannula either of gold or silver, or lead into the aperture ; and to permit the fore to heal over it, suffering the cannula to remain, or to come away by the nose.

For my own part I cannot say any thing to it, having never had occasion to try it, the cases of this kind which I have had under my direction, having generally succeeded under some of the methods already mentioned, which methods
will

given is clear in his intention, pursues it steadily, and properly, and refrains from doing too much, tho' I must again repeat what I have said before, viz. that there is no method of treating this disorder which is infallible, none that will absolutely and in all cases prevent a return, especially in scrophulous habits: yet when a just distinction is made between those cases which are in their own nature incapable of cure, and those which by being improperly treated are not cured, I am inclined to believe that the number of the former, will be found much smaller than it is generally imagined to be.

F I N I S.

Lately Published,

Sold by L. H A W E S and Co.

WRITTEN BY

P E R C I V A L L P O T T,

Senior Surgeon to St. Bartholomew's Hospital,

All neatly printed in Octavo.

1. **A** TREATISE on Ruptures in general. Price 4s.
2. An Account of a particular Kind of Rupture, frequently attendant upon new-born Children, and sometimes met with in Adults. Price 1 s. or bound up with the foregoing, Price 5 s.
3. Practical Remarks on the Hydrocele or watry Rupture, and some other Diseases of the Testicle, its Coats and Vessels; (illustrated with Cases :) Being a Supplement to a general Treatise on Ruptures, published in the Year 1756.
4. Observations on the Nature and Consequences of Wounds and Contusions of the Head, Fractures of the Scull, Contusions of the Brain, &c. Price 3 s. 6 d.

THE UNIVERSITY OF CHICAGO

TO THE PRESIDENT

OF THE UNITED STATES

AND TO THE SENATE

OF THE UNITED STATES

IN RESPONSE TO A RESOLUTION PASSED BY THE SENATE
ON FEBRUARY 2, 1940

AND TO THE HOUSE OF REPRESENTATIVES
ON FEBRUARY 2, 1940

AND TO THE SENATE
ON FEBRUARY 2, 1940

C A S E

O F

Sir JEREMY SAMBROOKE,
BARONET,

Fairly and Impartially STATED.

CONTAINING

A full and exact Account of the Singular
DISORDER in the EYE of this GENTLE-
MAN.

WITH

A faithful Relation of the great Variety of un-
common CIRCUMSTANCES attending his
CURE.

A N D

A clear Demonstration of the highest Probability of
his PERFECT RECOVERY.

Humbly Inscrib'd to HIMSELF.

By J. TAYLOR, Doctor of Physick; Oculist to his
Majesty; and Member of several of the Colleges of Phy-
sicians in Foreign Parts.

Qui dat Videre, dat Vivere.

L O N D O N :

Printed for J. CROCKATT, near Garraway's Coffee House in
Exchange-Alley; and Sold by M. COOPER, at the *Globe*,
in *Pater-noster-Row*. 1743.

[Price SIX-PENCE.]

1650240

TO

Sir Jeremy Sambrooke, Bart.

S I R,



VER since I first had the Honour of being consulted, and employed, in a Case, on which your Welfare and Happiness so immediately depended; as the good Opinion you expressed of my Judgment, and the great Confidence you have all along reposed in my Integrity, gave me the utmost Delight and Satisfaction; so they failed not to excite my best Endeavours, and encouraged me to try, by the strictest Application, extraordinary Pains, and unwearied Diligence

to be an Instrument of Good to a Gentleman of your Consequence and Merit, in a Matter of no less Importance, than the Restoration of that unvaluable Blessing of Sight.

INCITED by the Hopes of obtaining so desirable an End, I enter'd on the Work with Pleasure, and proceeded in it with Assiduity and Circumspection. Nothing was left untried, that Reason could dictate, or Art direct, in this so nice, and important an Affair. And Heaven had so far blessed my Undertakings, and succeeded my Endeavours, that I had happily surmounted the most material Difficulties, so that the Good I pursued was now in View; and I had, not without Reason, entertained

I had so successfully brought to this happy State, I should shortly see accomplished with equal Felicity ; namely, that You might reap the Benefit, and myself the Pleasure of having all my Labours crowned with a prosperous and good Event.

JUDGE then, Sir, if I must not be very sensibly touched at the Disappointment of all my well-grounded Expectations ; a Disappointment that affects me with less Sorrow on my own Account than Yours. For though I cannot but regret the Loss of that Pleasure and Reputation, I had so nearly acquired ; yet, as I am conscious of having taken such Measures, as all impartial Judges must approve, and would, I am persuaded, if continued,

med. Effects, I am but rightly concerned, with respect to myself, and nothing gives so much Pain, as that I am hindred from compleating the Happiness of a Gentleman, whose Worth and Goodness make him so deservedly Esteemed, and to be conducive to whose Welfare, was the Summit of my Wishes, and the utmost Height of all my Ambition.

THAT my Hopes of Success were not without Foundation, I presume to have fully made appear in the following Sheets, to every Man of Understanding and Probity. I have sufficiently justified all my Proceedings to the *Skillful*, who, as they cannot but see, how nearly the Cure was effected, and how easy it is still to compleat

compleat it, must therefore be sensible of the great Injury you would suffer to have it obstructed, or so much as retarded. And those amongst them, who have the Happiness of knowing You, must undoubtedly conceive just Indignation against any, who shall be so cruel and wicked, as to deter You from the Enjoyment of that Blessing, which Providence has in a Manner put into your own Hands, and seems as it were, to Court your Reception: And of which, you cannot be now deprived, without the greatest Injustice, and basest Inhumanity.

THAT you may still live to enjoy the Blessing of Sight, and every other Advantage, that can contribute to make your Life Happy, and render

der the Continuation of it as pleasant
to your self, as it is serviceable to
great Numbers of Others ; is the sin-
cere Desire, and fervent Prayer, of,

S I R,

Your most obliged,

Humble Servant,

*Great Queen-Street,
Lincoln's-Inn-Fields,
October 1, 1743.*

JOHN TAYLOR.

THE



THE
C A S E
OF

Sir Jeremy Sambrooke, Bart.

Fairly and Impartially Stated, &c.



S the Case of *Sir Jeremy Sambrooke*, has been the Subject of much Conversation, as well from the Singularity of the Disorder, and the great Variety of extraordinary Circum-

stances attending the Cure, as from the raised Expectation of a great Number of Persons of Worth and Distinction, peculiarly interested in the good, or ill Success, attending a Gentleman of such universal Estimation, and exten-

B

five

five Merit; whose many amiable Qualities render him so deservedly Dear to all who have the Honour of his Acquaintance, and who, consequently cannot but be very sensibly affected, with Pleasure or Uneasiness, proportionable to the happy or unhappy Event of an Undertaking, in which his Welfare is so immediately concerned: I was induced to think, that a full and particular Narrative of my whole Proceeding, in the Treatment of this Gentleman's Indisposition, would not be unpleasing to the Public; especially, as it is given by his own Permission, and published with no other View, than that of justifying all the Measures I have taken, and demonstrating, that a further Pursuit of them can hardly fail, of being attended with a happy Recovery of useful Vision.

ABOUT the begining of *June* last (in Consequence of the good Success, I was so happy as to obtain in a parrallel Case, with the Lady his Sister, and several Others *, whose Recovery had been enquired into, and known by some of that Honourable Family) I was consulted concerning a Disorder in the left Eye, under which this
Gentleman

* Mrs. Hooper, of Tunbridge. Mr. Crosby, of Bristol. Mr. Hammond, of Hatton-Garden, &c.

UPON Examination, I found the Cryſtalline totally Opaque, with all the Appearances of that Solidity in the Opacity; which conſtitutes, what is vulgarly term'd, *The Maturity, or Ripeneſs of the Cataract*, not without the ſtrongeſt Proofs, that can be, of the healthful Senſibility of the Immediate Organ of Sight, namely, the Mobility of the Pupil, and the Senſation of Light.

ENCOURAG'D by theſe ſtrong, preſumptive Arguments of Succeſs, I concluded (which I believe every one will allow to be a reaſonable Inference) That upon Depreſſing the Opaque Cryſtalline, and thereby recovering the Transparency of the Axis of the Globe, ſuch uſeful Sight, as is uſual in the like Caſes, would undoubtedly be reſtored.

UPON Enquiry made into the Patient's general Temperament, the chief Thing worthy of Obſervation, was, that he had for a Number of Years been afflicted with a Chronical Pain in the back Part of his Head. This one Objection would certainly have over-ballanced all other Encouragements, and have deterred me, from attempting

B 2

the

of Sight, from these unexceptionable Arguments already mentioned, which I have never experienced once to fail, amongst that great Number of Subjects, and variety of Circumstances, which I have had the Opportunity of treating in Cases of the same Nature.

NOTWITHSTANDING all these good Appearances in the Gentleman's Favour, the CrySTALLINE being depressed after proper Preparations, with great Facility, and all imaginable Success, in the Operation; to my extraordinary Surprise, I found, that on restoring the Axis of the Eye to its natural Pellucidity, he did not experience the Sensation of any additional Light, as might reasonably have been expected, from what had attended the Lady above-mentioned, and Numbers of Others in the like Circumstances.

EARLY in the following Morning, I found some Symptoms, which indicated a Return of the wonted Head-ach, and upon examining the Globe, I perceived the Axis to be transparent, and yet the Eye insensible of the Impression of the Object: I was now convinc'd, that this want of Sight, was owing to a Defect in the Brain,
obstruct-

more resolved to cut the Artery; judging this threatn'd return of his usual Pain, to be at least attended with an extraordinary Distention of the Blood-Vessels in the Brain: This prognostic proved just; for upon Dividing the Temporal Artery, the Patient was immediately freed from the Symptoms of his wonted Pain, and in a few Days after, his Eye had a Sensibility, (though confus'd) of the Impression of the Object.

It is very remarkable, that the Seperating the Artery, (though attended with so happy an Event,) was notwithstanding the only Source of all those succeeding Interruptions, in the Progress of the Cure; and the sole Reason, why this Gentleman, yet stands in need of a little more of my Assistance for a perfect Recovery of useful Sight; and all this from the indispensable Necessity, we lay under, of not admitting the Application of a general Bandage; lest the Return of the Head Ake, might unhappily deprive us of that Degree of Vision, which we had already obtained, and which there appeared such high probability of Improving to a great Perfection.

THE

on which at the wound; which put me under the necessity of contriving such a Bandage, as might make a more immediate Compression, on the Part, without affecting the whole Head, (not being willing to apply the Bandage usual in such Cases; for the Reasons abovementioned;) this was not followed with its desired Effect; for the next Day the Artery bled afresh: In short, being still unwilling to venture the usual Bandage, several Bandages were contrived, and successively applied, all tending to the same End, namely to compress the Part, without, affecting the whole Head; but Maugre all our Endeavours, for upwards of twenty Days; (during which the Patient underwent a great Number of Bleedings at the Wound;) nothing was Effectual, 'till at length we were reduc'd to the sad Dilemma of exposing the Gentleman to the loss, either of Sight, or Life: In this Case; there was no room for Deliberation; I was necessitated to apply the usual Bandage, which had, as far as regarded the wounded Artery, all desirable Success, but with respect to the Eye, was attended with Consequences of a very different Nature; the particulars of which I am now to relate.

Not

Not long after the Application of this Bandage, the so much dreaded ill Effects appear'd; the Patient's Head began to be afflicted with its usual Pain, and his Eye attended with all the Symptoms, that indicate an approaching Inflammation. On Inspecting the Eye, the second Day after the Application of the Bandage, I found a small Quantity of Matter collected in the lower Part of the Cornea, between its Interior Pellicles, not much unlike that, to which (from the Resemblance it bears to the White of the Root of a Human Nail) has been given the Name of Onyx. This was a very formidable Symptom, for it Incited, an extraordinary Pain, not only in the Nerves of the Immediate Organ of Sight, by the Preternatural Plethora of the Globe, but also in those Parts of the Brain, from whence these Nerves take their Origin. In this Situation, I well knew, that no Cessation of Pain could be expected, till either the Cause was remov'd by Discharging this Extraneous Matter from between the Pellicles of the Cornea, or till the Nerves of the Immediate Organ of Sight, had lost their Sensibility by the continuance of the Pressure: But this Operation, I durst not then Attempt, by Reason of the attending Inflammation,

mation, the Cause of which I was compelled to permit, least by removing the Bandage, the Bleeding of the Artery should be repeated ; and had I performed it, during the Continuation of the Bandage, and consequently of the Inflammation, it would be of no Use, as its return, after a short Time, would have been unavoidable. I therefore chose, rather than to permit a total Loss of Sensation in the Nerves of the Immediate Organ of Sight (an Effect which would infallibly have happened, had the Pressure continued long in the same Degree) to have Recourse to general Evacuations, in order to hinder, as much as possible, the extraordinary Supplies to the Arteries, in and about the Globe, consequent on the Pain incited by the Pressure : These Evacuations had their good Effect ; for though it could not be supposed that by them, any Part of the Matter could be discharged ; yet, as this Method mitigated the Inflammation, the Augmentation of the Matter was intirely prevented.

IN a few Days after, I found this Matter, (though not at all encreased in Quantity) had forced its Way, (a Thing not unusual) into the Anterior Chamber of the Aqueous, consequently was no longer the Onyx, but new constituted, that
Disease

ently shewn in a Lecture on this Subject. This Transition of the Matter, gave increase to the Pain ; not from any Augmentation in its Quantity, but from the Change of its Situation ; it being now lodged, immediately upon the Iris, and indicating a Necessity, for either a quick Discharge by Operation, or the Repetition of our wonted Evacuations ; I choosed the latter, being apprehensive, that the Matter might have acquired such an Acrimony, as would corrode the Sides of the Aperture in the Cornea, and leave it irrecoverably open ; the undoubted Effects of which, must have been a continual Efflux of the Aqueous Humour ; Instances of which I have frequently met with. Nor indeed was this a needless Precaution ; for after I had alleviated the Inflammation by proper Methods, I then perceiv'd, the Matter was not Acrimonious, by the Facility with which the Wound was healed immediately after its Discharge.

I did not venture to make an Opening large enough to discharge the Matter at one Puncture ; chiefly, that I might avoid the Danger of wounding the Iris on the sudden following of the Aqueous Humour, and the immediate Change of the

C

Situation

...of this Precaution, I was well acquainted with, from my having so often observ'd, That all those direful Consequences, which so frequently succeed the common Method of Operation, owe their Origin to this important Accident : I therefore made several Punctures, for its Discharge, to the Number of near twenty at different Times, till I was so happy as to entirely clear the Aqueous Chamber of this extraneous Matter, when in its stead, there remained (as is usual enough) a few Particles of Blood, proceeding probably from the Section of some few of those Blood Vessels, which empty themselves in that Part of the Aqueous Chamber, and from whose Extremities, the Blood, from which the Matter was collected, was at first separated ; this was Discharged in the same Manner, and with equal Success, as the precedent Matter ; till I had at length left the Aqueous Chamber entirely free from any extraneous Matter whatsoever.

DURING all this Time, 'till the Matter was Discharged, I was under the Necessity of continuing the Regimen hitherto observed, to secure the Immediate Organ of Sight, from the Loss of Sensibility, which must have been the inevitable

necessary Precautions.

HERE I cannot help remarking, that there is something very peculiar and extraordinary in this Case, and what perhaps has hitherto been unprecedented; that an Inflammation should be excited by a Cause, the Permission of whose Subsistence was unavoidable; and yet, such were the Methods taken, that the Nerves of the Immediate Organ of Sight, did all along from the first Section of the Artery, and do at this present Time, still maintain their healthful Sensibility, which without these Methods, could never have been effected,

THERE is yet another remarkable Circumstance, which must not be omitted; that during the Continuance of the Inflammation, occasioned by the extraneous Matter (which for the Reasons already given, could not be sooner Discharged) the Pupil gradually lost its Mobility and Figure, 'till at last it remained nearly closed.

I need not here observe, that there are four general Things essential to the Restoration of Sight :

C 2

To

lucidity of the Axis, to enable the Medium within the Axis, properly to transmit the Light; and To maintain the Figure and Mobility of the Pupil.

T H A T we have in this Gentleman's Case, obtained the three First (which are by far the most Difficult) is indisputable; since, during the Continuance of the Extraneous Matter in the Eye, 'till after its Discharge the Pupil became too contracted to admit sufficient Light; he was able to distinguish the Figure of the Object, which he could never have done, had not the Axis of the Eye been perfectly pellucid, for the Admission of the Light reflected from every Point of the Object, and its Medium, in a state proper to Transmit it, and had not the Nerves of the Immediate Organ of Sight, at the same Time maintained their healthful Sensibility; there is therefore, nothing further required for recovery of that Sight, which is usual after the Depression of the Cataract, (which with the Assistance of Glasses, we all know, is obtained to a great Perfection) than the bare Opening the contracted Pupil, and thereby obtaining a free Passage for the Light.

HERE

Case, which must not be pained over in Silence; and that is, that the Pupil is not only Contracted, but also the whole Iris is almost equally Elevated towards the Cornea; which happened the 2d Day after the total Discharge of all the extraneous Matter from the Aqueous Chambers, a Symptom not unusual in a long continued Inflammation; but this was not a Circumstance of Consequence sufficient to make us reject the Operation, because it will by no means render the Success less certain; for whether there be, or be not, any Aqueous Humour behind the Iris, the Opening the Pupil, in the Manner I am well known to Practice, must infallibly be followed with the Recovery of useful Sight; the Reasons are, that if on the one Hand, there is Aqueous Humour behind the Iris, as I have good Reason to believe there is, I then (by an Operation, I frequently Practice, with great Facility and no Danger) Reinstate the Iris so far, at least in its healthful Position, that it shall be no Impediment to the Restoration of useful Vision; and if on the contrary, there be no Aqueous Humour behind the Iris, then it
is

perit, (as is plain from that Sight, the Eye always maintained) and the extraordinary Elongation and Convexity of the Vitreous, will compensate for the Loss of the Aqueous: and he must be but a very superficial Judge in Optics, who does not know, that this Loss of the Aqueous, in such a State of the Eye, can no way impede a sufficient Refraction of Light properly Transmitted.

I am not ignorant, that it has been thought a disputable Point, whether or no the Globe has not lost something of its healthful Figure; but that it has not, seems very Evident, since it is not easy to conceive, how any Disfiguration of the Globe could happen without such Pressure on the Nerves of the Immediate Organ of Sight, as must inevitably have deprived them of their healthful Sensibility, which I have already shewn, is not here the Case: But even giving this large Concession, in order to remove every possible Objection, and admitting, the Globe has suffered some Alteration of Figure, yet even this is no Argument against the Probability of Success in the Operation, since
we

of the Immediate Organ of Sight do retain their healthful Sensibility, and there are several Diseases of the Eyes in which the Globe loses its Figure, without any Defect of Sight.

I am sorry, that I am obliged, in this Particular, to dissent from the Gentlemen with whom I had the Honour to be jointly Consulted; and whose extensive Knowledge and uncommon Abilities no One holds in higher Estimation than myself. But as I am more peculiarly interested in the present Case, and my Reputation more immediately concerned in the good or ill Success that shall attend any Measures concerted for its Treatment, and as my whole Life has been spent in Studies of this Nature, and my Experience for a great Number of Years, been confined to this particular Practice, those Gentlemen will Pardon me, if I appear even a little warm on this important Point; since the Arguments I have offered to prove, the Globe has not suffered any Disfiguration (and even on admitting it, that the Objection can be of no great validity) appear to me very cogent and powerful. And I am so far from urging them out of any Love to Dissention, or Disputation, that 'tis with the utmost Reluctance

to the Assertions of those Gentlemen, whose good Opinion I so greatly Prize; and whose Judgment I so highly Honour; that nothing would afford me greater Delight and Satisfaction, than to have my Sentiments concur with Their's, and any Measures I shall take, so happy as to meet with their Approbation.

HAVING thus far proved the reasonableness of expecting the perfect Recovery of this Gentleman's Sight, it remains for me to explain, how this good Effect may be attained without any ill Consequence, even that of a simple Inflammation. And this will evidently appear to any impartial Judge, if it be remembered, that there is nothing required here besides the Opening the Pupil, which is now contracted by a Multitude of little Cicatrices, all insensible, as Experience Daily shows: And as this Aperture through the Globe, is not made in or about, but much below the Horizontal Line, where none of the Nerves designed for the Service of the Iris, and Uvea pass, and when we are in the Globe, we pass only through the Vitreous Humour, the Ligamentum Ciliare being, in this State of the Eye, far enough out of Danger of being wounded,

'tis impossible any ill Consequence should attend the Operation, since there is no sensible Part wounded; all this is plain, not only from Daily Examples in this Operation, but also from the well known Experiment of peircing the Eyes of Animals without the least succeeding Inflammation.

HAVING thus explained the Method, and therein proved the great Probability (I had almost said certainty) of this Operation being attended with all desirable Success, without exposing the Patient to the Danger of the least ill Consequence, I appeal to every thinking Man how unjust, nay, how Inhumane it must be, to endeavour to deter this Gentleman from embracing the happy Opportunity, and pursuing but a little longer those means which may with so much Ease, and so little Danger, restore to him the Fruition of that inestimable Blessing of Sight, of which he has been so long deprived, and without which, all other Privileges and Possessions are tasteless, and Life itself scarce worth the keeping!

IT may, therefore, be reasonably expected, that all, who are so happy as to be any way peculiarly interested in this Gentleman's Welfare,
D
fare,

sure, will be so far from discouraging such a laudable Design, or retarding its Execution, that on the contrary, they will rather use the utmost of their Endeavours to persuade him to pursue that Happiness a step farther, which stands, as it were, expecting his Reception; as there now remains no Manner of Difficulty to remove that slight Obstruction, that lies between him and the Enjoyment of that Blessing, of the Loss of which, none can be truly sensible, but they who are so unhappy as to have been deprived of it; and I think I may reasonably affirm, that every Argument calculated to dissuade this Gentleman from completing his Pursuit, after this great Privilege of Life, now so nearly overtaken, may, if it proceeds from the fear of Danger; be termed *Weak*; but if designedly offered for any other end, must be called extreamly *wicked*.

FOR my own Part, after so much uncommon Pains taken in this Gentleman's Cure, and the great sollicitude I have undergone for its happy Success, which I had now so far affected, as to have removed all the material Difficulties, that opposed, and have so little remaining, to be farther done; should I be hindered from the Prosecution of those Measures I have

their side, of being attended with their desired Effects, I should undoubtedly be very sensibly affected with the painful Disappointment, since I know of nothing on Earth could afford me greater Pleasure, than the Recovery of a Gentleman of his Consequence, and Merit: and it would be prodigiously absurd to imagine, that I should be so desirous of attempting such an important Undertaking, in which my Reputation is so visibly at stake, where it not that I have all the Reasons, I can possibly desire, (and I think I may from my long Experience, and frequent Practice, without vanity, reckon myself a tollerable Judge of their Validity) to encourage me to hope for a good and happy Event.

F I N I S.

ON *Monday* the 21st of *November* next, about five in the Evening, Dr. TAYLOR will begin a Course of Lectures on the Nature and Cure of the DISEASES OF THE EYE (at his House in *Great Queen-Street, Lincoln's-Inn Fields.*) Containing, I. An Historical Account of all that been published on these Diseases, by the most celebrated Writers, both Antient and Modern. II. Impartial Examinations, and Remarks on all they have delivered concerning their Causes and Cure. III. The Author's own explanation of their Causes, together, with a great variety of Circumstances, distinguishing those Symptoms, which properly constitute each particular Species. IV. His method of curing every Disease in each Species and State, whether by Medicines or Operations: in which the manner of each Operation, as practised by himself, is fully described, its peculiar Advantages clearly set forth, and its difference from other Methods, truly demonstrated and justified. All founded on a long Experience and large Practice.

Now

A Treatise on the Mechanism of the Eye,
with several Plates.

Just Published.

Syllabus Cursus Anatomiae, Infirmorum atque Operationum Globi Ocularis, et Partium Contiguarum, D. D. Josepho Cervy, Regiae Majestatis Catholicae Praeclarissimo Medico Primario, Meritissimoque Praefidi, Proto-Medicatui Regnorum Hispaniae, &c. Dicatus; à D. D. Joanne Taylor, M. D. Regisque Magnae Britanniae Medico Oculario, Multisque in Academiis celeberrimis Socio, Authore.

A L S O

Testimonia, Seu Attestationes, Quibus Cathedrali, caeterique Doctores facultatem Medicinae professi, Regiae, insignisque Universitatis Conimbricæ Lusitaniae, De operationibus, methodoque oculorum infirmitatibus medendi Joannis Taylor, Medicinae Doctoris, Britannicaeque Majestatis Physici Ocularii, Attestantur, suasque sententias exponunt, Excelso D. D. Joanni V. Lusitaniae Regi Humillime oblata.

ALSO

L l'Anatomie du Globe de l'Oeil, avec l'usage de ses différentes Parties, & de celles qui lui sont contiguës. Ornê de planches graveës en Taille douce.

Dê dié à Monsieur le Premier Medecin du Roy de France.

A L S O

AN Impartial Inquiry into the Seat of the Immediate Organ of Sight: viz. Whether the *Retina* or *Choroides*. Being the Subject of a Lecture, in a Course lately given on the Nature and Cure of the Diseases of the Eye. To which is annexed, Five and Forty Queries on this Controverted Subject. Submitted to the Examination of the Curious.

All may be had at Mrs. COOPER's, at the *Globe* in *Pater-Noster-Row*; where also may be had the Sentiments of the Foreign Universities on Dr. TAYLOR's Success in the curing the Diseases of the Eye.

DISEASES

OF THE

Chrystalline Humour

OF A

HUMAN EYE:

Or, of the

Cataract *and* Glaucoma.

WITH A

New Theory of their CAUSES, and an Endeavour to demonstrate that there are no membranous Cataracts; but that all Cataracts are from an Alteration of the chrystalline Humour itself.

WITH

An exact Description of a new and more successful Method of making the Operations necessary to the Removal of the several Species of these Diseases.

Humbly address'd to Her MAJESTY.

To which is prefix'd,

A LETTER to the PHYSICIANS and SURGEONS of London and Westminster.

Qui dat videre, dat vivere. Cic.

By JOHN TAYLOR, M.D. Oculist,
FELLOW of the College of Physicians of
Basle in Swisserland, of the Imperial Academy
of Cologn in Germany, and of the Society of
Leige and Rheims.

L O N D O N:


Printed for JAMES ROBERTS, near the Oxford-Arms
in Warwick-Lane. M DCC XXXVI.

1650246



TO THE
QUEEN.

MADAM,

HE great Advantage of
a perfect Enjoyment of
Sight, and the deplora-
ble Condition of the
ind, are so obvious; that I need not
A 2 make

make an Apology for writing a Treatise on this Subject, since nothing of the Kind has yet appear'd in our Language worthy the least Notice. And I am so much the more encourag'd to publish it, that I have the Honour to lay it at your MAJESTY'S Feet, who are as capable of judging of its Solidity and Merit, as You are inclin'd to encourage ARTS and SCIENCES; and whose great Discernment in all Things is equal to those eminent Virtues which have made You the Glory of Your Sex, have added Lustre to Majesty, and inwardly adorn'd the Splendor of an outward Crown.

The Honour, MADAM, which
have had to attend Your MAJESTY
and

was allow'd to lay before You, permits me to hope that this W.O.R.K will have the good Fortune of proving acceptable to Your M A J E S T Y.

It would, M A D A M, be the greatest Presumption in me to undertake to celebrate Your M A J E S T Y's Praises, conscious as I am that I could never rise up to the Grandeur of the Subject; and therefore it better becomes me to admire with a respectful Silence those amiable Qualities that have gain'd Your M A J E S T Y the Esteem and Affection of Your People at Home, and made You the Envy

I am,

M A D A M,

With the greatest

Respect and Submission,

Your M A J E S T Y 's

most humble and most dutiful

Subject and Servant,

JOHN TAYLOR.

TO THE
PHYSICIANS
AND
SURGEONS
OF

London and Westminster.

GENTLEMEN,



F all the Senses, that of Seeing is allow'd to be most useful; yet 'tis a general receiv'd Opinion, that the Organ destin'd to serve to that noble End, is more expos'd to Dangers, more liable to Infirmities and Diseases than any other; and that such Diseases are with more Difficulty.

Use: But I believe the Objection may be soon answer'd, and Nature be found here in no wise to differ from her ordinary Method of acting; if we consider what little Attention has been given to this Branch of Knowledge, and how few have engaged in its Practice.

There is no Doubt but the Cause of this Ignorance arises from the great Nicety that attends its Study, and the little Success that has flow'd from its Practice; if, therefore, from the following Sheets, it may appear I have apply'd myself with more Diligence to the former; or if the Success which I have met with in the latter may be every Day known, I question not, but these will be sufficient Motives to recommend this Piece to your Perusal; and of this I am the more persuaded, because I apprehend, you, who are particularly distinguish'd in the learned World from your Superior Judgment in Physick, can't but be well pleas'd with any Industry, that may be conducive to the improving this most useful, and perhaps only Part, which has been hitherto least known.

Whatever may have been thought of the Method I have taken towards the Advancement of this Science, may be needle

dious Progress I have taken, which I find has given to some a Cause of Objection, has been the only Means and Foundation of my Knowledge and Improvement, as it has furnish'd me with an Opportunity of making my Remarks upon a greater Number of Subjects, in a few Years, than the whole Series of my Life, in a settled Way, could possibly have afforded me.

I wou'd be here understood, that from the many Experiments I have made during the uninterrupted Course of Eight Years (the Time I was on this Progress) I have been furnish'd with those Lights which have enabled me to form a more certain and extensive Theory than my Predecessors could pretend to, or my Contemporaries can boast of. And surely it must be granted, that in all Sciences, that which is supported by a Number of incontestible Facts, is preferable to that which is only founded upon a precarious, and often chimerical Hypothesis: Therefore, Gentlemen, I hope, what I have the Honour to present you, will be read and examin'd, without the least Regard to any private Ends, but purely in the View of rendring Service to the Publick.

Not

for the Cure of which we have hitherto had no true Method propos'd, and which I have endeavour'd to remedy, by a more easy and secure Way than has yet been found out; that of the Cataract, as being more general, is, for that Reason, more worthy of Remark; especially if we consider that it has been thought necessary to expose the Patient to wait sometimes many Years in a very miserable State, for the Maturity of it; in hopes of Success from an Operation which has always been acknowledged dangerous, and so very uncertain, by the ordinary Methods, that 'tis even hard to conceive why it should not rather totally destroy the Sight, than prove a Means of its Recovery.

If it shall appear from the Success some of you have seen, and the Account here given of the Manner of my removing these Disorders, that I am the first who have discover'd the Means to avoid that painful Delay of a Cataract's Maturity, and to remove the several Species of it with less Danger, I hope you will think me worthy your Esteem, as well as that of the Publick.

judged incurable; but if it is evident from my Practice, and in the Progress of this Work, that I have been so happy as to find out a Way of removing several Species of it, I hope, since I explain the Methods, and pretend to no Secret, there will be nothing in this Case that savours of Presumption.

It may not, perhaps, be amiss to observe here, that it has ever been a Custom with the profess'd Oculists of all Nations, to make a Secret of their Practice, and to refuse to communicate to any one a Share of that Knowledge they possess'd. For my Part, I declare publickly my Design of acting in a different Manner, and am so far from desiring to conceal my Practice, that I shall ever use my utmost Endeavours to make it more diffusive and universal.

I am far from affirming, that I have a compleat Knowledge of all the Diseases of the Eye; and own there are many Things I am unacquainted with: But it may be sufficient here to recommend my Practice and Theory to the Judgment of You, who, from your Education, can be no Strangers to the Subject; and I shall beg Leave to finish with this Reflection, That whatever Fate attends this Work,
I know

I am,

GENTLEMEN,

Your Obedient,

Humble Servant,


Suffolk-Street, Dec. 21.
1735.

JOHN TAYLOR.

ON THE
DISEASES

OF THE

Chrystalline HUMOUR.



CHAP. I.

Of the CATARACT in general.



WHAT the *Greeks* call *Hypochy-
ma*, and the *Latins*, *Suffusio*,
is one and the same Disease,
known by the Name of *Cata-
ract*.

Authors are not agreed about the Nature
of a Cataract; some pretend that it is a Sub-
stance of a membranous Appearance, form'd
from a diseas'd Alteration of the aqueous
Humour; others, that it is a Substance of a
membranous Appearance form'd from *Pus*,
B separated

ractions, but that all Cataracts are an Alteration of the chrySTALLINE Humour itself.

They who believe that a Cataract is a Substance of a membranous Appearance from a diseas'd Alteration of the aqueous Humour, tell us, That when any impure Particles are emitted into the aqueous Humour of an improper Quality, Figure, or Magnitude, to pass on in the usual Circulation; they continue moving about this Humour, till by their Levity or Viscidity they assemble together behind the Pupil, and by degrees obtain the Consistence and Opacity of a Pellicle or Membrane.

If we consider the Situation and different Quantity of the aqueous Humour in the two Chambers of the Eye, with the manner of its Supplies, we shall be encourag'd to believe that such a Substance of a membranous Appearance would rather form itself in the anterior than in the posterior Chamber; and since we have no Instance of a Cataract being form'd in the anterior Chamber, we may very reasonably conclude that no Cataract can be form'd from any diseas'd Alteration of the aqueous Humour.

They who believe that a Cataract is a Substance of a membranous Appearance form'd from Pus, separated by a Loss of Continuity

posterior Chamber of the aqueous, which by its Viscidity adheres to the Circumference of the Pupil, and forms a membranous Cataract.

Experience daily proves the Error of this Hypothesis; for if we observe the different Positions of a Cataract in the several Degrees of its Progress, we shall perceive that it begins in the Seat of the ChrySTALLINE, and as its Opacity increases, it appears to rise towards the Center of the Pupil, and that in its last State it admits Light to pass between the Capsula of the ChrySTALLINE and inner Edge of the Pupil, in a sufficient Quantity to discover Shades of Objects.

Whereas if a Cataract adhered to the inner Edge of the Pupil in all or part of its Circumference, it would not only proportionally hinder the Light from passing to the immediate Organ of Sight, but stop the free Motions of the *Iris*; and if we suppose from the Softness of its Texture it would not stop the Motions of the *Iris*, we must allow some Change in its Figure, depending on such Motions; and since Experience makes it evident that the Cataract maintains very nearly, if not exactly, the Figure of the healthful ChrySTALLINE in every State of this Disease, we may very reasonably conclude

They who maintain that a Cataract is an Alteration of the ChrySTALLINE Humour, have proved it by such a vast Variety of unquestionable Experiments, that I think (in the Time I write) 'tis a generally receiv'd Opinion. I shall therefore only endeavour to demonstrate that there are no membranous Cataracts, but that all Cataracts are an Alteration of the chrySTALLINE Humour itself.

The Question is then, Whether there are membranous Cataracts? Or whether all Cataracts are an Alteration of the chrySTALLINE Humour?

Messieurs *Hecquet* (a), *Woolhouse* (b), *St. Yves* (c), *Hovius* (d), *Morand* (e), *Geisler* (f),
Gastaldi

(a) 1. Remarques sur l'utilité de la saignée dans les maladies des yeux.

2. Sa Lettre sur les maladies des yeux pour expliquer ce qui en a été dit dans ses remarques sur l'utilité de la saignée dans les maladies des yeux.

(b) 1. Ses Dissertations sçavantes & Critiques sur la Cataracte & le Glaucome.

2. Ses Observations sur le Memoire Academique de *Monf. Morand*.

3. Son Memoire dans le Journal des Sçavans, *Decembre*, 1720.

(c) 1. Son Traité des maladies des yeux, p. 239.

2. Reponse à une Lettre Critique de son Traité des maladies des yeux.

(d) De circulari humorum motu in oculis, p. 86.

(e) Ses Observations sur la Cataracte dans l'histoire de L'Academie Royale des Sciences, l'année 1722.

(f) Sa Lettre écrite de *Nuremberg* sur la Cataracte.

both these Opinions, and assert, that there are membranous Cataracts, but that they are not so common as those from an Alteration of the ChrySTALLINE.

On the contrary, Messieurs *Heister* (a), *Maitre-Jan* (b), *Brisseau* (c), *Petit* (d), with many others, maintain that there are no membranous Cataracts, but that all Cataracts are from an Alteration of the chrySTALLINE Humour.

B 3

After

g) *Questio Medico-Chirurgica, &c. sub hac verborum*
ie, an Cataracta à vitio humoris aquei aut crySTALLINI ori-
r, &c.

b) Sa Lettre écrite à Mr. *Woolbouse*, &c. *Vid. Journal*
Scavans pour le mois de Juillet, 1722.

i) Ses Observations sur la Cataracte & le Glaucome.

k) Suite des Maladies Chroniques V. 5.

l) *Dissertatio Medica de Cataracta, &c.*

a) 1. De Cataracta, Glaucomate, & amaurosi Tractatio, &c.

2. Apologia & uberior explicatio systematis sui de Ca-
racta, Glaucomate, & amaurosi, contra *Wolbusti* ocularii
Parissensis cavillationes & objectiones, itemque *Parissensis*
ditor, &c.

3. *Vindiciæ sententiæ suæ de Cataracta, Glaucomate,*
maurosi, &c.

) *Traité des Maladies de l'Oeil.*

) *Traité de la Cataracte & du Glaucoma.*

y) 1. Sa Lettre dans laquelle il démontre que le Cristallin
est près de L'uvee, &c.

2. Sa Lettre contenant des réflexions sur ce que M.
Quet a dit dans ses Remarques sur l'utilité de la saignée
des maladies des yeux.

3. Sur les deux espaces que l'humeur aqueuse occupe
à l'oeil & sur le Cristallin & la Cataracte. V. *Histoire de l'*
Académie Royale des Sciences pour l'année 1722, 23, 25,
230.

the Existence of a membranous Cataract; in order to determine this Question, I shall enquire into the exact Situation of the ChrySTALLINE, the true Depth or Thickness of the Chambers of the aqueous Humour, and the Consequences that must attend the Removal of this supposed membranous Cataract from the posterior Chamber.

The celebrated *Dr. Petit* greatly favours the Opinion which maintains that all Cataracts are from an Alteration of the chrySTALLINE Humour, as appears in several of the Memoirs of the Royal Academy of *Paris* and he supports this Opinion chiefly from his having proved by the following Experiment the exact Situation of the ChrySTALLINE and Thickness of the Chambers of the aqueous Humour.

He separated a human Eye from its Fascia and Muscles, and in that Situation it weighed 147 Grains and a half, and its Axis was 11 Lines one Third of Length; this Eye was a little flat on its Sides, under the strait Muscles, which made four irregular obtuse Angles; it had 11 Lines one quarter, measured by two of its opposite flat Sides, from the Right to the Left, and 10 Lines one quarter by two of its opposite flat Sides from the superior to the inferior Parts.

Diameter measur'd from the exterior Part of the *Cornea*. But five Lines and a half when measur'd from the interior Part of the *Cornea*, where the *Cornea* grows wider, by reason of its Inclination.

The Pupil had one Line and a half of Diameter, this was remarked after the Removal of the *Cornea*, which had the Thickness of $\frac{1}{12}$ of a Line. The *Cornea* and aqueous Humour being removed, he took away with a fine Sponge, all that remain'd of the aqueous Humour about the Surface of the Capsula of the ChrySTALLINE.—He weigh'd this Eye, and found less Weight by five Grains, which was the Weight of the aqueous Humour this Eye contain'd.—He then measur'd the Axis of the Eye, and found it nine Lines $\frac{2}{3}$, that is, one Line $\frac{2}{3}$ from the most prominent Part of the *Cornea*, to that of the Capsula of the ChrySTALLINE; from which, if we allow $\frac{1}{12}$ for the Thickness of the *Cornea*, the Thickness of the aqueous Chambers will be exactly one Line $\frac{1}{2}$.

The ChrySTALLINE formed by its anterior Part a Portion of a Sphere, of nine Lines in Diameter; it had four Lines $\frac{1}{2}$ in Thickness, and two Lines $\frac{1}{4}$ in Thickness; the Height of its Segment, had a little more than $\frac{3}{5}$ of a Line.

cessus ciliares. This Capsula confin'd the ChrySTALLINE in the Cavity of the Vitreous Humour; it did not adhere to any Part of the ChrySTALLINE, but was (as it always is) moistened with a little quantity of Liquor.—The ChrySTALLINE weigh'd four Grains $\frac{1}{4}$.

The vitreous Humour which fill'd all the rest of the Eye, weigh'd ten Grains.—The Membranes weigh'd thirty-four Grains. If we put together the Weight of each of these Parts, the Total will be 147 Grains and a half for this Globe of the Eye.

There are several other ways of knowing the true Situation of the ChrySTALLINE, and Thickness of the aqueous Chambers. If we make an Opening into the right Side of the Globe, with the Point of a Lancet, about three Lines below the Edge of the *Cornea*, and another in the opposite Side, exactly the same Distance from the *Cornea*; if afterwards having enter'd a Needle at one of these Openings, and pass'd it in a direct Line thro' the other, we cut round the *Sclerotis* immediately below the *Cornea*, remove the *Uvea*, open the Capsula of the ChrySTALLINE, and take the ChrySTALLINE from its Seat, we shall perceive that the Needle has pass'd the posterior Part of the ChrySTALLINE, without having touch'd the ChrySTALLINE in its Passage.

Edge of the *Cornea*, and another in the opposite Side, exactly at the same Distance from the *Cornea*; if after having enter'd at one of these Openings a small *plano-Convex* Needle, with its plain Surface directed towards the fund of the Globe, and pass'd it, in a direct Line, through the other, we cut round the *Sclerotis*, immediately below the *Cornea*, and remove the *Uvea*, we shall perceive the plain Surface of the Needle immediately placed on the anterior Surface of the Capsula of the ChrySTALLINE,

And we may determine with great Exactness, the Thickness of the aqueous Chambers, by examining with Attention the Origine, and Insertion of the *Ligamentum ciliare*, and remembering that this Ligament never exceeds $\frac{3}{4}$ of a Line in Length.

By these Experiments, we learn the impossibility of removing this supposed membranous Cataract from the posterior Chamber of the Aqueous, without violently breaking or separating the *Ligamentum ciliare* from its Adhesion to the Capsula of the ChrySTALLINE.

And by these Experiments we learn, that if to avoid breaking or separating the *Ligamentum ciliare*, in an Endeavour to remove
this

its Insertion, to the Capsula of the Chry-
stalline, we must make such an Opening in-
to the posterior and anterior Parts of this
Capsula, as to render it incapable to con-
tinue the Chrystalline in its healthful Seat.
— Thus the Chrystalline being no longer
confined in its Capsula, would leave its Si-
tuation in the *Vitreous*, and fall on the *Uvea*.

But even this Care is never taken by those
who favour the Existence of a Membra-
nous Cataract; for by what I have seen, and
by all the Accounts that have come to my
Knowledge, the Operation which has ever
been universally practised, consists only in
passing a Needle a little behind the *Cornea*,
and directing it to the superior Part of the
Cataract, and afterwards in endeavouring
to place the Cataract a little below the infe-
rior and inner Surface of the *Uvea*. Thus from
the above Experiments it appears, That the
Needle must act as much in the Seat of the
Chrystalline, as in the aqueous Chamber,
and consequently must act as much on the
Chrystalline, as in this supposed Membrane.

And suppose the Needle was enter'd into
the Posterior Chamber of the aqueous,
without wounding the *Ligamentum ciliare*,
or Capsula of the Chrystalline, it will be
very difficult to conceive how a Needle of
 $\frac{1}{3}$ of a Line in its Diameter, should move

in

the Removal of this supposed
Cataract; since it appears by the most exact
Enquiry, that the Thickness of the posterior
Chamber of the aqueous never exceeds $\frac{1}{4}$ of a
Line, when the aqueous Humour, (which is
at this Time discharged) is contained in it.

From what I have said it appears, that
those who favour the Existence of a mem-
branous Cataract, must not only necessarily
admit of a different manner of removing it,
from that of a diseased Chrystalline, but
also have some certain way of knowing the
one from the other. — For if before the
Operation they should judge it a Cataract,
from an Alteration of the Chrystalline,
when it was a Membrane in the posterior
Chamber of the aqueous: By endeavouring
to remove this Cataract, they must effectually
destroy the healthful Chrystalline, by
forcing it through its Capsula, or forcing it
with its Capsula, from the *Ligamentum ci-
liare*. And, suppose, during the Progress of
such an Operation, they find their Mistake,
and immediately return their Needle into
the superior Part of the posterior Chamber
of the aqueous, in order to remove this
supposed Membrane, it would be impossible
for them, (as I have already observed) to
pass the Needle into that Seat of the aque-
ous, without violently breaking or separa-
ting

and the aqueous Humour being uncharged, the Chryſtalline, with its Capſula, muſt neceſſarily fall on this ſuppoſed Membrane, and force it ſo irregularly on the *Uvea*, that no Endeavours could remove it, without deſtroying the Uſes of the Chryſtalline, and expoſing the Patient to great Danger.

Thus they would not only be effectually diſappointed of any immediate Succeſs, but in all Probability occaſion an irrecoverable Loſs of Sight.

And if before the Operation, they ſhould judge it a Cataract, from an Alteration in the poſterior Chamber of the Aqueous, when it was a diſeaſed Chryſtalline; by endeavouring to remove this Cataract, they muſt ſo change the healthful Situation of the diſeaſ'd Chryſtalline, or force it through its Capſula on the *Uvea*; that on finding their Miſtake, it would be impoſſible for them to depreſs this alter'd Chryſtalline, without expoſing the Patient to the utmoſt Danger.

And if it appear'd that they had any certain Signs of knowing this Membranous Cataract, from that of a diſeaſed Chryſtalline. The Situation of the Chryſtalline, its Capſula and *Ligamentum ciliare*. — The Smallneſs of the poſterior Chamber of the Aqueous. — The Impoſſibility of moving
a Nee-

the *Uvea*, in the Progress of the Operation from the Discharge of the aqueous Humour. — The wounding or forcibly separating the Parts of the *Ligamentum ciliare*, and the Change that must necessarily happen in the Situation of the ChrySTALLINE, sufficiently prove that Sight could never be restor'd by any Operation possible to be attempted to remove this suppos'd Membrane; but on the contrary, every such Attempt must bring on the most dangerous Consequences.

By this way of Reasoning I was at first inclined to believe, that all Cataracts were from an Alteration of the ChrySTALLINE; and was afterwards strengthened in my Opinion, from the vast Number of Errors in the Sentiments of those who favour the Existence of a membranous Cataract. And my Opinion receiv'd further Support, from having never found any Alteration in the posterior Chamber of the Aqueous, that had the least Analogy to the Account we have of a Membranous Cataract; but have always remarked on examining the Pupil, from the first Appearances of every Species of this Disease, Degrees of Whiteness, and Opacity in the Seat of the ChrySTALLINE; and that in the Progress of their Symptoms, this Whiteness
and

actly, the Figure of a healthful Chry-
stalline.

Besides, if a Cataract could be formed by any Alteration in the posterior Chamber of the Aqueous, I might reasonably have expected to have seen one Instance in such a vast Number of Disorders of this Kind, which have presented to my Care; and I must of Necessity have known it from that of a diseas'd Chrystalline, from its Figure, Situation, and Opacity, consider'd from the Make of the posterior Chamber of the aqueous.

Since therefore not one Instance of a membranous Cataract has presented to my Observation, notwithstanding I have been long favour'd with a very extensive Practice in the Diseases incident to the Eye; and that if such an Instance could be found, it appears impossible to account for its Formation, or to remove it, without effectually destroying the essential Parts of Sight, I am perswaded that there are no Cataracts but what are from an Alteration of the chrystalline Humour.

C H A P.

C H A P. II.

Of the Definition of the Cataract.

BY a Cataract I understand a diseas'd Alteration of the ChrySTALLINE, attended with Degrees of an unequal Opacity and Colour, Loss of Diameter, preternatural Change of its Consistence, Magnitude, Figure, Gravity, and Situation, where the ChrySTALLINE maintains one unequal Continuity, while thro' all its Parts its Capsula still continues to have its healthful Transparency; and in another State of it, it has Degrees of equal Opacity and Colour, Increase of Diameter, preternatural Change of its Consistence, Gravity, and Situation; and here the ChrySTALLINE suffers a Solution of Continuity in some or all of its Parts, with a diseas'd Alteration of its Capsula. The former I call the true, and the latter the false Cataract.

C H A P.

*Of the Causes of the several Species of
the true Cataract.*

AUthors are not agreed about the Causes of the true Cataract; some pretend that it is owing to Degrees of a preternatural Viscidity, in that Fluid brought for the Supplies of the ChrySTALLINE, which by possessing Particles of an improper Magnitude, may be so irregularly lodg'd in its Vessels, as to destroy that Equality so necessary to Transparency. Others, on the contrary, deduce this Opacity of the ChrySTALLINE from an Inflammation of the Blood, and preternatural increas'd Momentum of that Fluid with which the ChrySTALLINE is supply'd; for in that Case gross Particles, inconsistent with Transparency, may be impell'd into the lymphatick Vessels, of which it is compos'd. And others attribute this diseas'd Alteration of the ChrySTALLINE, to a Defect in that Liquor which is found between the ChrySTALLINE and its Capsula, and observe, that when the Vessels intended to convey the Supplies to the
Chry-

the Alterations observable to attend the several Species of this Disease.

Since this Disease happens to those who are the most healthful and temperate, and the most free from viscid or inflam'd Blood, great Variety of Objections must necessarily be remov'd to make any of these Opinions reasonable; which has induc'd me to believe, that these Diseases are owing to a preternatural, forcible, and irregular Contraction of any one or more of the Muscles of the Globe, by which the Parts of the Crystalline are more or less violently and irregularly press'd; and that the Degrees of their Consistence, Magnitude, Figure, Colour, Gravity, Situation, and Opacity, depend on the Degrees of Strength, and Uniformity of this Pressure; and that the Cause of this preternatural, forcible, and irregular Contraction of any one or more of the Muscles of the Globe, is owing to the Deficiency of the Supplies of that Fluid, so necessary to their Sensation and Motion; and that the Cause of the Deficiency of such Supplies, is owing to a long, constant Direction of the Axis of the Eye to particular Objects, by requiring in a limited Time a greater Supply to particular Muscles of the Globe, than

C

what

of the Globe, may occasion such Muscles to act with a degree of Strength different from those which act in Opposition to them; whence follows an Irregularity in the general Action of all these Muscles.

From this preternatural Action of these Muscles, the Globe suffers different Changes in its Figure, by which the several Parts of its Contents are differently forced from their healthful Situation; and the *Ligamentum Ciliare* being fix'd to the Circumference of the Capsula of the Chrystalline, and the aqueous Humour lying immediately on the anterior Parts of this Ligament, and the vitreous on its posterior; the chrystalline Humour being thus confin'd in its Capsula, between the two Surfaces of these Humours, the preternatural Pressure which is thus made on them, must necessarily be continu'd to the two Surfaces of the Capsula and the Chrystalline; and in proportion to the Degrees of Strength, and Uniformity of this Pressure, the Chrystalline must not only suffer a diseas'd Alteration in some or all of its Parts, but the whole Chrystalline must be chang'd from its healthful Situation in its Capsula.

From the Degrees of the Irregularity and Force of this preternatural Pressure, with

degrees of each Species of this Disease.

That the preternatural, forcible, and irregular Contraction of one or more Muscles of the Globe, is the Cause of the several Species of the true Cataract, is further evident from those Alterations of the ChrySTALLINE which sometimes succeed a Blow receiv'd near the Eye; for the Alterations of the Figure of the Globe, from the irregular and violent Action of any one or more of its Muscles in such a Case, are the same in a greater degree, as the irregular and violent Action of any one or more of these Muscles arising from any other; because from the violent, sudden and irregular Change of the Figure of the Globe arising from this Blow, the ChrySTALLINE is press'd with such Violence towards the *Uvea*, as not only to change it from that Situation in its Capsula necessary to receive its Supplies, but also to force it into the posterior Chamber of the aqueous; thus there is such an entire Stop put to the healthful Circulation of the fluid contents of the ChrySTALLINE, and such a change made in the Continuity of its Parts, that all the Parts of the ChrySTALLINE immediately obtain a greater degree of Opacity than could have happen'd in a long time by a less degree of this Pressure, where the Chrystal-

and is caus'd from its continually changing its Situation behind the *Uvea*.

That the preternatural, forcible, and irregular Contraction of any one or more of the Muscles of the Globe is the Cause of the several Species of the true Cataract, is further evident from observing that it always affects such as have employ'd much Time in reading, or viewing particular Objects.



C H A P. IV.

Of the diagnostick and prognostick Signs, and Cure of the several Species of the true Cataract.

THE diagnostick Signs of the several Species of the true Cataract are known from the Situation, Colour, and Degrees of an unequal Opacity of the anterior Surface of the diseas'd ChrySTALLINE.

In the first State of this Disease, we perceive Degrees of a small Cloudiness more im-

And the reason of this small Cloudiness or Opacity appearing near the Center of the Seat of the ChrySTALLINE, is from the central Diameter of the ChrySTALLINE being greater, and consequently containing a greater Number of these alter'd Parts.

In the second State of this Disease, we perceive the Opacity of the ChrySTALLINE to be increas'd, and to be continu'd thro' all the Seat of the ChrySTALLINE, but more immediately in its Center, to maintain the healthful Figure of the ChrySTALLINE, to be advanc'd towards the Center of the Pupil to near half the Thickness of that Part of the posterior Chamber of the aqueous, between the inner Edge of the Pupil and the most prominent Part of the Capsula of the ChrySTALLINE, to have degrees of a Mixture of a pale white and blue Colour, and to continue to impower the Patient to see Shades of Objects in a rect Line from the Axis of the Eye.

In the last State of this Disease we perceive the Opacity of the ChrySTALLINE to be greatly increas'd, and to be very unequal in its anterior Parts, to maintain very nearly the Figure of the healthful ChrySTALLINE; to be so considerably diminish'd in its Volume, to have the Thickness of that Part of the posterior Chamber of the aqueous, between the inner Edge of the Pupil and the most

to have degrees of a Mixture of various Colours, and to permit a sufficient Light to pass between the Edge of the Pupil and anterior Circumference of the Capsula of the ChrySTALLINE, so as to impower the Patient to see Shades, Colours, and sometimes the Figures of many Objects.

The prognostick Signs of this Disease being curable, depend in general on the Perfection of the immediate Organ of Sight; and even in its first State it is seldom remov'd without the Operation.

To remove the first Symptoms of this Disease, I only recommend the Patient to abstain from all such Work as may command great Attention of Sight, such as reading, writing, sewing, &c. to keep the Eye from receiving Light from any very luminous Object, and to cause a Shade to be fix'd before the Eyes so as to prevent the Light from falling immediately within the Axis of the Eye.

The second and last State of this Disease will admit of no Remedy but the Operation.

C H A P. V.

Of the Causes of the several Species of the false Cataract.

THese diseas'd Alterations of the Chry-
stalline are always preceded by an ap-
parent Plenitude, Distention, and preterna-
tural Change of the Contents of the sanguine
and lymphatick Arteries in all the Parts of
the Globe, arising from some diseas'd Alte-
ration of the Blood, by which its Momen-
tum is so increas'd as to force into the Vef-
sels of the Chrystalline (from the Extremi-
ties of those Arteries employ'd in the Service
of their Supplies) certain Particles of an im-
proper Quality to maintain their healthful
continuity.

And from the different Quality and Num-
ber of these diseas'd Particles, some or all
the Parts of the Chrystalline pass thro' such
various diseas'd Alterations of Plenitude,
solution of Continuity, and Opacity, as to
become some, or all of its Parts, chang'd into the
consistence of Milk, Cream, and sometimes
a Matter which has some Analogy in its

of the Capsula, as to render it opaque, and this alter'd Chrystalline being contain'd in its Capsula as in a Bag, has obtain'd various Denominations, such as, *the bag, the milky, and the purulent Cataract.*



C H A P. VI.

Of the diagnostick and prognostick Signs, and Cure of the several Species of the false Cataract.

THE preceding *Optthalmia* being entirely removed, the Patient continue to complain of Degrees of Pain from Light and of a very great Diminution of Sight. on examining the Pupil, we perceive Degrees of a small Opacity continu'd thro' the whole Seat of the Chrystalline, and that this Opacity maintains the Figure of the healthful Chrystalline, and is of a clear blueish Colour.

In the second State of this Disease the Patient complains of such a Defect of Sight, a

the Pupil, we perceive the Opacity of the Chryſtalline to be very greatly increas'd, to continue maintaining the healthful Figure of the Chryſtalline, and to be advanc'd towards the Center of the Pupil to near half the Thickneſs of that Part of the poſterior Chamber of the aqueous, between the inner Edge of the Pupil and the moſt prominent Part of the Capsula of the Chryſtalline, and to have Degrees of a white and blue Colour.

In the laſt State of this Diſeaſe the Patient complains of ſuch an entire Loſs of Sight, as to be only ſenſible of a very little Light to paſs between the inner Edge of the Pupil and Circumference of the Capsula of the Chryſtalline, but not ſufficient to diſcover the Shade or Colour of any Object; on examining the Pupil, we perceive the alter'd Chryſtalline to be ſo conſiderably increas'd in its Volume, as to appear to be immediately placed on the inner Edge of the Pupil; to continue maintaining the Figure of the healthful Chryſtalline, and to be of a dark white and yellowiſh Colour.

The prognostick Signs of Cure in this Diſeaſe are always very doubtful, and any Remedy very precarious, according to the different Temperament of the Patient, and the Probability of being injudiciouſly treated.

To

ly bleeding of the Temporal Artery, Cupping behind the Ears, &c. with the Assistance of Collyriums, and spirituous Fomentations with Camphire, — to keep the Eyes from receiving Light from any very luminous Objects, and to cause a Shade to be so fix'd before the Eyes, as to stop the Light from falling immediately within the Axis of the Eye.



C H A P. VII.

Of the Definition of the Glaucoma.

BY a *Glaucoma* I understand a diseas'd Alteration of the ChrySTALLINE, where the ChrySTALLINE maintains one exact equal Continuity thro' all its Parts, with a diseas'd Alteration of its Capsula, attended with degrees of a very equal Opacity and Colour, very great Increase of Diameter, preternatural Change of its Consistence, Gravity, and Situation; and in its last State with an Elevation, Dilatation, and Immobility of the Pupil, and *Gutta Serena*.

C H A P.

*Of the Causes of the several Species
of the Glaucoma.*

THese diseas'd Alterations of the Chry-
stalline are always preceded by a pre-
ternatural Viscidity of the Blood; and are
owing to the Degrees of the Quantity, Mo-
mentum, and Viscidity of that Fluid brought
for the Supplies of the Vessels of the Chry-
stalline.

And from the Degrees of the Quantity,
Momentum, and Viscidity of this Fluid, all
the Vessels of the Chrystalline suffer various
Degrees of an equal Plenitude, Opacity and
Colour, till the Circulation of their Contents
entirely cease.

And this preternatural Plenitude of the
Contents of these Vessels by degrees so in-
creases the Volume of the whole Chrystal-
line, as to place its anterior Surface imme-
diately behind the inner Circumference of
the Pupil.

And in a further Increase of the Plenitude
of the Contents of these Vessels, the Volume
of the Chrystalline is so greatly augmented,

as

And by this great Increase of the Volume of the Chryſtalline, the Plenitude of the Globe is ſo greatly augmented, as to occaſion Degrees of a preternatural Preſſure on the immediate Organ of Sight.

And this preternatural Preſſure on the *Uvea* and immediate Organ of Sight, is attended with Degrees of a violent Pain immediately in the Fund of the Globe, which continues in degree till in the laſt State of this Diſeaſe there is a Stop put to the Circulation of the Contents of the Veſſels of the Chryſtalline, and conſequently to the Augmentation of it, and Plenitude of the Contents of the whole Globe.



C H A P. IX.

Of the diagnostick and prognostick Signs, and Cure of the ſeveral Species of the Glaucoma.

WHEN the Patient begins to complain of a Diminution of Sight, on inspecting the Pupil, we perceive Degrees

so increas'd, as to appear immediately plac'd behind the inner Circumference of the Pupil.

In the second State of this Disease, the Patient complains of Degrees of the most violent Pain in the fund of the Globe, and of such a Diminution of Sight, as to be unable, from any Direction of the Axis of the Eye, to see more than the Shades and Colours of certain Objects. On examining the Pupil, we perceive the Volume of the Chrystalline to be so greatly augmented, as to have rais'd the Circumference of the Pupil towards the *Cornea*, to near $\frac{1}{4}$ of the healthful Thickness of the anterior Chamber of the aqueous, that the alter'd Chrystalline continues to maintain its healthful Figure, and appears of a dark bluish Colour.

In the last State of this Disease, the Patient complains no longer of Pain, but of such an entire Loss of Sight, as to be even insensible of Light. On examining the Pupil, we perceive the Chrystalline to be so greatly augmented in its Volume, as to have rais'd the Circumference of the Pupil towards the *Cornea*, to near $\frac{1}{2}$ of the healthful Thickness of the anterior Chamber of the Aque-

Disease are always exceeding dangerous, and never admit of any certain Remedy for the Recovery of Sight, but in its first State; and then only from the Operation.

However, the Cure may be attempted in its first State, by a warm attenuating medicinal Diet, Bleeding, Cupping, repeated Emeticks, &c. with the Assistance of Colliriums, and Spirituous Fomentations, — to keep the Eyes from receiving Light from any very luminous Objects, and to cause a Shade to be so fix'd before the Eyes, as to keep the Light from falling immediately within the Axis of the Eye.

The second and last State of this Disease, will admit of no Remedy, for the Recovery of Sight; and the palliative Cure consists in an attenuating Medicinal Diet, warm Bathing, Bleeding in the temporal Arteries, Cupping behind the Ears, &c.



C H A P. X.

Of the Regimen necessary to precede my new Operations for the several Species of the true Cataract, and the several Species of the Cataract called the shaking Cataract. --- For the several Species of the false Cataracts, and the several Species of that State of the Glaucoma, where the Iris, and immediate Organ of Sight, maintain their healthful State.

SINCE a perfect Knowledge of the Animal OEconomy, is in no Case more required, than in the Treatment necessary to these Operations, (which, of all others, are infinitely the most difficult, and attended with the greatest Variety of Accidents)

cuations some Days before the Operation.



Of the Manner of making my New Operations necessary to the Removal of the several Species of the true Cataract, and the several Species of the Cataract call'd the Shaking Cataract. For the several Species of the false Cataract, and for the several Species of that State of the Glaucoma, where the Iris and immediate Organ of Light maintain their healthful State.

C H A P.

C H A P. XI.

*Of the Manner of making my New
Operation for the several Species of
the true Cataract.*

HAVING plac'd my Patient in the usual
Position, I fix my Speculum, and
with a Lancet make a small Longitudinal
Incision, of about half a Line in Length,
through the outward and inward Membranes
of the Globe, into the vitreous Humour. I
make this Incision about one Line and a half
below that Part of the lateral and outward
Edge of the *Cornea*, situated about two
Lines below a Line drawn Parallel to the
greater and lesser *Canthus* of the Eye. I
then pass, in a perpendicular Direction to
the Globe of the Eye, a *plano* - Convex
Needle of about one sixth of a Line in
thickness, through its most Convex Part,
and one third of a Line in its greatest Dia-
meter, with its Convex Surface towards the
superior Part of the Orbit, into the Open-
ing

Vitreous, then very slowly bring my Needle forward with its Surfaces in the same Direction, 'till its Point is brought about a Line below a Line drawn Parallel to the inferior Insertions of the *Ligamentum Ciliare*, with the Capsula of the ChrySTALLINE, and about the same Distance from the Capsula. I then continue my Needle about one Line forward, in the same Direction, and thus pass its Convex Surface, under the inferior Surface of the Capsula of the ChrySTALLINE, about a Line below it, and about the same Distance below the inferior Insertions of the *Ligamentum Ciliare*. I then gently raise my Needle in the same Direction, 'till its Convex Surface suffers some small Resistance from the ChrySTALLINE, which I judge with great Exactness, by inspecting the Pupil, and observing a Motion in the alter'd ChrySTALLINE. And being thus assured that the Convex Part of my Needle is placed immediately under some Part of the Capsula of the diseas'd ChrySTALLINE, I gently raise the ChrySTALLINE about a third or fourth Part of a Line, to learn, with some Exactness on what Part of the Capsula the Convex Part of my Needle is placed; and if on inspecting the Pupil, I perceive that the ar
teric

recedes farther from it, I very slowly continue my Needle towards the inferior Part of the Globe, with its Surfaces in the same Direction, and its anterior Edge at the same Distance from the inner Surface of the *Uvea*, and Insertions of the *Ligamentum Ciliare*, 'till I judge the Convex Part of my Needle, about the same Distance below the inferior Surface of the Capsula, and Insertions of the *Ligamentum Ciliare*, as it was immediately preceding its being first placed under this Capsula. I then replace the Convex Surface of my Needle, on the inferior Surface of the Capsula of the Chrystalline, nearer or farther from the inferior and inner Surface of the *Uvea*, or nearer or farther from the Insertions of the *Ligamentum Ciliare*, according to the Direction I received from the Motion of the alter'd Chrystalline: And when I find by such a Motion, that the Convex Surface of my Needle is so exactly placed under its Capsula, that on raising it towards the superior Part of the Orbit, the anterior Surface of the Capsula of the Chrystalline would be continued, in a direct Line, at the same Distance from all the inner Surface of the *Uvea*: I again raise the Chrystalline about

faces in the same Direction and Distance from the superior and inferior Part of the Globe, and thus permit the inferior Surface of the Capsula of the ChrySTALLINE to return to its former Situation. — The most prominent Part of the inferior Surface of this Capsula, being now about the same Distance below the Point of my Needle, as it was raised from its Situation by the Convex Part of it, I direct the Point of my Needle, with some Violence, about two Lines forward, with its Surfaces at the same Direction and Distance from the superior and inferior Parts of the Globe: And thus the Point of my Needle passes through the Capsula and ChrySTALLINE, about a Line forward from its Entrance. I then forcibly break through those Parts of the ChrySTALLINE, and its Capsula, over which my Needle is placed, and thus make an Opening in the inferior Part of the Capsula; observing by this Action, that my Needle forcibly enters about two Lines into that Part of the Vitreous. — I then withdraw my Needle about three Lines in the same Direction it falls in, and gently raise its Point towards the superior Part of the Globe, about a Line above a Line drawn parallel to the superior and lateral Surface of the Capsula of the ChrySTALLINE,

forward, with its Surfaces in the same Direction, 'till I judge its Point be about a Line above the most prominent Part of the superior Surface of the Capsula. I then gently raise my Hand, 'till the inferior surface of my Needle suffers some small Resistance from the ChrySTALLINE; which I judge of, as before, with great Exactness, by inspecting the Pupil, and observing the Motion of the alter'd ChrySTALLINE. And being thus assured that the plain Surface of my Needle is immediately placed on some part of the superior and lateral Surface of the Capsula of the ChrySTALLINE, I gently depress the ChrySTALLINE about a third or fourth Part of a Line, to learn, with some Exactness, on what Part of the Capsula my Needle is placed: And if on inspecting the Pupil, I perceive that the anterior Surface of the ChrySTALLINE is not continued, in a direct Line, at the same Distance, from all the inner Surface of the *Uvea*, but approaches nearer to, or recedes farther from it; I very slowly raise my Needle towards the superior Part of the Globe, with its Surfaces in the same Direction, and its anterior Edge at the same Distance from the superior and inner Surface of the *Uvea*, and Insertions of the *Ligamentum Ciliare*, 'till I judge the plain Surface of my Needle about the same

face of the Capsula, and Insertions of the *Ligamentum Ciliare*, as it was immediately preceding its being placed on this Capsula. — I then replace the plane Surface of my Needle, on the superior and lateral Surface of the Capsula of the ChrySTALLINE, nearer or farther from the superior and inner Surface of the *Uvea*, or nearer to, or farther from the superior and lateral Insertions of the *Ligamentum Ciliare*, according to the Direction I received from the Motion of the alter'd ChrySTALLINE; and when I find by the Motion of the alter'd ChrySTALLINE, that the plane Surface of my Needle is so exactly placed on its Capsula, that on continuing it towards the inferior Part of the Orbit, the anterior Surface of the Capsula of the ChrySTALLINE would be continued in a direct Line, at the same Distance from all the inner Surface of the *Uvea*; I again depress the ChrySTALLINE about half a Line towards the inferior Part of the Globe; and by this Motion of my Needle, I enlarge the Opening I made in the inferior Part of the Capsula. I then once more very slowly raise my Needle towards the superior Part of the Globe, with its Surface in the same Direction, and its anterior Edge at the same Distance from the superior and inner Surface of the *Uvea*, and Insertions of the *Ligamentum Ciliare*.

the plane Surface of my Needle, about a
Line farther towards the posterior Part of
the Capsula, or about a Line farther from
the superior Insertions of the *Ligamentum*
lenticulare. My Needle thus placed, I conti-
nue it very slowly, directing its plane Sur-
face immediately towards the Opening I
made in the inferior Part of the Capsula of
the ChrySTALLINE; and when I have thus for-
tunately pass'd the greater Circumference of
the diseased ChrySTALLINE, the remaining Part
of it instantly follows. Thus the alter'd
ChrySTALLINE falls into those Parts of the Vi-
trous through which I pass'd my Needle,
after having broke through the inferior
Parts of the ChrySTALLINE, and its Cap-
sula.

During this Change of Situation of the
alter'd ChrySTALLINE, I continue the plane
Surface of my Needle on the superior and
posterior Part of its Capsula, till I perceive
in its Passage on the Capsula about the
Center of the Axis of the Eye, when instant-
ly raise the anterior Edge of my Needle to-
wards the superior Part of the Orbit, and in
this Direction pass it with great Exactness
behind the inferior Surface of the *Uvea*. My
Needle thus plac'd, with great quickness, I
turn its plane Surface towards the inferior

My Needle thus plac'd on the present superior Surface of the Chryſtalline, I inſtantly direct its anterior Edge towards the inferior Part of the inner Surface of the *Uvea*; and by thus changing the Direction of my Needle, its plane Surface is brought about the ſame diſtance towards the preſent anterior Surface of the alter'd Chryſtalline. The plane Surface of my Needle being now obliquely ſituated on the poſterior Part of the Globe, I continue my Needle in the ſame Direction to about a Line below a Line drawn parallel to the Center of the Globe and thus place the Chryſtalline near the inferior and lower Parts of the *Vitreous*. — I continue my Needle about a Minute thus plac'd on the preſent anterior Surface of the diſeas'd Chryſtalline; and after raiſing the Needle's anterior Edge, till its convex Surface is immediately directed towards the ſuperior Part of the Orbit, I very ſlowly take it out of the Eye in the ſame Direction.

By thus paſſing the diſeas'd Chryſtalline thro' the Opening I made in its Capſula, the ſuperior Part of this Capſula being ſuſpended by its Adheſions to the *Ligamentum Ciliare*, the ſuperior and poſterior Parts of the alter'd Chryſtalline paſs with great Facility from under its Capſula, not only becauſe

necessarily comes forward with some degree of Force, to give place for the sudden Reception of the alter'd ChrySTALLINE.

And from this slow and regular Passage of the alter'd ChrySTALLINE thro' the Opening I made in its Capsula, the concave *Vitreous* must necessarily change its Situation, and rise with great Exactness immediately behind the superior and posterior Surface of this Capsula.

The ChrySTALLINE being thus pass'd out of the Axis of the Eye, and the superior and posterior Parts of its Capsula thus forc'd on the superior and inner Surface of the anterior, it is thus the anterior, and Part of the Circumference of the posterior Part of the Capsula, become immediately seated before the Membrane of the now convex *Vitreous*; and thus the Figure of the now convex *Vitreous* is the same with that of the healthful ChrySTALLINE.

If we believe that the *Ligamentum Ciliare* is in any manner necessary to our seeing Objects at different Distances, whether from altering the Figure, or changing the Situation of the ChrySTALLINE; such Uses of it are very nearly preserv'd after this Operation, as in a healthful Eye; for the diseas'd ChrySTALLINE thus pass'd thro' an Opening in its Capsula, and the *Ligamentum Ciliare* maintaining
its

ful Plenitude; it follows that the *Ligamentum Ciliare*, by altering the Convexity of the Capsula, must necessarily place the Surface of the convex *Vitreous* nearer to, or farther from the immediate Organ of Sight, with very nearly, if not exactly, the same Degrees of Strength, as when the Capsula possess'd its healthful Contents.

But if we consider, that in the ordinary Operation for the Removal of the Cataract, the *Ligamentum Ciliare* is always forcibly separated from its Adhesions to the Capsula of the ChrySTALLINE, and by consequence such Uses of it effectually destroy'd; we shall have very good reason to believe that the *Ligamentum Ciliare* has no such Power, since we have many Instances of the Sight having been so restor'd by it, as to have enabled the Patient to see Objects at different Distances.

C H A P. XII.

Of the Reasons of making my new Operation necessary to the Removal of the several Species of the true Cataract, as directed in the preceding Chapter.

I Pass my Needle in a perpendicular Direction to the Globe of the Eye, with its convex Surface towards the superior Part of the Orbit, that by the succeeding Motions of my Needle I may with the greater Certainty place its convex Surface under the Capsula of the Chrystalline. I endeavour to place the convex Surface of my Needle about a Line below the inferior Surface of this Capsula, and about the same distance below the inferior Insertions of the *Ligamentum Ciliare*, that I may afterwards with the greater Facility properly place its convex Surface under the inferior Surface of the Capsula of the Chrystalline.—My Needle thus plac'd, I raise its convex Surface till it meets with a sufficient Resistance from the Chrystalline to
move

be and that my Needle is plac'd immedi-
ately under the Chryſtalline, but learn with
great Exactneſs under what Part of the
Capsula my Needle is thus plac'd. I endea-
vour by the ſucceeding Motions of my
Needle, to place its convex Surface immedi-
ately under a certain Part of the Capsula of
the Chryſtalline, that I may be the more
exact in the Place proper to make the Open-
ing into this Capsula. — The convex Part
of my Needle being properly placed under
the Capsula of the Chryſtalline, I raiſe the
Chryſtalline about a third of a Line towards
the ſuperior Part of the Orbit, and inſtantly
withdraw my Needle about two Lines with
its Surface in the ſame Direction, that by
thus permitting the Chryſtalline to fall a
certain diſtance below the Point of my Nec-
dle, I may with greater Certainty make an
Opening of a proper Length in the inferior
Part of its Capsula. — After having open'd
the Capsula, I continue my Needle about
two Lines towards the inferior Part of the
Orbit, that by thus forcibly breaking into
thoſe Parts of the *Vitreous* over which my
Needle was plac'd, there may be a Place pre-
par'd for the more eaſy Reception of the
diseas'd Chryſtalline. I endeavour by the
ſucceeding Motions ſo to place the plane
Surface of my Needle on the ſuperior and
lateral

surface of the Capsula of the diseas'd Chry-
stalline may maintain the same Distance and
Direction from all the inner Surface of the
Uvea. I endeavour by the succeeding Mo-
tions of my Needle to replace its plane Sur-
face about a Line farther, towards the poste-
rior Part of the Capsula of the ChrySTALLINE,
or about a Line farther from the superior
Insertions of the *Ligamentum Ciliare*, that I
may not be expos'd in passing the Chrystal-
line thro' the Opening I made in its Capsula,
to break any of those Parts of the Capsula on
which my Needle is plac'd, or separate this
Capsula from any of its Adhesions with
the *Ligamentum Ciliare*; which might have
appen'd, had I endeavour'd to force the
ChrySTALLINE thro' the Opening in its Capsu-
le, when the plain Surface of my Needle
was in its preceding Situation: And to the
more effectually answering this Intention,
I continue my Needle in this oblique Dire-
ction of its plane Surface on the superior and
posterior Part of the Capsula of the Chry-
STALLINE, till I perceive it on its Passage on
the Capsula about the Center of the Axis of
the Eye, and then I raise the anterior Edge of
my Needle towards the superior Part of the
Orbit; and in this Direction pass it behind
the inferior Surface of the *Uvea*.

The

inner Surface of the *Crystalline*, I turn this plane Surface of my Needle with great Quickness towards the inferior Part of the Globe, and endeavour by all the succeeding Motions so to place the Chrystalline, as it may maintain its Seat out of the Axis of the Eye. I continue the plane Surface of my Needle for some little time plac'd on the present anterior Surface of the alter'd Chrystalline, to give time for the Parts of the new convex *Vitreous* to possess their new Seat with the greater Exactness, and thus render the alter'd Chrystalline less liable to change from the Situation in which I have plac'd it. I then raise the anterior Edge of my Needle, till its convex Surface is immediately directed towards the superior Part of the Orbit, and draw my Needle out of the Eye in the same Direction, that I may not be expos'd to change any of the present Situation of the Contents of the Globe in its Passage.

C H A P. XIII.

Of the Manner of making my new Operation, necessary to the Removal of the several Species of that Cataract which arises from a Blow receiv'd near the Eye; where the alter'd Chrystalline is pass'd thro' the anterior Part of its Capsula, and being continually changing its Situation behind the Uvea, is known by the Name of the shaking Cataract.

I Proceed exactly as in the Operation for the several Species of the true Cataract, till my Needle is pass'd about two Lines into the *Vitreous*; I then withdraw my Needle in the same Direction, till I judge its Point about one Line from its Entrance into the Globe; I then gently raise the posterior Edge of my Needle, and very slowly direct my Hand backwards, till I judge that its

within the Axis of the Eye. I then continue to direct my Hand backwards, till I judge that the convex Surface of my Needle is obliquely situated from its Point to the inner and outward Surface of the *Uvea*. I then continue my Needle forward with its Surfaces in the same Direction, till I perceive its Point immediately before the present anterior Surface of the diseas'd ChrySTALLINE; I instantly direct the superior Edge of my Needle towards the superior Part of the Globe, till its plain Surface is plac'd on or near the superior and lateral Surface of the alter'd ChrySTALLINE; I gently depress the ChrySTALLINE about a third or fourth Part of a Line, to learn with some Exactness on what Part of the ChrySTALLINE my Needle is plac'd; and if on inspecting the Pupil, I perceive that the anterior Surface of the ChrySTALLINE is not continu'd in a direct Line at the same distance from the inner Surface of the *Uvea*, but approaches nearer to, or recedes farther from it, I then replace the plane Surface of my Needle nearer to, or farther from the Surface of the *Uvea*, according to of the Direction I receiv'd from the Motion the alter'd ChrySTALLINE; and when I find by the Motions of the alter'd ChrySTALLINE,

would be continu'd in a direct Line, at the same distance from all the inner Surface of the *Uvea*. I continue my Needle with its Surfaces in the same Direction thro' the Axis of the Eye, and thus forcibly remove the alter'd ChrySTALLINE into those Parts of the *Vitreous*, over which it was immediately plac'd. My Needle thus plac'd on the present superior surface of the alter'd ChrySTALLINE, I instantly direct its anterior Edge towards the inferior Part of the inner Surface of the *Uvea*, and finish the Operation as in that for the removal of the several Species of the true Cataract.

C H A P. XIV.

Of the Reasons of making my new Operation necessary to the Removal of the several Species of the Cataract known by the Name of the Shaking Cataract, as directed in the preceding Chapter.

MY Needle having pass'd about two Lines into the *Vitreous*, I withdraw in the same Direction, till I judge its Point

E

about

rior Edge of my Needle, and very slowly direct my Hand backward, till I judge that its convex Part is brought parallel to the inner Surface of the *Uvea*, that on directing its Point into the Axis of the Eye, I may not be expos'd to wound any of the Parts of the *Ligamentum Ciliare*. I endeavour by the succeeding Motions of my Needle so to place its plane Surface on the superior and lateral Surface of the alter'd Chrystalline, that on directing my Needle towards the inferior Part of the Orbit, the anterior Surface of the alter'd Chrystalline may continue at the same Distance and Direction from the inner Surface of the *Uvea*. My Needle thus placed, I forcibly remove the alter'd Chrystalline, and finish the Operation as in the true Cataract, for the Reasons given in that Chapter.

*Of the Manner of making my new
Operation necessary to the Removal
of the several Species of the false
Cataract.*

I Proceed exactly as in the Operation for the several Species of the true Cataract, and I find by the Motion of the alter'd Chrystalline that the plane Surface of my Needle is so exactly plac'd on the superior and lateral Surface of the Capsula of the alter'd Chrystalline, that on continuing it towards the inferior Part of the Orbit, the anterior Surface of the Capsula of the alter'd Chrystalline might be placed in a direct Line at the same distance from the inner surface of the *Uvea*. I again depress the alter'd Chrystalline about half a Line towards the inferior Part of the Globe; and by this Motion of my Needle I enlarge the Opening I made in the inferior Part of its Capsule, and lessen the Plenitude of its Contents. I then once more raise my Needle towards the superior Part of the Globe, with its Sur-

E 2

faces

the superior Part of the Capsula of the Chry-
stalline. I then replace the plane Surface of
my Needle about a Line farther, towards
the posterior Part of the Capsula of the
Chrystalline, or about a Line farther from
the superior Insertions of the *Ligamentum*
Ciliare. My Needle thus plac'd, I continue
it very slowly, directing its plane Surface im-
mediately towards the Opening I made in
this Capsula, and thus forcibly remove the
alter'd Chrystalline into those Parts of the
Vitreous thro' which I pass'd my Needle, af-
ter having broke thro' the inferior Parts of
the Chrystalline and its Capsula.

And as the Contents of the Chrystalline are
thus pass'd thro' the Opening in this Capsu-
la, I continue the plane Surface of my Nee-
dle on the superior and lateral Surface of the
Capsula of the Chrystalline, till I judge it in
its Passage on the Capsula about the Center
of the Axis of the Eye, when instantly I raise
the anterior Edge of my Needle towards the
superior Part of the Orbit; and in this Di-
rection pass it with great Exactness behind
the inferior Surface of the *Uvea*. My Nee-
dle thus plac'd, I turn with great Quickness
its plane Surface towards the inferior Part of
the Orbit; thus it necessarily falls on some
of

then direct the plane Surface of my Needle obliquely towards the posterior Part of the Globe, and forcibly continue it about a Line below a Line drawn parallel to the Center of the Globe, and thus mix the Contents of the alter'd Chrystalline with some of the inferior Parts of the Vitreous.

By thus passing the diseas'd Chrystalline thro' the Opening I made in its Capsula, the superior Part of this Capsula being suspended by its Adhesions to the *Ligamentum ciliare*, the alter'd Chrystalline passes with great Facility from under its Capsula, not only from the Assistance it receives by the Needle, but from that of the Vitreous, which at this time necessarily comes forward with some degree of Force, to make room for the Reception of the alter'd Chrystalline.

And from this slow and regular Passage of the alter'd Chrystalline out of the Axis of the Eye, the concave Vitreous must necessarily change its Situation, and rise with great Exactness immediately behind the superior and posterior Surface of this Capsula, and become convex; and as the superior and posterior Part of this Capsula is thus forced on the inner Surface of the anterior, the whole Capsula of the alter'd Chrystalline is immediately seated before the Membrane of the now convex Vitreous; and it is thus the Figure of the convex Vitreous is nearly, if

E 3

not

ful Chryſtalline.

After having thus remov'd from the Axis of the Eye the Contents of the alter'd Chryſtalline, I proceed to remove this opaque Capsula from before the Membrane of the now convex Vitreous; and for this End I withdraw my Needle, till I judge its Point about one Line from its Entrance into the Globe; and after having gently rais'd the posterior Edge of my Needle, I very slowly direct my Hand backwards, till I judge that its convex Surface is brought parallel to the inner Surface of the *Uvea*. I then continue my Needle forward with its Surface in the same Direction, till I judge its Point to be enter'd within the Axis of the Eye. I then continue to direct my Hand backwards, till I judge that the convex Surface of my Needle is obliquely situated from its Point to the inner and outward Surface of the *Uvea*. I then continue my Needle forward with its Surface in the same Direction, till I perceive its Point enter'd into the posterior Chamber of the Aqueous, immediately before the anterior Surface of the opaque Capsula. I then lessen the Pressure of my *Speculum*, and continue my Needle thro' the posterior Chamber of the Aqueous, with its Surfaces in the same Direction, till its Point is carried about half a Line under the inferior and lateral Edge of the Pupil, when the Pupil is neither contracted

fula over which the inferior Edge of my Needle is plac'd, and thus separate with great Exactness all the inferior and lateral Extremities of the *Ligamentum Ciliare* towards the greater *Canthus*, from their Adhesion to this opaque Capsula. I then continue the plane Surface of my Needle towards the Center of the opaque Capsula, and direct its Point forward till arriv'd about the same distance under the superior and lateral Edge of the Pupil, as it was in the preceding Motion of my Needle under the inferior; and by gently pressing with the superior Edge of my Needle, I separate in like manner all the superior and lateral Extremities of the *Ligamentum Ciliare* towards the greater *Canthus*, from their Adhesion to this opaque Capsula. I then withdraw my Needle till its Point is brought from behind the Pupil, and continue to raise the superior Edge of it towards the superior Part of the Orbit, and thus pass its plane Surface immediately before the anterior Part of this opaque Capsula, till its Point is arriv'd about the same distance under the Center of the superior Edge of the Pupil, as it was in the preceding Motion under the superior and lateral; and by gently pressing with the superior Edge of my Needle, I in like manner

opaque Capsula.

The Opaque Capsula of the ChrySTALLINE being thus separated on all Sides from its Adhesions to the *Ligamentum Ciliare*; and the plane Surface of my Needle, now placed on the superior and lateral Part of the anterior Surface of this opaque Capsula, I very gently press with the superior Edge of my Needle on this Capsula; and by continuing its Surfaces in the same Direction towards the inferior and inner Surface of the *Uvea*, the opaque Capsula passes with great Exactness from before the Surface of the now Convex Vitreous. — My Needle with the opaque Capsula of the ChrySTALLINE being now out of the Axis of the Eye, and seated immediately behind the inferior and inner Surface of the *Uvea*, I continue it, with its Surfaces in the same Direction, to about a Line below a Line drawn parallel to the Center of the Globe, and thus place this opaque Capsula near the inferior and lower Part of the Vitreous; I then instantly raise the anterior Edge of my Needle, 'till its Convex Surface is immediately directed towards the superior Part of the Orbit, and very slowly take it out of the Eye, in the same Direction.

C H A P.

C H A P. XVI.

Of the Reasons of making my new Operation necessary to the Removal of the several Species of the false Cataract, as directed in the preceding Chapter.

I Proceed exactly, as in the Operation for the several Species of the true Cataract, 'till the plane Surface of my Needle is properly placed on the superior and lateral Surface of the Capsula of the alter'd Chrystalline; that by the succeeding Motions of my Needle in pushing the Chrystalline towards the inferior Part of the Globe, the anterior Surface of its Capsula may maintain the same Distance and Direction from the inner Surface of the *Uvea*. I then endeavour to replace the plane Surface of my Needle, about a Line farther towards the posterior Part of the Capsula of the alter'd Chrystalline, or about a Line farther from the superior Insertions of the *Ligamentum Ciliare*,
that

Parts of the Capsula, on which my Needle is placed, or separate this Capsula from any of its Adhesions with the *Ligamentum Ciliare*; which might have happen'd, had I endeavour'd to force the alter'd Chrystalline through the Opening in its Capsula, when the plane Surface of my Needle was in its preceding Situation. — The alter'd Chrystalline being pass'd through the Opening I made in its Capsula, I endeavour, by all the succeeding Motions of my Needle, to mix its Contents with those of that Part of the Vitreous, which is seated out of the Axis of the Eye; that notwithstanding its Fluidity, no Part of it shall be able to enter the posterior Chamber of the Aqueous, during the Progress of the Operation, necessary to the Removal of the Capsula from before the Surface of the now Convex Vitreous.

Having thus placed the Contents of the Chrystalline, I withdraw my Needle, with its Surfaces in the same Direction, 'till I judge its Point about one Line from its Entrance with the Globe, for the Service intended by the succeeding Motion. I gently raise the posterior Edge of my Needle, and very slowly direct my Hand backward, 'till I judge that its Convex Surface is
brought

Axis of the Eye, I may not be expos'd to wound any of the Parts of the *Ligamentum Ciliare*. My Needle being enter'd into the posterior Chamber of the Aqueous, I lessen the Pressure of my Speculum, that by the present Loss of the Plenitude of the Globe, from having open'd the aqueous Chambers, I may so considerably enlarge its Diameter, and lessen its Convexity, as to be able to pass with the greatest Facility the plane Surface of my Needle immediately before all the Surface of the opaque Capsula, without being expos'd to alter in its Passage the present Position of any of the Parts of the Capsula or Vitreous. — I intend, by all the succeeding Motions of my Needle, to remove with great Exactness the opaque Capsula, without wounding, or violently breaking any of the Parts of the *Ligamentum Ciliare*, and effectually place it out of the Axis of the Eye.

C H A P.

C H A P. XVII.

Of the Manner of making my new Operation necessary to the Removal of the several Species of that State of the Glaucoma, where the Iris and immediate Organ of Sight maintain their healthful Perfection.

I Proceed exactly, as in the Operation for that Alteration of the Chrystalline, call'd the *Shaking Cataract*, 'till I perceive the Point of my Needle within the posterior Chamber of the Aqueous. I then separate the Capsula of the Chrystalline from its Adhesions to the *Ligamentum Ciliare*, exactly after the same Manner as I have directed in that Part of the Operation for the several Species of the false Cataract, where I remove the opake Capsula from before the Convex Vitreous. — The Capsula of the Chrystalline being now separated on all Sides from its Adhesions to the
Liga-

line about a third or fourth Part of a Line,
to learn with some Exactness, on what Part
of the Capsula of the Chrystalline my Needle
is placed: And if on inspecting the
Pupil, I perceive that the anterior Surface
of the Capsula of the Chrystalline is not
continued in a direct Line, at the same Di-
stance from the inner Surface of the *Uvea*,
but approaches nearer to, or recedes farther
from it; I then replace the plane Surface of
my Needle nearer or farther from the inner
surface of the *Uvea*, according to the
Direction. I received from the Motion of
my Needle the alter'd Chrystalline; and when I find
by the Motion of the alter'd Chrystalline,
that the plane Surface of my Needle is so
exactly placed, that on continuing it to-
wards the inferior Part of the Orbit, the
anterior Surface of the Capsula of the al-
ter'd Chrystalline would be continued in a
direct Line, at the same Distance from the in-
ner Surface of the *Uvea*; I continue to move
my Needle very slowly through the Axis of
the Eye, till its Convex Surface is placed im-
mediately behind the inferior and inner Sur-
face of the *Uvea*, and thus forcibly remove
the Chrystalline with the Capsula, into the
tenuous.

By

give Place for the Reception of the alter'd Chryſtalline; and become Convex.

My Needle thus placed with its plane Surface on the preſent ſuperior Surface of the Capſula of the alter'd Chryſtalline, I inſtantly direct its anterior Edge towards the inferior Part of the inner Surface of the *Uvea*; and by thus changing the Direction of my Needle, its plane Surface is brought about the ſame Diſtance towards the preſent anterior Surface of the Capſula of the alter'd Chryſtalline.

The plane Surface of my Needle being now obliquely ſituated towards the poſterior Part of the Globe, I continue my Needle in the ſame Direction, about a Line below a Line drawn parallel to its Center, and thus place the Chryſtalline near the inferior and lower Parts of the Vitreous. — I then continue my Needle about two Minutes thus plac'd on the preſent anterior Surface of the diſeaſed Chryſtalline: And after raiſing its poſterior Edge till its convex Surface is immediately directed towards the ſuperior Part of the Orbit, I very ſlowly take it out of the Eye in the ſame Direction.

C H A P.

Of the Reasons of making my new Operation necessary to the Removal of the several Species of that State of the Glaucoma, where the Iris and immediate Organ of Sight maintain their healthful Perfection.

I Proceed exactly as in the Operation for that Alteration of the ChrySTALLINE, call'd the *Shaking Cataract*, 'till I perceive the Point of my Needle within the posterior Chamber of the aqueous Humour; that I may with the greater Certainty enter this Chamber, without being expos'd to wound any of the Parts of the *Ligamentum Ciliare*. I intend by the succeeding Motions of my Needle to separate with great Exactness the *Ligamentum Ciliare* from all its Adhesions to the Capsula of the ChrySTALLINE, without wounding or violently breaking any of its Parts. I endeavour by the succeeding Motions of my Needle so to place its
plane

Part of the Orb, the anterior Surface of the Capsula of the alter'd Chrystalline may continue at the same Distance and Direction from all the inner Surface of the *Uvea*.

The Chrystalline with its Capsula being plac'd out of the Axis of the Eye ; with the plane Surface of my Needle on the present superior Surface of the Capsula of the alter'd Chrystalline, I endeavour by all the succeeding Motions so to place the alter'd Chrystalline, as it may maintain its Seat out of the Axis of the Eye.

C H A P.

C H A P. XIX.

f the Regimen necessary to succeed my new Operations for the several Species of the true Cataract, and the several Species of the Cataract call'd the shaking Cataract. For the several Species of the false Cataract, and the several Species of that State of the Glaucoma, where the Iris and immediate Organ of Sight maintain their healthful Perfection.

AFTER the Operation I cause some Drops of a Mixture (made of a Tincture of *Bals. Peruv.* with warm Water) to be dropt into the Eye for some Minutes ; and immediately after I cause to be apply'd over the Lelids a warm Cataplasim, made with some Drops of the above-mention'd Mixture, and the Pulp of *Cassia*. These Drops and Cataplasim are to be repeated every four Hours (except at the time of Sleep) till the second
F Morning

ous Fomentation with Camphire. I then give free Power to the Motion of the Eyelids, and cause a Shade to be so fix'd before the Eye, as to admit of its receiving very little Light, and entirely to prevent the Light from falling immediately within the Axis of the Eye. I cause this Fomentation to be repeated twice a Day for about fifteen Days, and the Shade to be continu'd in different Directions before the Axis of the Eye, till the Eye can perceive Objects in the ordinary Degrees of Light without Pain.

The Patient is to be continu'd for about twenty Days in the medicinal attenuating Diet, accompany'd with gentle Evacuations as mention'd necessary to precede these Operations.

F I N I S.





ADVERTISEMENT.

IN about two Months the Author of this Work will publish a new Treatise on the Anatomy, Supplies, and Uses of the several Parts of the Globe of a human Eye, with its Glands, Coverings, and lachrymal Canals.

To which will be added some Observations on that important Question, *viz.* Whether the *Choroides* or *Retina* is the immediate Organ of Sight?

This Work will be follow'd with all possible Expedition, by new Treatises on the Diseases of the immediate Organ of Sight, *Cornea*, *Uvea*, lachrymal Canals, and Eyelids.

Handwritten text at the top of the page, possibly a title or header.

First main paragraph of handwritten text.

Second main paragraph of handwritten text.

Third main paragraph of handwritten text.

Fourth main paragraph of handwritten text.

Fifth main paragraph of handwritten text.

MECHANISM OF THE E Y E.

Wherein its Power of Refracting the Rays
of LIGHT, and Causing them to Converge at the
RETINA; is Consider'd: With an Endeavour to
ascertain the true Place of a CATARACT, and
to shew the good or ill Consequences of a Judicious
or Injudicious REMOVAL of it.

By John Tayloz, Surgeon in *Norwich*.

*Hear him ye Deaf, and all ye Blind behold!
He from thick Films shall purge the Visual Ray;
And on the sightless Eye-Ball pour the Day!*



N O R W I C H :

Printed by Henry Cross-grove, and Sold by the Book-
sellers of London and *Norwich*, 1727.

12217113

Y. E.

1650248



TO
Mr. WILLIAM CHESELDEN,
Surgeon to St. *Thomas's* Hospital,
AND
FELLOW of the ROYAL SOCIETY.

S I R,



Have heard, that
the Watch-Word
in the Time of
War, is frequent-
ly the Name of One of the
Generals; and that he who
A 2 makes

ing Demanded, *Who Goes there?*) is allow'd, without further Questions, to pass on about his Business.

This engag'd me to ask the Liberty of Prefixing your Name to this Trifle; which however unworthy it may be of the Honour, will yet derive a double Advantage from it; first to Itself, then to it's Author: It will Itself, by this Means, pass unquestion'd into the World; and it's Author will have an Opportunity of paying due Acknowledgments, for the
Benefits

Benefits he has receiv'd from
your Industry and Ingenu-
ity. The Success which has
so frequently attended the
Use of your Needle in this
Delicate Operation, suffici-
ently shews the Niceness of
your Hand; and the little
Damage you have ever done,
where Unavoidable Acci-
dents have conspir'd to ren-
der the Operation unsucces-
ful, is a great Evidence of
the Accuracy of your Judg-
ment.

The Knowledge I have of
this Branch of my Professi-
on, such as it is, I am wholly

A 3

in-

the subject I have had on a
Great Number of Subjects
within these two Years, will
I hope prove that I have fol-
low'd no ill Director.

If I have in any Degree
made it probable, that the
Opacity of the Crystalline
Humor is the True CATA-
RACT, I must here own
that I had the First Hint of
it from You: And if I have
done no Injustice to your
Opinion by unworthily de-
fending it, I hope shortly to
see some One of my Bre-
thren, more able than My-
self,

for the Discovery.

I am,

Worthy SIR,

Your most Obliged

and most Obedient

Humble Servant,

JOHN TAYLOR.



for the Discovery.

I am

Worship Sir,

Your most Obedient

and most Obedient

Humble Servant,

JOHN TAYLOR.





THE
P R E F A C E.



Think it Necessary for my own Vindication to say Something by Way of Preface, and I expect to be pardon'd for it by all Honest Impartial Men; for 'tis to such only I appeal, having already suffer'd as much as I could suffer, from Men of a Character quite contrary.

I am sorry I have it to say, That the worst Usage I have met with, has
been

mine whilst an *Embryo*, has been sufficiently ridicul'd by them: 'Tis, perhaps, still shapeless and unform'd; but if there are found in it the Features of a Man, 'twill at least demand some Regard from all those who would deserve the Character of Men. I have been driven to such a *Dilemma* by their Unnatural Treatment, that for some Time I have suspended the Publication, and was once resolv'd intirely to suppress it. This following has been their Way of Arguing, *If 'tis his Own, 'twill not bear the Light; if 'twill bear the Light, 'tis not his Own*; either Side of this would be equally prejudicial to me: But I shall endeavour to prove, that if I have not made a very ill Use of
my

When I have done this in the best Manner I am capable, I shall desire this following Fair Challenge may be annex'd to my Account of Myself: That if any of my *Unkind Brethren* have any Thing to say, which They think will prove me too ignorant for my Profession, that they may do it in the same publick Manner which I have made Use of to prove that I am not so.

The Worthy Gentleman to whom I have made an Offer of this Trifle in the *Dedication*, will be my Witness of the Truth of the following Particulars: That I attended the Practice of *St. Thomas's* Hospital regularly

nish'd at that Time with a Sett of the
Best and most Rational Practitioners
which have been ever known there.

I think according to the Account
of the Hospital, that there are very
seldom less than 4 or 5000 Subjects
cured of their several Diseases in a
Year; and I believe it will very rea-
dily be granted by all unprejudic'd
People, that where Accidents are so
various and numerous, a young Sur-
geon may improve himself more in
five or six Months, than he can with
the best private Master in the King-
dom in as many Years: Here more-
over a Man has the Opportunity of
learning every Branch of his Profes-
sion; and he who does not endea-
your

to be suspected of great Negligence in his Observation, or Want of Ability to see into the Reason of the Practice ; either of which will certainly make him fearful of undertaking it, because the Success will always appear doubtful to him : But when a Man has by diligent Observation on the Chirurgical Part, and a competent Knowledge of the Rational, qualify'd himself to discharge the Duty of his Profession with Ability and Honesty, I think his endeavouring to do so, should not only not render him obnoxious to the Gentlemen of his Profession, but even make him claim some Share of their Regard, and Protection too from the Insults of the rest of the World :
But

When I came to practise in this City, full of Expectation of the Usage just now mention'd, I soon found myself miserably deceiv'd; for instead of That, I was immediately surrounded with *Calumny* and *Detraction*: And from what Quarter? Why even from those Gentlemen by whom I thought to be *Protected*: There was immediately a *Party* raised against me, who not contented with aspersing my Abilities, fell foul upon my Manners; and would fain have insinuated, (if they could have got any Man of Sense to have believ'd them) that if a Man wore tolerable good Cloaths, and lik'd the Company of a Friend, 'twas impossible

who are themselves

----- *Dully grave, insipidly serene,
And carry all their Wisdom in their Mien.*

And in short, they like Nobody, but such as are like themselves : So that an Awkard *Saint-like* Behaviour, a Solemn Gait, an *Hypocritical* Elevation of the Eye, and an *Affected Religious* Grimace, is sufficient with them to set a Man at the Head of his Profession. If a Man is recommended to any of them, the First Question is, *Is he One of Us?* If they are answer'd in the Negative, they immediately fly ; and tho' they know Nothing of the Man, swear he has no Faith, no Honour, no Capacity,

ly d to be employ d in any Thing.

'Tis a great Hardship to fall into the Hands of these Gentlemen, and yet to be quite void of all those valuable Talents which must recommend one to their Favour. For my Part, if I am call'd to a Patient, I expect my Business is with his *Body*, and not with his *Soul*: I can Set a Broken Leg, as well as my Neighbours; but as to *Cases of Conscience*, I must leave them to be set right by Gentlemen whose Profession qualifies and authorizes them to do it.

But I am afraid, if we come to examin the Matter a little more closely, we shall find that this is generally
all

n odd Way of impoling upon the
World, and giving of it one Term for
another: Their pretended *Gravity*
should be call'd *Natural Dulness*, for
is so in Fact; and where there is
more, it will shew itself: Their Pre-
ference to *Religion*, in Opposition to
Human Reason, is not because they
have *more* of the First than their
Neighbours, but *less* of the Latter:
Their *Canting* Custom of Asking a
peculiar Blessing on every Thing
they undertake, is because they know
their Skill will stand very much in
Need of it: Their Disaffection to
Human Learning is, because they
know Nothing more of it than the
Name; and so of every Thing else:

B

In

ty: copie to truit Mens Lives and
Limbs with.

I have often wonder'd, that when
the ACT was made for Examining
all Surgeons (before they were ad-
mitted to practise in and about *Lon-*
don) at *Surgeons-Hall*, the same Or-
der did not pass thro' every County,
that the Country might have had the
same Advantage of having a Sett of
Able Men amongst them as the Town
has: Many and great would have been
the Benefits of it, to the Profession
in General, to every Able Professor
of it in Particular, and to all those
whose Misfortunes oblige them to
have Recourse to It or Them for Assi-
stance.

being a to be diligently attentive in
his Business, in order to qualifie him-
self to pass his Examination; and
this certainly in Time would be of
infinite Advantage to the Profession:
Every Able Professor of it would
certainly be known, by that very
examination, to be so; and by this
means would be plac'd in his pro-
per Sphere, and fear no Danger from
the *Canting Blockheads* beneath him.
The Benefit of the Patient, is too ob-
vious to be insisted upon; his Limbs
very frequently, and often his Life
so would be preserv'd by it.

But as a *Capacity* is not Every Thing
which is requir'd of a Surgeon, I wish
there was any Method as certain as

others who have had the First and wanted the Latter; but in order to make a Good Surgeon, 'tis absolutely necessary that a Man should want Neither of them. I'll beg the Liberty to relate a Story or Two of a Gentleman who has been remarkably kind to me, and I am very glad I have this Opportunity of returning his Civility, and letting the World know that he is a Man of *Real Merit*: This Friend of mine was sent for to a Child that had dislocated its Knee, he carried with him Plaisters, Bandage, &c. in order to reduce it, which he endeavour'd to do in the best Manner his Capacity would admit of; after the Business was over,
he

he call'd to see his little Patient, and found the Family much pleas'd at his *great* Skill, declaring that the Child's Complaint in that Part was intirely remov'd: The Doctor finding in this Case that he had not spent his Skill in vain, began to exert himself upon it, and undertook a Case of much more Difficulty, (which prov'd a *Dislocation* in Fact) in which he assured them of the like Success, and that he would cure that as expeditiously as he did the other; which, to prove his Assertions true, he actually did; by which *wonderful* Cure he has demonstrated his *Genius* in a *laudable* Manner. Another Brother of mine (some small Distance from this Place) had the Care of a Gentlewoman, whom

than he found her; she applied herself to a Surgeon of my Acquaintance, who made a perfect Cure of her in Fourteen Days: This *Wonderful* Man was ask'd some Time after, How it came about that a Man of his *pretended* Abilities and long Practice, after a Trial of Years, should miss of Success in a Case where a Young Fellow just beginning to practise, succeeded to the Surprize of the whole Town in a few Days? And whether he did not think it ought to diminish his Character, and add to the Character of the other? He had a Conflict with himself for some Time whether he should sacrifice his *Capacity* or *Honesty*; at last it went on the Side of *Dear Capacity*,

him, and hasten to tell you the Reason of my publishing the following Account of a CATARACT.

I am more sensible than any other Man can be, of my own Inability to perform accurately what I have propos'd in my *Title-Page*; my Design not being at all to instruct those whose regular Education in Physick and Mathematicks have qualify'd them to understand these Things infinitely better than I can pretend to do: And I declare before I enter upon the Work, that I have no other View in it than to satisfy the World that I am no *Empirick* (as I have been vilely and maliciously represented to

the Nature and Structure of the EYE,
as is requir'd of any one who pre-
tends to the Operation.

The Reason why this nice and necessary Operation is so rarely performed by Gentlemen regularly bred to the Profession of Surgery, I take to be this; That it having been a long Time the distinguishing Excellency of several of our famous Mountebanks, the regular Surgeons, by undertaking it, are under the dreadful Apprehensions of being rank'd with itinerant Operators, commonly known by the Name of *Quacks*: The Fault is absolutely their own; for did they not join with these *Empiricks* (whom upon
on

entirely despise, to apprehend, that a
Fellow who dares attempt a Thing
which is not the general Profession
of the whole Fraternity, whatever
Pains he has taken, or whatever Ad-
vantages he has had to make himself
Master of the Performance; did they
not join to do this, I say, how ineffe-
ctual would all the little Malice and
Calumny of these contemptible *Wret-*
ches be towards ruining the Reputa-
tion of a regular Surgeon! And how
easie would it be for us, (for 'tis We
only who can do it) by making a
Collection of the injudicious and
barbarous Practices of those illite-
rate murdering *Rascals*, so to destroy
their Reputations, (if 'tis not a Scan-
dal to the Word *Reputation* to use
it

and some other Branches of our Profession Home to ourselves, the Original Proprietors, without any Manner of Danger of the foregoing Imputation.





A N
ACCOUNT
OF THE
Mechanism of the EYE.



S I have propos'd to give
some Account of the Me-
chanism of the EYE, I
know no properer Way
of beginning it, than
by an Anatomical Disse-
ction of the EYE Itself;
and after I have shewn

some few general Diseases that the particu-
lar Coats and Humors of the Eye are sub-
ject to, I shall proceed to shew how the
said Coats and Humors are instrumental to
Vision,

Place of a **Cataract**, and the Consequence
of a *judicious* or *injudicious* Removal of It.

The Globe of the Eye is moved by Six Muscles, Four Streight, and Two Oblique, viz. *Elevator*, *Depressor*, *Adductor*, *Abductor*, *Obliquus Superior*, and *Inferior*: The First of these arises out of the Bottom of the *Orbit*, and after it has ascended and grown tendinous, is inserted into the Upper Part of the *Tunica Sclerotis*, near the *Cornea*. The *Depressor* arises and is inserted directly opposite to the *Elevator*. The *Adductor* arises from the Bottom of the *Orbit*, and is inserted on the Side next the Nose, into the *Tunica Sclerotis*. The *Abductor* arises and is inserted directly opposite to the *Adductor*. The *Superior* Oblique Muscle arises between the *Elevator* and *Adductor*, at the Bottom of the *Orbit*, thence ascending by the *Sutura Transversalis* becomes a Round Tendon, which passes thro' a little Annular Cartilage call'd *Trochlea*, (from its bearing the Form of a Pulley) at the upper and inner Part of the
Orbit,

cle arises from the *Os Maxilla Superioris*, thence passing over the *Depressor*, is inserted at the Bottom of the Eye near the *Abductor*. The Four Streight Muscles are so fix'd to the *Sclerotis*, that they lift up the Eye, depress it, bring it towards the Nose or from it, hold it fast, turn it round, just as they act either separately or conjointly.

The *Superior Oblique* Muscle moves the Eye round towards the Nose about its *Axis*, and pulls it upward and inward, and thereby directs the Pupil outward and downward; and thus the Eye is dispos'd to see Things placed near it. The *Inferior Oblique* Muscle turns the Eye towards the outward Corner, and also upwards; if these Two act together, they fix the Eye fierce and protuberant out of the *Orbit*: Moreover, another Use of these Muscles is, to suspend the Eye, and render it very fit to perform the Motions of the other Four. Between these Muscles Fat is interspers'd, which serves to lubricate and facilitate their Motion.

next

call'd the White of the Eye, and is full of Blood-Vessels; which are very visible in an *Ophthalmia*: The Symptoms of this Disease are Heat, Pain, Pulsation, and an apparent Distention of these Blood-Vessels; it may be occasion'd externally, by any Thing forc'd against or coming into the Eye, irritating or compressing the Fibres of this Coat. If this Inflammation be only external, the Case symptomatical, and the Patient not worne out by other Diseases, 'tis easily Curable: On the Contrary, if it becomes so great as to tumefie the Face, and the Humors very pungent and corrosive, we judge it difficult to cure. Its Cure depends on Bleeding, lenient Purgation, Blistring, Cupping, &c. together with the Assistance of a cooling Diet, topical Application, and proper *Collyria*.

The next Coat is the *Sclerotis*: It expands itself nearly into a Globe, membranous and hard almost like Leather, and encompasses the whole Eye, growing gradually thinner, clearer, and prominent in the Fore-part, which

Humors. This Coat is sometimes diseas'd by a little White Speck of a Cartilaginous Substance, covering Part of, and sometimes the whole *Pupilla*: If this Excrescence is only superficial, it may be remov'd by an Operation in Surgery call'd *Leucotomy*; but if the *Cornea* is throly affected, I believe it incurable. 'This Operation is perform'd in the following Manner, *viz.* I place the Patient in a proper Position, and fixing the *Speculum Oculi* invented for that Purpose in order to keep the Globe without Motion, pare off the Excrescence with a small curv'd Knife, leaving as few Inequalities as possible; and having prevented an Inflammation by proper Repellents, I blow a Powder into the Eye; which, assisted by the Motion of the Eyelid, smooths off the Inequalities left by the Knife.

Next to this Coat lies the *Tunica Choroides*: 'Tis a Membrane of little Firmness, being much thinner than the former; 'tis ting'd by a dark Liquor separated from the Blood-Vessels,

the Circumference of this Hole, and which lies upon the Side of the Cryſtalline Humor, is call'd *Uvea*, compos'd of circular and ſtreight Fibres; 'tis qualified by this Means to contract and dilate, as Light and Objects make different Impreſſions upon it. The Outſide of this laſt Membrane is call'd *Iris*, where appear different Colours: On the Inſide of it from its Circumference ariſe the *Proceſſus Ciliares*, made of ſhort Fibres like Lines drawn from the Circumference to the Center.

The laſt Coat is the *Retina*, ſo call'd from its Reſemblance to a Net; it covers the whole Fund of the Eye, and is nothing more than a fine Expansion of the Medullary Fibres of the Optick Nerve, upon which the Rays of Light meeting in a Point, convey the Images of outward Objects to the Brain. The Diſeaſe of this Coat is call'd *Gutta Serena*; the Eye ſeems clear in Appearance, and there are no other Signs of Blindneſs except a preternatural Dilatation of the *Pupilla*.

Between these Coats are contain'd Three Humors, the *Aqueous*, the *Crystalline*, and the *Vitreous*: 'The First is call'd *Aqueous* because it resembles Water, being a thin transparent Fluid; it serves to extend the *Cornea* into an equal Convexity; besides, it serves moreover for the *Crystalline* Humor to move forward or backward in, as Occasion requires us to see Objects nearer or farther from us.

Next is the *Crystalline*: It's compounded of numberless spherical Segments, closely united and transparent; its Shape is a depressed spheroid, it's inclos'd in a fine Membrane call'd *Aranea*, and is much thicker than either the *Aqueous* or *Vitreous*: Its Diseases and Uses I shall speak of in another Place.

The next is the *Vitreous*: It's very clear and tenacious, it lies behind the *Crystalline*, and fills the greatest Part of the Eye; it's vascular on every Side, its Foreside concave for the *Crystalline* Humor to lodge in, and

C

Backside

This Humor is sometimes so vitiated as to hinder or utterly destroy Sight, which Disease is often mistook for a *Gutta Serena*.

I shall now proceed to speak of the Parts of the Eye, as they are subservient to Vision.

The Use of the Muscles in general is to direct the Globe of the Eye, and fix it in such a Position, as that the Rays of Light reflected from any Object which we would have visible to us, may pass thro' the *Pupilla* and paint the Image of it on the *Retina*.

To form the Image of any Object, 'tis necessary that the Rays of Light coming from it should converge and meet in a *Focus*: And we judge of the comparative Magnitude of all Bodies, by the Angle which is subtended by those Bodies before the Eye.

To qualifie the Eye to perform this, Nature has given it a spherical Form, and compos'd it of transparent Coats and Humors; the latter to transmit the Rays coming from
the

bring them all together in a Point on the
Retina.

All convex diaphanous Bodies refract the
Rays of Light by Attraction, and the Ray
bends towards that Part of the Body where
there is the greatest Quantity of Matter;
this in convex Bodies is the Middle.

Therefore all Rays on whatever Parts
of such Bodies they fall, are refracted to-
wards a Line drawn thro' the Center of
those Bodies, and made to converge behind
them at a Distance proportionable to the
greater or less Convexity of the Bodies them-
selves: Thus when the Eye is too flat, as
in old People, (from a Decay of the Humors,
or a Shrinking of the Coats, or both) the
converging Rays do not coalesce at the *Re-*
tina; but would pass on if suffer'd, and not
come to a Point but at some Distance beyond
: By this Means Vision becomes weak
and indistinct.

gibbous, and the Convexity of them is so great, that the Rays meet and cross one another before they arrive at the *Retina*. The former Eyes are help'd by Convex-Glasses, which put the Rays into a State of easie Refraction before they fall upon the Eye: The latter are help'd by Concave-Glasses, thro' which the Rays passing do diverge, and fall upon the Eye in such a Manner, as to require a greater Degree of Refraction than they would in their natural Course: In the former Case the Eye is not help'd by the Object's appearing bigger, but more distinct; for very small Objects are render'd distinctly visible to such an Eye by the Help of a Convex-Glass, whereas very large ones cannot be seen enough to be distinguish'd by the Naked Eye.

I don't propose to give any new Theory of Vision, nor to be able to describe the Manner of it better than 'tis already done by an eminent Author; for which Reason I shall make use of his own Words, so far as they are serviceable to my present Purpose, and
conductive

a Cataract.

All the Rays which come from one Point of an Object, are by the *Cornea* and Humors of the Eye united in a Point of the *Retina*, which is in a streight Line, drawn from the same Point of the Object, thro' the Center of the Eye; and consequently all the Rays, which come from all the Points of an Object, are united on the *Retina*, in the same Order and Proportion as the Points of the Objects are from whence those Rays come.

Therefore the Interposition which these Rays make upon the *Retina*, must be the Image of the Object.

And thus Vision in general is perform'd : But to know what the several Parts of the Globe contribute hereunto, it's needful to observe that the *Cornea* is more convex than any other Part of it ; by which Means all the Rays are gather'd to pass thro' the *Pupilla*, and none of them are lost upon the *Uvea*.

either the *Ligamentum Ciliare* contracts, or both the Oblique Muscles squeeze the Middle of the Bulb of the Eye, to render it oblong, when Objects are too near us.

The streight Fibres of the *Uvea* dilate the *Pupilla*, when there are but few Rays of Light; and the circular Fibres contract it, when there are too many.

When the *Pupilla* is contracted, we see most distinctly; when it is dilated, we see most clearly.

The *Glassy* Humor keeps the *Crystalline* at such a Distance from the *Retina*, as is necessary for uniting the Rays, which come from one Point of the Object, exactly in one Point of the *Retina*. The Impression of the Object is made upon the *Retina*.

The *Choroides* is tinctur'd black, that the Rays of Light which pass thro' the *Retina* may not be reflected back again to confuse the Image of the Object.

Being

of an Object exactly in one Point of the *Retina*; and that the Rays which come from Objects at different Distances, are united at different Distances, behind the *CrySTALLINE* Humor; they cannot both be exactly united upon the *Retina*, therefore the Eye cannot see equally distinct, at the same Time, Objects at different Distances.

The Optick Nerves pierce the Globe of the Eye, a little on the Inside of the Optick *Axes*. Their external Coat, which is a Production of the *Dura Mater*, is continu'd to the *Sclerotis*, as their internal from the *Pia Mater* is to the *Choroides*; and their medullary Fibres passing thro' all, are expanded into the *Retina*, upon which the Images of Objects are painted.

The Center of this Expansion is insensible, and all Rays which fall upon it are lost; and consequently that Point of the Object from which those Rays come, is invisible to the Eye: The Reason of which proceeds probably from the Blood-Vessels, which enter with

But whatsoever its Cause is, there is a manifest Advantage in the Optick Nerves being inserted on the Inside of the Optick *Axes*: For if they had pierc'd the Eye in the *Axis*, then the middle Point of every Object had been invisible; and where all Things con-
duce to make us see best, there we had not seen at all.

We must likewise have lost some Part of an Object, if the Optick Nerves had been placed on the Outside of the Optick *Axes*: Because an Object may be so placed, as that all the Rays which come from one Point, may fall upon the Outside of both Eyes: But it is impossible they should fall upon the Inside of both Eyes; and therefore that Point which is lost in one Eye, is visible by the other,

This shall serve for the general Theory of VISION: I shall now come to speak of a few Diseases of the Eye, by which Vision is render'd incompleat, or totally

Before I define a **Cataract**, 'twill be necessary to premise a few Things, in Order to clear the Way for my Definition: For since the *Hypothesis* I go upon is New, and not yet clearly demonstrated, it would be look'd upon as a strange Peice of Presumption to overlook in the Gross all the different Objections which may be made against it, and satisfy myself with only a few Probabilities (collected out of my own little Experience and Observation) which may seem to make for it.

Objection I. 'Tis Demonstrated by the Mathematicians, that neither Corpuscles in the *Aqueous* or *Crystalline* Humor can be perceiv'd on the *Retina*; and 'tis said by Physicians, that a Beginning of a **Cataract** is a suffusion of the Eye, when little Clouds, Motes, and Flies, seem to float about in the Air: Now if these Assertions are both true, one would be apt to conclude, that neither the *Aqueous* nor *Crystalline* Humor can be the Seat of a **Cataract**: The first is indisputable,

For I have known a good Number of Subjects with real Confirm'd **Cataracts**, who never remember to have been thus affected at the Beginning of them.

And altho' some at the Beginning of **Cataracts** are thus affected, 'tis no Proof that the Disease may not be in one or the other of these Humors: For the same Cause which will produce a **Cataract**, may at the same Time the **Cataract** is forming affect also the Fund of the Eye; and by disordering the Blood-Vessels, here and there stop the Influx of the Animal Spirits into the little Filaments of the Optick Nerve which help to make the *Retina*. This will certainly produce them Clouds, Motes, &c.

And I have the more Reason to believe this, because I have Couch'd several who after the **Cataract** has been well suppress'd, have remain'd blind with a *Gutta Serena*; which is, I suppose, a total Stoppage of the Influx of the Animal Spirits into the nervous Filaments afore-mention'd.

Objection II.

the Assistance of the *Crystalline* Humor there could be no Vision ; that is, except that Humor lay always ready to receive the Rays and refract them after they had pass'd the *Pupilla*, they would still proceed with so small a Degree of Refraction that their Place of Converging would be beyond the *Retina* ; and by Consequence Vision would be render'd very indistinct, or perhaps the Eye would only perceive a confus'd Glare of Light ; if this Humor was taken away, and this could be demonstrated, 'twould not destroy my Supposition : For I freely acknowledge, that after the Removal of a **Cataract**, (or the *Crystalline* Humor, which as I proceed I shall use synonymously) Vision is never so perfectly perform'd as it was before the Eye was diseas'd.

And if the Subjects I have hitherto perform'd the Operation upon were of Consideration enough to have given me Pains beyond my Needle, I make no Doubt but I could have fix'd them with Glasses which would have still assisted their Sight, and
made

Objection III. If Vision can be perform'd without this Humor, and the Maxim is true *Quod Natura Nihil agit frustra*, Why was it placed in the Eye?

I answer, 'Tis placed there to enable the Eye to see equally well at different Distances: For since the Rays proceeding from the luminous Points of nigh Objects diverge more than those from more remote Objects, the Power of making them equally converge at the *Retina* is very probably lodg'd in the *Crystalline* Humor; which, according to the different Distances of Objects, approaches nigher to, or removes farther from the *Retina*, and perhaps too assumes a different Convexity.

And this I look upon to be unquestionable: Because I have made it my constant Observation, That after a successful Depressure of a *Cataract*, the Patient sees best at a Distance something greater than the Eye requires when all its Parts are in their natural Situation;

Eye, Vision is render'd indistinct or confus'd.

Objection IV. But if the *Crystalline* Humor is depress'd, what becomes of it afterwards? And how is the Cavity fill'd up, which the said Humor possess'd in its natural Situation?

Answer. When I pass my Needle into the Eye, I press directly forward to the *Crystalline* Humor, and endeavouring to disengage it from the *Ligamentum Ciliare*, I depress it to the Bottom of the *Vitreous* Humor; when the *Pupilla* immediately seems clear to me, and such a Torrent of confus'd Light rushes into the Eye, as sometimes affrights the Patient even more than the Puncture of the Needle.

Common Sense tells us, That if a solid Body which is immers'd in a Fluid should be remov'd from one Part of that Fluid into another, it makes no Alteration in the Bulk of the Whole; for whatever Place the Solid is remov'd to, it fills up but so much Space precisely as it did in the Space which it was

queous turns in and fills up the Cavity.

N. B. From the different Specifick Gravity of the *CrySTALLINE* Humor, which in all Probability is different in different Stages of the Distemper, we may account for what is call'd the Rising again of the **Cataract**: As to what becomes of the *CrySTALLINE* Humor after 'tis depress'd, 'tis not unlikely that being broke from the Vessels which nourish'd it, it shrinks by Degrees, and as it were melts into the *Vitreous*.

Dr. Turner says, A **Cataract** is a Disease of the Eye, in which a Film or Skin growing before the *Pupilla*, like a Curtain drawn against a Hole letting the Light into a Room, hinders the Passage of the same, till it is depress'd by the Needle, as in Couching.

There is no one of my Profession who pays a greater Veneration to the Authority of that worthy, learned, and ingenious Gentleman, than myself; I am beholden to him that I have a regular, methodical, and I may venture

always more Danneis in other Parts of Surgery than a Man of his studious Temper could well attend to) very rarely, if ever perform'd the Operation, I hope 'twill not be thought Presumption if when I come to speak of the Operation itself, I should venture to give some Reasons which may seem to thwart his Opinion, and establish my own.

That there are such Films as the Doctor mentions on this Side the *Pupilla*, and in the *Aqueous Humor*, I can readily grant, having met with several of them in the little Course of my Practice: These the Practitioners are allow'd to call *Suffusions*, *Films*, &c. but not **Cataracts**; for I am determin'd to call no Disease of the Eye by this latter Name, except that particular One which is to be remedied by the Use of the Needle: But the Case as mention'd by the Doctor can receive no Benefit that I know of by the Operation, nor did I ever attempt to Couch a Patient (tho' I have had several offer'd me) when the Circumstances were as before-mention'd: This I shall endeavour to demonstrate, in describing the Operation itself. The

a Film separated from the *Crystalline* Humor, and floating in the *Aqueous*; that all its Parts are existent in its first Conformation, that it is a *Cystis*, and that *Cystiffes* are never form'd *de novo*, but are only Membranes disengag'd from the neighbouring Parts, which increasing in Thickness receive Nourishment from the Parts to which they adhere.

I shall beg Leave to examine this Opinion a little, and however specious and refin'd it may seem, I am afraid we shall find as little Reality in it, as there is in the ordinary Profession of Friendship amongst the Gentlemen of their Country.

First they say, 'tis a Membrane disengag'd from the *Crystalline*, and floating in the *Aqueous*; the first they don't offer to prove, and the latter is false in Fact: For if this pretended disengag'd Membrane should float in the *Aqueous*, the Patient would have at least a Chance of seeing now and then a little, by its being remov'd wholly or in part from before the *Pupilla*; but such a Removal we find

n a constant equal Blindness, till the Cause
is remov'd by the Needle: I must beg them
then to fix their floating Island, before they
pretend to make a rational *Hypothesis* of it;
and if they do, I hope they'll be so kind as
to let us know to what Membranes they'll
join it, and in what Manner it receives Nou-
rishment from them, then we'll talk farther
with them.

What have led the *Frenchmen* into the
Talk of its being a *Cystis*, and that no *Cy-
stisses* are form'd *de novo*, is the late Disco-
veries made by the Improvements of Micro-
scopes. They are wonderfully pleas'd with
and fond of the Opinion, that every Thing
which is proceeds originally from an Egg or
Seed; but however true this may be of per-
fect Beings, I don't expect that the same will
be asserted of the *Fungus's* of Trees, and
other supernatural Excrescences: I can be
very well content to believe, that Sloughs and
Mucus may be detach'd from several Mem-
branes in the Body, and may afterwards in-
crease by an Addition of the same Matter.

D

and

prove presently, that 'tis impossible a **Cataract** should be in this Part of the Eye; and that will at once put an End to the Controversie.

What has given me frequent Occasion of reflecting lately, has been the odd Manner of our Authors of General Theories; they undertake to describe and define every particular Disease, with as much Freedom and Positiveness, as if they had been acquainted themselves with a good Number of Patients labouring under such Disease, and had made the Causes, Symptoms, Cure, &c. of it, their peculiar Study.

How just a Proceeding this is, will appear by the following Definition of a **Cataract**, in a late modern Writer of this Sort.

A Suffusion, says he, is a thick Foulness or Excrecence, of the Tunica Cornea, Adnata, or Aqueous Humor, which when confirm'd makes the Cataract.

According to this Account, any thing which thickens the *Cornea*, or makes the *Aqueous Humor*

Cataract; for he seems to use the Word **Cataract** not as any particular Disease, but only as something which shall darken and obscure the Sight, by hindring the Transmission of the Rays of Light. Thus the *Albugo*, *Pterygium*, *Hyopyon*, or the Viscidity of the *Aqueous* Humor, frequently consequent upon an *Ophthalmia*, may be justly deem'd a **Cataract**. But how this squares with the real Notion of this Disease, and its Cure, I shall leave my Readers after a few Pages to determine.

But passing over all the different Definitions and Surmises of others, I shall take the Liberty (for Reasons which I shall give afterwards) to define it in the following Manner.

A **Cataract** is a Disease of the *Crystalline* Humor, whereby the said Humor is totally or at least one Side of it render'd opake: The Humor in this diseas'd State reflects all or most of the Rays of Light, instead of transmitting them to the *Retina*; in Consequence of which, the Humor appears whitish, bluish, yellowish, &c. and the Eye is blind.

Any Blow or Shock, which may alter the particular Position of the Particles of which this Humor is compos'd : For to make a Body diaphanous, 'tis not only necessary that it should be made of such particular Particles; but that those Particles should be combin'd and united after a particular Manner. Thus Salts, wet Paper, or the *Oculus Mundi* Stone; grow opake by being dried; Horn, by being scrap'd; Glafs, by being reduc'd to Powder; or otherwise flaw'd; Turpentine, by being stirr'd about with Water till they mix imperfectly.

The Matter which is separated for the natural Nourishment of this Humor from the Blood, may from the too great Viscidity of the Blood itself, either be separated in too small a Quantity, (the Glands and Strainers of these Parts being exceeding fine) and by this Atrophy the Humors may grow dry and opake, as Paper, Salts, *Oculus Mundi* Stone, &c. Or otherwise upon a Fever or *Ophthalmia*, the Vessels leading to this Humor may be so enlarg'd, as to admit Particles of an improper
Figure

figure or Magnitude: The Heterogeneous
Particles, (if I may so call them) instead
of properly nourishing the Humor, may be
lodg'd irregularly in its Pores, (as Water and
Turpentine as above) and by destroying that
Equality which we are told is necessary to
Transparency, render the Humor opake.

It may be caus'd by violent Grief, which
occasions a more than natural Secretion of
Tears from an adjacent Part; this may from
Continuance make so great a Revulsion,
that the *Crystalline* Humor may probably
grow dry, or suffer Diminution, for Want of
its usual Supplies. I have Reasons to imagine
this, because in Eyes thus diseas'd I have
found this Humor much smaller than ordi-
nary.

It may and often does proceed from an un-
skillful Treating of the Eye in an *Ophthalmia*, whilst there is a more than ordinary Flux
of Matter upon the Parts which seems to in-
crease a Passage thro' the Coats: Instead of al-
luring it forwards by kind Discutients, I have
too frequently known it drove back by cold
repellents; and by lodging itself in the Coats

of its Use and Beauty.

Dr. *Pitcarn*, in speaking of a **Cataract**, says, *Some ascribe it to the thickning or coagulating of the Aqueous Humor, and others to that of the CrySTALLINE: But these know Nothing of the Nature of the Eye; for Corpuscles floating in the Aqueous or CrySTALLINE Humor, can't be perceiv'd on the Retina.*

For once I'll run the Risque of having my Understanding censur'd, and venture to say, That tho' the Corpuscles themselves are not seen being within the Eye, yet they can hinder the Eye from discerning every distinct Part of any Object which is without it: And these little Specks of Darknes, amidst a Glare of Light, are mistaken by the Patient for Something positive, when they are only Privations; and the Effect would be the same, if there were any Stoppages in the small nervous Filaments which compose the *Retina*.

Thus the Doctor's Philosophy is true, when he says they can't be seen, &c. But if he would pretend to inferr from thence, that they can
be

Things, his Consequence is rare, and
demonstrated to be so, by Mr. Cheselden
at the End of his ANATOMY.

DIAGNOSTICK.

'Tis known by a Diminution of Sight in the Patient, and by Something which appears like a Bluish Cloud behind the *Pupilla*, tho' the Colour of it often varies: I do not yet know whether there be any Reason given for the Variety of Colours, but I have observ'd that the Green and Yellow are generally in old Subjects.

PROGNOSTICK.

Not so much the Age of the Person is to be consider'd, as the Age and Ripeness of the **Cataract**: I have seen both very young and very old Subjects couch'd with Success, tho' the best Prognostick is certain to be made in a middle Age and good Habit.

If a Surgeon understands the Operation, and is at all able to judge when the **Cataract** is ripe for it, he has a great deal of Reason to promise Success. If

which he can't foresee, and sometimes 'tis impossible for him to prevent, he is not to be blam'd if he misses of Success, tho' he had seem'd pretty positive in promising it both to his Patient and himself.

Regimen *before the Operation.*

In Order to prevent an *Ophthalmia*, use proper Evacuations, as Bleeding, Lenient Purgatives, &c. a few Days before the Operation.

OPERATION.

The most temperate Seasons are the best, for which Reason I make Choice of *Spring* or *Autumn*. 'Tis requisite the Day should be fine and clear, and the Light well chosen. I oblige the Patient to recline his Head backwards upon a Pillow, the Pillow is supported by the Back of a Chair of a proper Height; behind the Patient an Assistant is plac'd, to keep his Head without Motion; 'twill be necessary too for a Person on each Side, to secure the Hands of the Patient; for I have seen very ill Consequences attend the Neglect

of this: Having thus prepar'd the Patient, I place myself on a Stool, rais'd by Pillows according to the Heighth of the Subject, till I have a full Command of the Patient's EYE; then bidding him turn his Eye towards his Nose, so that the *Tunica Conjunctiva* may be sufficiently apparent, I plunge my Needle directly into the *Vitreous* Humor, thro' the *Tunica Conjunctiva*, about the Distance of the Tenth Part of an Inch from the *Cornea*; then thrusting it boldly to the upper Part of the *Crystalline*, gently turn the flat Part of my Needle on the Top of the **Cataract**; by which Means I have a greater Surface upon it, and the **Cataract** consequently less liable to slip from under my Needle: I depress it very slowly, and place it as I think at the Bottom of the *Vitreous* Humor; I don't immediately remove my Needle off it, but continue it depress'd for about Half a Minute, for Reasons which I shall give when I come to speak of the Causes of a **Cataract**'s rising again: After this, I draw my Needle back about the Sixth Part of an Inch; and gently raising the Point, take it quite out of the EYE.

Regimen

Immediately after the Operation I apply a Repellant Compress, and order the Patient to be put to Bed, raising his Head something higher than usual; and after suffering of him to remain quiet a few Hours, I renew my dressing, and begin to treat him with a cool *Regimen*, proper Diet, &c. as in a symptomatical Fever: This I do to second the Intention of the former *Regimen*; notwithstanding which, an *Ophthalmia* in Spite of all I can do will sometimes supervene, and then it must be treated as such.

These *Ophthalmia's* are sometimes so violent as to become difficult of Cure, and often produce all the Effects, and perhaps too really bring that Distemper upon the Eye call'd *Gutta Serena*. This last mention'd Disease is seated on the *Retina*, and may be brought on by any Thing which will enlarge the Capillary Arteries of that Membrane; for they lying contiguous to the Nerves, and their Diameters being increas'd, the intermediate Nerves are so press'd betwixt them, that the Influx of the Animal Spirits is intirely prohibited,

hibited, and all Communication is cut off betwixt them and the Original of the Optick Nerves in the Brain, which is the *Commune Sensorium*: Thus tho' the **Cataract** continues well depress'd, and the Eye in all Appearance perfectly clear, yet the Patient will remain Blind, and 'tis too often beyond the Reach of Art to give him any Assistance.

However, I generally use all the Evacu-
ations proper in a stubborn *Ophthalmia*, with Calomel, Purgatives, &c. and keep the Patient to an attenuating Diet; obliging him to abstain from all viscid Nutriments, and to use much Exercise: This Method I have known to meet with its desir'd Success.

There are many other Causes besides this Common One, which may render the Operation unsuccessful; first, the Rising again of the **Cataract**,

The **Cataract** will sometimes rise again, even before the Needle is out of the Eye: There are many specious Reasons to be assign'd for it, tho' perhaps none of them true; however I shall mention what I have hitherto imagin'd,

any Thing I shall here say if I find it contradicted by future Practice.

The first Reason (which is perhaps more obvious to others than myself) is, That the *CrySTALLINE* Humor being diseas'd, is become specifically lighter than the *Vitreous*; and that being disengag'd from the Membranes to which it in some Measure adhered, and depress'd to the Bottom of the *Vitreous*, upon Removal of the Needle it ascends by its own specifick Levity, and places itself at or near the Place from whence it was remov'd.

The next is, That consisting of very glutinous Parts, and being join'd perhaps to the *Uvea*, the Needle only removes the main Body, which still remains attach'd to the *Uvea* by certain small elastick Filaments; which as soon as the Needle is remov'd contract themselves, and again lift up the *CrySTALLINE* Humor into its natural Position.

If it remains depress'd for some Days and afterwards rises again, but being a second or third Time depress'd it continues where
you

increas'd Viscidity in the Humor itself, or from a Rarefaction of that Fluid into which it's immers'd; both of which will perhaps be consequent upon an Inflammation, perhaps from an Elasticity in the *Cornea*, whereby the *Aqueous* Humor may be immediately thrust into its Place by Removal of the *CrySTALLINE*; which Elasticity and Contraction of the *Cornea*, may possibly be too great to be overcome by the different specifick Gravity of the *CrySTALLINE*.

But I must confess, if I was to give my Opinion of the general Cause of the Rising again of the **Cataract** immediately upon its being depress'd, I should imagine it to be the lifting up the Point of the Needle before I withdraw it very near back again to the Place from whence I remov'd the **Cataract**. For I imagine that the specifick Gravity of the *CrySTALLINE* Humor, (being very little different if at all from that of the *Vitreous*) would never be sufficient to overcome the Resistance of the said *Vitreous* Humor, which is a very viscid and tenacious Fluid, unless there was a Passage made for it to rise again by that Motion

press'd; and therefore hold it necessary that after the **Cataract** is depress'd, the Needle ought to be kept some Time upon it, that the Fluid above it may have Time perfectly to reunite; so as becoming all over equally tenacious, it may be impower'd to keep the **Cataract** in the Position we would have it.

Another Cause of ill Success is, the wounding the *Uvea* or *Iris*; the Consequence of which will be an irregular Form of the *Pupilla*, some of which I have seen intirely clos'd: However if they are not so, there is always such a Callosity upon the wounded Membrane, as stops up the *Pupilla* in one Part, and leaves such an Opening of it on the other, that by stopping up the Rays where they should pass, and suffering them to pass where they should not, the Eye is render'd intirely useles.

These Parts too, when wounded, sometimes ulcerate; by which Ulceration, the whole Coats and Humors of the Eye become one common Mass of *Pus*, and the Eye intirely sinks.

The

wound the *Uvea* and *Iris* is, when he would endeavour to remove a Film which lies in the *Aqueous Humor*, or on this Side the *Pupilla*. And this, till I know better, I shall call an *Injudicious Practice*.

It often happens that the said Coats are wounded thro' the Fault of the Patient, even when the Operation is the most judiciously perform'd; for he not having Courage to keep his Eye in the Position he was directed, and by endeavouring to fly from the Pain which is given by the first Puncture of the Needle, which he imagines is only to be done by Motion, and not being able to move his Eye any farther towards the great *Canthus*, turns it back against the Point or Edge of the Needle; and by this Means, in Spite of all the Art or Address of the Surgeon, will with that Motion wound the Coats above-mention'd: But this last Accident does not happen to a Man who knows what he is doing, perhaps above Once in Twenty Times.

Another Reason of ill Success is, the **Cataract** proving Milky: To be able to determine

with my Needle I have not yet learn'd, and should be glad to be inform'd by any Body who has.

Another Cause of ill Success, which is also sometimes attributed to the Surgeon, may be a strange Fondness of Sight, which engages the Patient after having been couch'd Three or Four Days to pull off the Dressings in the Absence of the Surgeon; by which he exposing the Eye to the Assaults of the open Air, brings on that *Ophthalmia* which we were endeavouring to prevent: In this Case he himself is blamable, and not the Surgeon, for all the ill Consequences that may attend it, one of which I have found often to be the intire Loss of Sight.

Another Misfortune, which too frequently attends the unhappy Patient, is, when the **Cataract** is accompanied with a *Gutta Serena*, perhaps of as long standing as the **Cataract** itself; this we find when upon depressing the **Cataract**, (and there's no Possibility of judging of it before) and the *Pupilla* remaining clear, the Patient notwithstanding continues
in

plorable. I should be unwilling to think that there was any Part of Mankind such Strangers to common Sense, as to imagine this to be a Fault of the Surgeon; but, I myself have more than once been calumniated on this very Account: I can't help saying, I have been unfairly dealt with by some of the Gentlemen of my own Profession, who have misrepresented the Cases, and deluded the common People, as well as injured me; this was the Case of Three Subjects that I lately Couch'd in *Yarmouth*.

N. B. I have met with the same Treatment on Account of a Subject which I Cut for the Stone, where I will undertake to demonstrate to any Man in *England*, that it was as impossible to meet with Success in that Operation as it was circumstanc'd, as it is to succeed in the Operation of Couching where the **Cataract** is attended with a *Gutta Serena*; for unless a Man could directly look into the Body in the one Case, or down to the Bottom of the Eye in the other, he can be no more able to make a positive Prognostick, than he would be to alter the Circumstances.

I shall make here another Observation,
That where there is a **Cataract** in one Eye,
and the other Eye sound and good, the Pa-
tient don't ought to be encourag'd to suffer
the diseas'd Eye to be Couch'd; because the
Symptoms which may possibly attend it, may
be the Occasion of the Loss of both.

I shall take the Liberty of mentioning a
Case of my own, wherein I have met with
some of the afore-mention'd Accidents. A-
bout Five Months since, a Patient came to
me with a Confirm'd **Cataract** in one Eye,
and a Suffusion in the other. I found the **Ca-
taract** good, and (after the common Prepa-
rations) depress'd it with no apparent ill Ac-
cident. The Eye continu'd clear, and the
Patient saw for some Days: At which Time
an Inflammation came on, which equally af-
fected both Eyes; and, notwithstanding all
I could do to abate it, the Coats and Hu-
mors were so thicken'd by it, that the Eye
which had the Suffusion in it became totally
blind: And some Time after, that which I
had couch'd was in like Manner intirely de-
priv'd of Sight; tho' the **Cataract** remain'd
depress'd, and the *Pupilla* perfectly clear.

Observation.

the Suffusion above-mention'd was Nothing more than the Beginning of a **Cataract**, caused by the *Crystalline* Humor growing thick and muddy: And that this Inflammation, by increasing the Viscidity of the Matter separated for the Nourishment of the said Humor, increas'd the Thickness so, as to render the Humor intirely opake; or in other Terms, ripen'd the **Cataract**. The Reason why I did not venture to Couch it afterwards was, an uncommon Dilatation of the *Pupilla*, which had been from the Beginning of the Suffusion: The Disease of the other was a direct *Gutta Serena*, caus'd by the Inflammation in the Manner before describ'd.

The Subject I have been treating of being altogether new, I shall beg Leave to add the following Queries, to be consider'd by the Gentlemen of my Profession, not troubling myself whether they make for or against my *Hypothesis*.

Query I. Whether the Reason of the Dilatation of the *Pupilla* in one Eye and not the other may not be this: That the *Crystalline*

E 2

Humor

rent than formerly; the Patient finds that the Rays of Light do not strike so clearly and forcibly as they us'd to do; or in other Words, the Eye is darker than in its natural State: To amend which, he naturally and one may almost say involuntarily dilates the *Pupilla*; in order to receive more Rays of Light, and recompense the Defect of the diseas'd Humor?

The same is done whenever we want to see Objects in the Dark; but in this case the Cause remaining constant, the Dilatation is constant too, till the *Pupilla* by the frequent Use of this Motion becomes actually wider than it ought naturally to be. In such a Dilatation as I here mention, we can't promise any great Success from the Use of the Needle: For was the Humor itself perfectly clear, such a dilated *Pupilla* would admit such a Glare and Flow of Rays to the *Retina*, as would render Vision confus'd and indistinct; much less can we hope to have a good Eye, when that Humor so necessary to Vision is intirely removed.

Query II.

times appearing at first in a white Point, and afterwards spreading itself like a Line drawn from a Center to a Circumference, is not a Proof that 'tis a transparent Body already form'd, growing thick and cloudy, rather than the Beginning of a new Membrane?

N. B. The Composition of the *Crystalline* Humor, in the Anatomy of the Eye.

Query III. Whether **Cataracts** very old and discolour'd, declining in the same Manner, and becoming angulous and uneven, is not a Proof of the same Thing?

Query IV. Whether it would not be worth while to observe after the Couching a young Subject, if the same Subject is ever troubled with a second **Cataract**, properly distinguishing whether it be the same **Cataract** risen again or not? And this I think is best to be done, by observing the Interval of Time which passes between his being Couch'd and the Appearance of this second **Cataract**. Whether if this Observation be just, and a second **Cataract** be form'd in the same Eye, and

Query V. Whether our not being able to remove a **Cataract** at its first Appearance, is not a Proof that 'tis not the *Crystalline* Humor? Because for all what we know certainly, the *Crystalline* Humor is equally removable at all Times.

Query VI. Whether or no in that Stage of the Disease which we call a *Milky Cataract*, there may not be such a Solution of Continuity of the Parts of the *Crystalline* Humor, as may suffer the Needle to pass backwards and forwards in it, without being able to remove it out of its Place?

Query VII. And whether in a farther Stage of this Disease, the Parts may not again close, and stick together in such a Manner, as that the whole Body of the Humor may be at once remov'd, by only pressing upon a particular Part of it?

Query VIII. Whether there may not be a Gradation of Colouring settled from Sir Isaac
Newton's

and ending at a greenish Yellow; whereby we might be able to determine the exact Age of a **Cataract**, as soon as we see it? And whether such a particular Colour will or will not successfully bear the Needle?

Query IX. Whether 'tis not possible to invent a *Speculum Oculi*, which may keep the Globe of the Eye without Motion, and without giving that Pain to the Patient which too frequently contributes to the bringing on of an Inflammation? I mention this, because I have seen so many Inconveniencies attend the Use of the Common *Speculum Oculi*; that, rather than use it myself, I have chose to have Nothing to assist me, except the Pressure of the Thumb and Finger?

Query X. Whether a Trembling, which I have observ'd in the *Aqueous* Humor some Time after Couching, does not proceed from a Loss of Part of that Humor in Couching? And whether this Trembling is one Reason why the Rays of Light pass confus'd into the Eye?

Needle to pass between the *CrySTALLINE* Humor and the *Uvea*, without removing the former so as to spoil the Eye, or cutting or lacerating the latter?

Query XII. Whether, if this last is practicable, there's not a Possibility of removing a Film that may be in the Forepart of the *Pupilla*, notwithstanding the Needle enters in at the Backpart of it?

Query XIII. Whether there might not be a nice Experiment made, by placing an Eye like a *Lens* in the Hole of a dark Chamber? I think (if I remember rightly) I have met with the following Account in an Author of Experimental Philosophy. Stripping the Fund of the Eye of its Integuments, he plac'd it as above-mention'd; and holding a Peice of white Paper within the Room, at a small Distance from the Eye, he distinctly perceived the Images of all Objects from without painted upon the Paper; as they would have been upon the *Retina*, if the Eye had been in its natural Position.

Had

have gon on the Outside and with a Needle as in Couching deprefs'd the *Crystalline* Humor, and if notwithstanding this Depression the Person within could still perceive the Images as before, it would be a certain Proof that the Eye might see, notwithstanding the Loss of the *Crystalline* Humor.

The Eye should be taken down, and dissected; and by observing in what Part the *Crystalline* Humor after its Removal was lodg'd, we should receive a great Light into the Theory of Couching, and much Satisfaction as to the Truth of this *Hypothesis*.

I only mention this as it seems to me possible, without being able to determine whether 'tis practicable or not.

CONCLUSION.

Before I conclude, I thought it my Duty to mention, (and in the Words of a Great Man) That as a candid and sincere Behaviour ought to be preferr'd to all other Considerations, I would not let my Heart reproach

gain'd even the Possibility of a Cure, which is not in every Particular my Right : I therefore thought proper to acknowledge, that I am oblig'd to a very ingenious Gentleman (whom I have the Honour to be acquainted with) for Two or Three Hints that he has given me in the latter Part of this short Dissertation, which otherwise probably might have been omitted.

N. B. I believe it will be necessary to observe to such Persons as are not Judges of the Subject I have been treating of, That I have therein demonstrated the Reason of my not meeting with Success in some Patients that I have Couch'd, particularly in *Yarmouth*.

F I N I S.



E S S A Y

UPON

SINGLE VISION WITH TWO EYES:

TOGETHER WITH

E X P E R I M E N T S

AND

OBSERVATIONS

ON

SEVERAL OTHER SUBJECTS IN OPTICS.

By WILLIAM CHARLES WELLS, M. D.

L O N D O N:

PRINTED FOR T. CADELL, IN THE STRAND.

1792.

1650251

C O N T E N T S.

E S S A Y

UPON

SINGLE VISION WITH TWO EYES.

P A R T I.

Of the different Opinions concerning Single Vision with two Eyes; and principally of those of Dr. Smith and Dr. Reid, - p. 1.

P A R T II.

Of a new Theory respecting Visible Direction, and of a Solution hence derived of the Question, why Objects are seen single with two Eyes, - - - P. 34.

P A R T III.

Of some Consequences from the foregoing Theory of Objects being seen single with two Eyes; together with the Explanation of several other Phenomena of Vision, p. 63.
EX-

EXPERIMENTS *and* OBSERVATIONS

ON

SEVERAL SUBJECTS IN OPTICS.

ARTICLE I.

On Visible Position and Visible Motion; p. 85.

ARTICLE II.

On a supposed Consequence of the Duration of Impressions upon the Retina; and the Effects of accurate Vision being confined to a single Point of that Membrane, p. 106.

ARTICLE III.

On the Connection between the different refractive States of the Eyes, and the different Inclinations of the Optic Axes to each other, - - - p. 116.

ARTICLE IV.

*On the Limits of Perfect or Distinct Vision,
p. 131.*

E S S A Y

UPON

SINGLE VISION WITH TWO EYES.

P A R T I.

*Of the different Opinions concerning single Vision with
two Eyes; and principally of those of
Dr. Smith and Dr. Reid.*

THE end I have chiefly in view, in this
essay, being to offer a new solution of the
question, why objects are perceived single
with two eyes, I think it incumbent upon
me, in the first place, to show, that none of
the opinions I have met with upon this
subject, can be admitted as just.

These opinions, or such of them at least
as have gained any considerable reputation,

B

may

may be reduced into two classes. The first comprehends those of Galen, Alhazen, Rohault, Dr. Briggs, and Sir Isaac Newton, all of whom have regarded the question I have mentioned as equivalent to the following one : Whence comes it, that the mind should be affected with only one perception from two impressions upon the external organs of sight, since either of those impressions is, of itself, sufficient to produce a similar perception ? Their universal answer has been : Because the two impressions are united before they are communicated to the mind. And the only difference among these authors, has been with respect to the manner in which such an union takes place. To the second class are to be referred the opinions of those, who hold it as certain, that an object is seen single by both eyes, because it is seen by each of them in the same external place ; and who profess to point out some law, or constant rule of vision, from which this sameness of place is to be derived as a necessary consequence. Aguilonius, I believe, first gave
this

(3)
this view of the question, which has since
been adopted by Dechales, Dr. Porterfield,
Dr. Smith of Cambridge, and Dr. Reid of
Glasgow.

In ~~reference~~ ^{opposition to} the opinions of the
first class, more especially as they have
been repeatedly examined by others, I
think I need only say, that they must all be
considered as mere conjectures, founded
upon certain supposed changes in the brain
and nerves, the existence of which it is im-
possible, from the nature of the parts,
either to demonstrate, or to refute by expe-
riments; and that no one of them, though
admitted to be true, is yet sufficient to ex-
plain the phenomena on account of which
it was framed.

The opinions of the second class being
built, as their authors think, upon experi-
ments and observations, both allow and
demand a more accurate investigation. I
shall proceed, therefore, to examine such of
them as I am acquainted with, beginning
with that of Aguilonius; and what I
shall observe concerning it will apply also

to those of Dechales and Dr. Porterfield, who have done little more than copy what he has said.

If a line be drawn through the point of the mutual intersection of the optic axes, parallel to the interval between the eyes, Aguilonius calls it, from its office, the *horopter*; and if through this line, a plane be made to pass at right angles to that of the optic axes, he names it the *plane of the horopter*. After defining these terms he asserts, that, by a law of our constitution, all bodies which we see with one glance or look, whatever are their real places, appear to each eye to be situated in this plane.— And if this be granted to him, he easily and satisfactorily shows, why some should be seen single with two eyes, and others double. For since, according to a second opinion maintained by him, and not contradicted, I believe, by any other writer upon vision, the two lines of direction, in which an object is seen when we employ both eyes, can meet each other only in one point, it follows, that all bodies which are

4

really

really situated in the plane of the horopter, must necessarily appear single, as the lines of direction in which any one of them is perceived by the two eyes, coincide in that plane, and no where else; and that all bodies; which are not situated in the plane of the horopter, must as necessarily appear double, since, in this case, the lines of their visible directions intersect each other, either before or after they pass through it.*

Against the truth of this explanation, only one argument need be offered.—Were the visible places of all bodies to be contained in the plane of the horopter, these would appear of magnitudes proportional to the angles which they subtend at the eye. A finger, for instance, held near to the face, would seem as large as the part of a remote building it might conceal from the sight. But as this is contrary to experience, the principle from which it is derived, must be rejected, together with all its consequences. To Aguilonius, however, the merit is due of being the first, who

so

* Aguilonii Optica, p. 110, 148, 331, 344.

so far generalised the phenomena of single and double vision, as to observe, that those objects alone are seen single, which are really situated in the plane of the horopter.

The opinion of Dr. Smith is the next in the order of time. * " If it
" be asked (says that author) why in
" seeing with both eyes we do not al-
" ways see double, because of a double sen-
" sation, I think it is sufficient to say,
" that in the ordinary use of our eyes,
" in which the pictures of an object are
" constantly painted upon † correspond-
" ing places of the retinas, the predomi-
" nant

* Compleat System of Optics. Vol. I. p. 48.

† Dr. Smith gives the following definition of *corresponding points*. " When the optic axes are parallel or meet in a point,
" the two middle points of the retinas, or any points which are
" equally distant from them, and lie on the same sides of them,
" either towards the right hand or left hand, or upwards or
" downwards, or in any oblique direction, are called *correspond-*
" *ing points*." Vol. I. p. 46. According to this definition,
points correspond which have a certain agreement in situation.
No contradiction is, therefore, implied in this system by saying,
that an object may appear single, though its pictures should fall
upon points which do not correspond. Dr. Reid's definition of
the same term is very different.

“ nant sense of feeling has originally and
 “ constantly informed us that the object is
 “ single. By this means our idea of its
 “ outward place is connected with both
 “ those sensations, as is manifest by its
 “ appearing in two places when its pictures
 “ are not painted upon corresponding places
 “ of the retinas; which is only a direct
 “ consequence arising from our general ha-
 “ bit of seeing.” Should any one now en-
 quire whence it is, that, to produce single
 vision, all men agree in directing their eyes
 toward the object in such a manner, as to
 receive its pictures upon corresponding
 points of the retinas, since custom might
 have connected the sensations of any other
 two points with the information of its unity
 from feeling :* This answer may be given
 in the words of Dr. Smith : † “ When we
 “ view an object steadily, we have acquired
 “ a habit of directing the optic axes to
 “ the

* This objection is made to Dr. Smith's theory by Dr. Reid,
 —who seems to have overlooked the answer. Reid's Inquiry into
 the Human Mind, 3vo. p. 232.

† Vol. I. p. 46.

(8)
the point in view ; because its pictures
“ falling upon the middle points of the
“ retinas, are then distincter than if they
“ fell upon any other places ; and since
“ the pictures of the whole object are equal
“ to one another, and are both inverted
“ with respect to the optic axes, it follows
“ that the pictures of any collateral point
“ are painted upon corresponding points of
“ the retinas.”

Such is the solution which Dr. Smith has given of this celebrated question, and such the reply, which his general account of vision furnishes to one objection against it. But there are others which, in my opinion, cannot be so easily repelled. Before I offer these however, I beg leave to remark, that although it were proved, as I think it may be, that he is mistaken in the fact of objects appearing single, when their pictures fall upon the middle or other corresponding points of the retinas, still the truth of what is peculiar to him * of the solution
he

* Dr. Reid attributes to Bishop Berkeley the opinion, that objects appear single to two eyes, from an experienced connection
between,

he gives, might remain unshaken.— Objects, it may be said, are constantly seen single when we direct our eyes to them in a particular manner. Their pictures must, consequently, in every such case, fall upon the same places of the retinas; and whether these be corresponding or not, the unity of the visible appearances will be owing to the connection, which has uniformly been observed between the sensations of those places, and the information from feeling, that the objects which cause them are single. What I shall say, therefore, upon his opinion, will tend to show, that, admitting the fact respecting corresponding points to be true, his explanation of it ought however to be rejected.

For *first*, it may be observed, that, if we are taught by *feeling* to see objects single,

between particular sensations of sight, and the informations of touch. But I nowhere find it mentioned in the works of that author; and I even think it probable, that he purposely avoided treating of the question, as he found, that the solution of it, which naturally flowed from his principles of vision, was with difficulty to be reconciled to other conclusions he had derived from the same source.

notwithstanding a sensation in each eye, the informations of the former sense ought to be uniform, or else one set of visual appearances would be associated with different reports from feeling, and no certain mark afforded us which of them we should trust. Now Dr. Smith himself is obliged to confess, that we sometimes *feel* double; “ as
“ in the dark, when a button is pressed
“ with two opposite sides of two contiguous
“ fingers laid across; for this reason, that
“ those opposite sides of the fingers have
“ never been *used* to feel one but always
“ two things at a time.” * He adds, “ We
“ have learned, therefore, by experience of
“ both senses compared together, to make
“ their informations consistent with each
“ other.” Here then we find him to allow, that feeling is not always the predominant, but sometimes the inferior sense; that its informations are not constant and original, but changeful and derived; positions directly

* Vol. I. p. 48. Dr. Smith however has, from the influence of system, I suppose, mistaken this fact; for the button is *felt* double, when pressed in the manner above mentioned, though we should not be in the dark, and should even *see* it to be single.

rectly contrary to those he had immediately before maintained. But in the first instance of difference between the informations of the two senses, what rule had we for determining which was the most worthy of credit? How does a blind man correct his errors of touch? If the button be felt double, because pressed by two parts *not* accustomed to feel the same thing at the same time, there must have been a period in the life of every person, when a body pressed by any two parts would have been felt double, by three parts triple, and so on. Nor could sight have corrected those deceptions, if they can be called such; for every thing by the same hypothesis must then have also been seen double. How came we therefore, both to feel and see things single? Surely not by comparing the informations of the two senses together.

But *secondly*; were we to grant, that the sense of touch has originally and constantly informed us that objects are single, it would not follow, that we are thence taught to *see* them also single. For since the place;

which an object seems to either eye to possess, manifestly depends both upon its apparent distance, and its apparent direction from that eye, if visible place be, in the language of Dr. Smith, only an *idea* of real or tangible place, visible direction must bear the same relation to tangible direction ; a consequence of which is, that we can never have a more accurate knowledge of the direction, in which an object may lie from any part of our bodies, by sight than by touch. Facts however prove the contrary. Let any person, for instance, taking a pin in his hand, endeavour, without looking, to bring its head upon a level with either of his eyes ; and there are many chances to one but he will fail in the attempt, of which sight will inform him, when he turns his eye to the object. This to me is a convincing argument, that external bodies are not seen in certain directions, because they have been previously felt in them ; and consequently, that visible place, of which visible direction is a component part, is not merely a representative of the place perceived

ceived by touch. But if the place, in which an object appears to each eye separately, does not entirely depend upon any lesson from feeling, the inference is, that when an object appears in one and the same place to both eyes together, neither is this effect to be attributed solely to the informations of that sense.

Thirdly; in whatever direction an object may appear to either eye, it certainly cannot be seen in the same place by both, except at some point common to the two directions. Dr. Smith acknowledges this, and says, * that when an object is perceived single with both eyes, it is seen at the mutual intersection of the two visual rays; the visible direction of any object coinciding, according to him, with the visual ray, or the principal ray of the pencil which flows from it to the eye. Should we then even allow, that all we know by sight of the places of bodies has been borrowed from feeling, it will still be easy to show, that the rule of vision for each eye, which he
has

has derived from such experience, that of our seeing objects in the directions of their visual rays; is inconsistent with many of the phenomena of sight with two eyes; and consequently, that he has left unremoved the chief difficulty of his subject, which was to explain the single appearance of objects to both eyes, from those laws, or rules of vision, which affect each of them singly. For it is a well known fact, that if two bodies of the same shape, size, and colour, be placed, one in each optic axis, they appear but as one body, provided they be at equal distances from the eyes. Agreeably to the theory of our seeing objects in the direction of their visual rays, this cannot happen, except the united body appear at the intersection of the optic axes. Dr. Smith accordingly, * maintains that it does. Now, in the first place, I appeal to experiment for a direct proof that it does not; and in the second, I observe, that, as the two bodies in the optic axes appear as one, whether they be situated within or beyond the

* Vol. II. Remarks, p. 86.

the concurrence of those lines ; and as a right line joining the bodies, and extended both ways, appears at the same time to the sight as a right line ; it follows, upon admitting the fact which I have denied, that all objects in the plane of the optic axes which are seen in one position and state of the eyes, however near to us, or however remote they may in reality be, must appear to be equally distant, or rather in a line drawn through the concurrence of the optic axes, parallel to the interval between the eyes, and named by opticians the *horopter*. Again, if a right line be made to pass through any part of the plane of the optic axes, at right angles to it, the portions above and below this plane are perceived to be in the same right line with the point which is situated in it, and the whole appears perpendicular to the plane. But the point in the plane is seen, by the last article or proposition, in the horopter ; the whole, therefore, of the perpendicular line must be seen in a plane passing through the horopter at right angles to that of the optic axes ; or in other

words, in the *plane of the horopter*, in which consequently all bodies will have their visible places. But this was the very opinion of Aguilonius, to which he was probably led by a similar train of reasoning; though, as a teacher, he might choose rather to ground it immediately upon an original law of our constitution.

It is probable, however, that Dr. Smith did not perceive the conclusions which might be drawn from his doctrine of objects being seen in the directions of their visual rays, since he has nowhere spoken of them. At any rate, it is manifest he did not admit them, as he has mentioned the following circumstance as a fact,* to which they cannot be reconciled; that, when an object is seen double, both its apparent places are situated between its real place, and the mark at which we look. For, if this were just, together with what he has elsewhere advanced, phenomena ought in many cases to be observed, very different from those which are in truth found to exist. Thus,
for

* Vol. I. p. 48.

for example, if a right line be any where placed in the plane of the optic axes, it follows, from what he has said in one part of his book, that those points of it, through which the axes pass, must be seen united at the mark we look at, the axes crossing each other there; and from what I have just quoted, that every other point must be seen by each eye between its real place and that mark. The appearances, therefore, of all the points, if they do not lie disjoined, but are connected together in some orderly manner, will be arranged in the forms, either of two curves, both passing through the intersection of the optic axes, or of four right lines meeting one another at that point. If the right line be placed nearer to the face than the mark we look at, the apparent lines, whether curved or straight, will approach toward us from their common point, but recede from us, if the real line be situated beyond the mark. Such are the phenomena which ought to follow upon the admission of these two parts of Dr. Smith's theory of vision with two eyes, but

D

which

which are not found to exist in nature.—Aguilonius was at least consistent when he maintained, that all objects are seen in the plane of the horopter; while Dr. Smith, by deserting that opinion in part, seems only to have involved himself the more deeply in error.

Having now said what, I hope, will be thought sufficient to show, that the reason given by Dr. Smith, for our seeing objects single with both eyes, is neither grounded on well-attested facts, nor adequate to the explanation of the phenomena observed, I pass to the examination of the opinion of Dr. Reid.

As this neither rests upon, nor includes any new fact in vision, I need only mention, in order to give an account of it,* that its author maintains with Dr. Smith, that an object is seen in the same place with both eyes, and consequently single, when its pictures fall upon the centres of the retinas, or upon points in them, which are similarly situated with respect to the centres; but
differs

* Inquiry into the Human Mind, c. v. sect. 13.

differs from him in this, that he makes the property to be original, by which any two places in those membranes exhibit only one object, while Dr. Smith derives it altogether from custom.†

In my examination of the opinion of Dr. Smith, I took occasion to remark, that the truth of what distinguished it from all others might remain unshaken, though it were proved, that objects do not appear single, when their pictures occupy any of the corresponding points of the two retinas, since custom might have associated the perceptions of touch, with the sensations of any other parts whatsoever of those membranes. The same observation will not apply with equal justice to the opinion of Dr. Reid. On the contrary, could it be shown, that the places of the two retinas, which repre-

† They differ also with respect to the meaning of a term ; Dr. Smith calling *corresponding points*, such as have the position just mentioned, whether they represent objects single or not ; whereas Dr. Reid says, that those points correspond, whatever their position may be, which represent objects single ; and he appears to me not always to attend to the double use of the same term, when he speaks of the opinions of Dr. Smith.

sent an object single when each receives its picture, are not the centres, or such others as are similarly situated, an obvious inference would be, that the single appearance of the object is not occasioned by a property in those places, bestowed upon them for this special purpose by nature; it being reasonable to expect, that such a property should be found, if any where, in those parts of the retinas which are the most like to each other. I have, therefore, reserved till now, the observations which have occurred to me upon this subject, and which, when stated, must, at least, raise some doubt concerning what has been regarded as true by Dr. Smith and Dr. Reid, and by almost every other writer on vision, since the time of Kepler.

Anatomists have commonly taught, that the centres of the spheres, to which the cornea, the ball of the eye, and the two portions of the crystalline belong, are all placed in the same right line, hence called the optic axis, and that this being produced both ways, passes through the centres of
§ the

the cornea and retina, considered as surfaces. Opticians, on their part; observe, that an object appears single to both eyes, when the axis of each is accurately directed to it; from which they infer, that the centres of the retinas agree in suggesting but one object, though each receives its picture.—Again; since it is known by experience, that, while any object is seen single, to which the optic axes are turned, others at the same distance from the eyes likewise appear so; and since the pictures of these lateral objects fall upon points in the two retinas, equidistant from their centres, and both upon the same side, that is, both to the right or left of the centres, or both above or below them, opticians conclude, that every two places of the retinas, which are similarly situated with respect to the centres, must also agree in exhibiting but one object, though pictures are received by both.

But the whole of this reasoning is built upon a circumstance in the fabric of the eye, which has been shown by some of the
most

most eminent anatomists not to have place, For Varolius * long ago observed, that the crystalline is not situated in the middle of the eye, but more *inwardly*; and the accurate Zinn † has more lately mentioned, that if the eye be divided into a right and left half, the centre of the crystalline will be found in the inner portion. Haller ‡ confirms this fact; and Winslow's || observation, that the centres of the pupil and iris do not coincide, but that the former is nearer to the nose than the latter, is connected with it; since both Zinn and Haller agree, that the centre of the pupil is placed in the axis of the crystalline, while that of the iris is evidently in the common axis of the cornea and globe. Now, a consequence of this position of the crystalline is, that, contrary to what I believe is universally maintained, no ray of light whatsoever can pass unbent to the retina from the atmosphere, or any other medium differing in refractive

* Varolii Anatomia, 12mo. p. 16.

† De Oculo, 4to. p. 127.

‡ Elementa Physiologiae, tom. v. p. 403.

|| Winslow's Anatomy, vol. ii. p. 379. English edition, 8vo.

fractive power from the aqueous humour. If, then, the line joining the centres of the cornea and globe of the eye be what is called the optic axis, and if it be true, that objects appear single when we direct both these axes to them, it must be evident, to such as are acquainted with the common rules of optics, that the pictures of those objects do not fall upon the centres of the retinas, but more internally; and, therefore, that the centres and all the other points of those membranes, which by the present system are supposed to represent objects single, do in fact exhibit them double.

It will be said here, perhaps, that the line * passing from each eye, which we turn to objects when we see them single, is not
a pro-

* I am of opinion, that this line, or at least the line which we turn to objects when we see them most distinctly with one eye, is not the common axis of the globe and cornea. For I find, that, when I place the flame of a candle between either of my eyes, and a plane mirror, in such a manner that it may conceal its own image in the mirror from the sight of that eye, or rather that it may be a little below this image, but in the same vertical plane with it, the image of the flame, seen by reflection from the cornea, does not appear upon the middle point of this coat, but upon that point of it which is opposite to the centre of the pupil.

a production of the common axis of the cornea and globe, but some other, disposed in such a manner, that the pictures of those objects are received by the centres of the retinas. I answer; I readily grant the possibility of the thing, but I assert at the same time, that we have no proof of it, which is a sufficient reason for rejecting every conclusion that depends upon its truth.

Admitting, however, that objects are represented single, when their pictures fall upon the centres of the retinas, or upon any other two points which are equally distant from the centres, and both upon the same side, it appears to me, notwithstanding, to be in violation of all analogy, to ascribe this effect, with respect to the points, at least, on the right and left sides of the centres, to any peculiar property which they possess from nature. For when anatomists find, in a new species of animals, organs similar in structure to those of others they are already acquainted with, they immediately conclude, that they are also similar

lar in regard to their use. In animals of the same species, they believe with certainty, that the organs they see in one have the same properties, as the corresponding organs of another; and, if it be possible, they attribute with greater certainty the same properties to two organs of the like kind, which are found in the same individual. Such is the influence of the rule, that resemblance of property is implied by resemblance of structure. Now it is an universal fact, that if an animal be divided into a right and left half, the corresponding parts of those organs, which exist in pairs, are found at equal distances from the plane of partition. Thus, for instance, in respect to the eyes, the two optic nerves penetrate their outward coat at the same distance from this plane. Their muscles, blood-veffels, and every other of their component parts and appendages, are arranged in the like manner; those nearest to the dividing plane, or the innermost, in the one, being similar in structure to the innermost in the other, the outermost to the outermost, and the inter-

E

mediate

mediate to the intermediate, It is surely, therefore, natural to expect, that such parts should also be similar in their properties; and we in fact find this similarity to exist, wherever it can be clearly ascertained what the properties are. Every person, for example, admits, that the internal straight muscle of the right eye performs the same office, with respect to that eye, as the other internal straight muscle does with respect to the left eye. What judgment are we then to form of the opinion of Dr. Reid, which attributes the same original properties, or rather the joint possession of one original property, to places in the retinas situated at unequal distances from the general plane of partition; which makes an *external* point in one to correspond, in use, with an *internal* point in the other, and this too by a principle implanted by nature? If such things exist, they may, at least, be said to stand opposed to a most extensive analogy.

To these arguments, *a priori*, against the opinion of Dr. Reid, I shall now add others, which

which are derived from a consideration of its consequences.

First; Since visible place, as was formerly observed, includes in it visible distance, it is evident that, if both eyes, by virtue of an original property, see an object in the same place, distance must also be originally perceivable by sight. Dr. Reid, * however, has himself so ably shown, that we would never have acquired, by means of our eyes, any knowledge of distance, unless they had been assisted by the sense of feeling, that I forbear to say any thing more upon this head, than that the existence of no property can be admitted, which leads to the conclusion I have stated.

Secondly; If distance be not immediately perceivable by sight, the only manner, in which an original property of the eyes can affect the visible places of bodies, is by occasioning them to appear in certain directions. Now Dr. Reid maintains, † that every external point is seen in the direction of a line

* Inquiry into the Human Mind, chap. 6. sect. 3 & 20.

† Ibid. chap. 6. sect. 12.

passing from its picture on the retina, through the centre of the eye. If, therefore, this direction be the same as that suggested by the original property so often mentioned, the latter law is merely another expression for the former, and ought to be rejected as superfluous. If it be different, and should the two laws exist together, objects seen with both eyes might sometimes appear quadruple, sometimes triple, but never single. Were they to exist successively, one when we employ one eye, the other when both, an object, though at rest, should always appear to move when viewed alternately by one and by both eyes ; neither of which conclusions is agreeable to experience.

Thirdly ; To show in a different way, and one perhaps more easily understood, that the opinion of Dr. Reid is not consistent with the phenomena of vision it ought to explain, I shall suppose an experiment to be made upon a person who squints. But I must premise, that it appears, from the

observations of Dr. Jurin * and himself, † that all such persons have one eye of a weaker sight than the other; that when both eyes are open, the weaker is turned away from objects, which are attentively viewed; but that when the strong eye is closed, the weaker is pointed to objects, exactly as the former would be in the same situation; and that it likewise perceives them in similar directions. Let now the ordinary position of the person's eyes, upon whom the experiment is made, be such, that the optic axes intersect each other about an inch or two from the face; and while the other is closed, let the flame of a candle be placed in the axis of the weak eye, which I shall call the left, at the distance of some feet from it, and on the right side of the body. The flame will consequently appear in the same direction, as if his eye had no fault, and will be seen on his right, where it is in reality situated. Both eyes retaining the same position with respect to his head and each other, let the

weak

* Smith's Optics. Vol. 2. Remarks, p. 30.

† Inquiry, chap. 6. sect. 16.

weak eye be afterward shut, and the right opened, and let another object be placed in the axis of the latter, an opaque body being at the same time so disposed, as to hide from it the candle which is in the axis of the left eye. This object in the right axis will consequently appear on the left side. Now, since the two objects, which have been thus viewed separately, are situated, one to the right, and one to the left; and since they have been also *seen* in those positions, their visible places must be two, as well as their tangible, and must be remote from each other. How then should these objects appear, if, instead of being viewed alternately, each by the eye in the axis of which it is placed; they were seen by the two together; the positions and internal states of the eyes being in both cases the same? Dr. Reid must answer; They will possess but one visible place, since their pictures fall upon the centres of the two retinas, points endowed with the original property of representing objects single. But where is this one place

to

to be round? In the axis of the right eye, or in that of the left, or between the two? In any of these cases, or in any other that can be imagined, the law of visible direction, so much insisted upon by Dr. Reid, that objects appear in the perpendiculars to their pictures upon the retina, and in truth every other law of visible direction hitherto published, must be suspended with respect to one or both eyes; unless, indeed, the united object be referred to the intersection of the optic axes, about an inch or two from the face. This, I believe, Dr. Reid would not readily admit; but if he should, another case of squinting may be imagined, in which the optic axes recede from each other, and where the same reasoning will apply without the possibility of its force being thus eluded. It now remains for me to mention, that the experiment here stated by the way of supposition, in which the optic axes cross each other near to the face, was actually made by Dr. Reid, with this result, that the two objects appeared in different places, when seen by both eyes together;
and

and that the other experiment, in which the optic axes are supposed to diverge, was made by myself, with a similar event. Dr. Reid, however, instead of being led, by the termination of his experiment, to impute a fault to the principle from which he had expected a different one, concluded from it, that there was something unnatural, beside the squinting, in the person's eyes, upon whom it was made; though it had been previously ascertained, that objects appeared in the ordinary manner to each of them, when separately employed.

My examination of the second class of opinions, respecting the cause of the single appearance of objects to two eyes, being finished, some person, perhaps, will now say; Granting that no error can, at first sight, be shown in your arguments against those of Dr. Smith and Dr. Reid, is it not a sufficient reason for believing them fallacious, that they prove too much? If objects appear single neither from custom, nor an original property of the eyes, have we not an effect without a cause, and must there
not

not be something wrong in the facts or reasoning which lead to such a conclusion? The answer I make is as follows : Since visible place contains in it both visible distance and visible direction, it is not necessary that the single appearance of an object, to both eyes, should depend altogether either upon custom, or an original principle of our constitution ; for its visible distance to each eye may be learned from feeling, and its visible direction be given by nature ; in which case, the unity of its place to the two eyes, will be owing to neither of those causes singly, but to a combination of both ; and this I regard as a sufficient reply.

THE END OF PART I.

E

AN

AN
E S S A Y
UPON
SINGLE VISION WITH TWO EYES.

PART II.

Of a new Theory respecting Visible Direction, and of a Solution hence derived of the Question, why Objects are seen single with two Eyes.

I NOW proceed to offer a new opinion, why objects are seen single with two eyes ; or in other words, why they appear in the same place to both, this being the light in which I view the fact to be explained.

In every part of natural philosophy, accidents often lead to discoveries, which reason alone might not easily have reached. Under this cover I hope to shelter myself from the charge of presumption, in venturing

turing to give the solution of a problem, upon which the talents of many persons of great learning and genius, have been unsuccessfully employed ; for should I prove more fortunate than such men have been, this must be attributed to the knowledge of a circumstance I observed by chance, in repeating some very common experiments.

The visible place of an object being composed, as I have already several times remarked, of its visible distance and visible direction, to show how it may appear the same to both eyes, it will be necessary to explain, in what manner the distance and direction, which are perceived by one eye, may coincide with those which are perceived by the other : and first with respect to the distance.

In judging of distance by sight, we frequently make considerable mistakes, even when the objects are not very remote ; but no person, I believe, has ever observed, that while an object seemed to one of his eyes at a certain distance, it has appeared to the other to be at a different distance, and from this circumstance alone has been seen double;

or, to express the same thing in another way, that while the visible appearance of an object to one eye, covered the visible appearance of the same object to the other eye, the two appearances did not seem entirely to coincide, and make one, but were seen separate by the two eyes. I do not stop to give the reason of this fact, which must be plain to those who are acquainted with Bishop Berkeley's theory of visible distance; but proceed to mention, that the difficulty in finding a true and sufficient cause for the union of the two visible places of one or two objects to two eyes, must therefore consist altogether in showing, in what manner the two apparent directions may coincide, consistently with the attending phenomena.

Since Kepler's great discovery of the seat and manner of vision, there have been, as far I know, only two theories offered respecting the apparent directions of objects. One is, that they are perceived in the direction of lines passing from their pictures on the retina, through the centre of the

eye; * the other, that their apparent directions coincide with their visual rays. But both of these theories are inconsistent with the phenomena of single vision with two eyes. For according to neither of them can an object, placed at the concurrence of the optic axes, be seen single, unless we have a most accurate knowledge of its distance; nor will either admit two objects to be seen as one, which are situated in the optic axes, whether on this side, or beyond where they meet, unless the united object be referred by sight to their very point of intersection; both of which conclusions are contradicted by experience. It is evident, therefore, that some other theory of visible direction is required, which shall not be liable to these objections; and such a theory, I hope, I shall bring forward in the following propositions,

* Mr. D'Alembert has said (*Opuscules Mathematiques*, Tom. I. p. 265) that all optical writers before him had regarded it as an axiom, that every visual point is seen in the direction of its visual ray. But the assertion is not well founded. For Kepler long ago taught (*Paralipomena in Vitellionem*, p. 173) that objects are perceived in lines passing from their pictures upon the retina, through the centre of the eye; in which he was followed by Dechales and Doctor Porterfield; to the latter of whom Dr. Reid improperly attributes the discovery of the *same* supposed law.

sitions, after mentioning the meanings which I affix to several terms I shall frequently employ.

EXPLANATION OF TERMS.

I. When a small object is so placed with respect to either eye, as to be seen more distinctly than in any other situation, I say it is then in the *optic axis*, or the axis of that eye; and if another small body be interposed between the former and the eye, so as to conceal it, and if a line joining the two be produced till it falls upon the cornea, I call this line the *optic axis*, or the axis of the eye; leaving for future determination the precise point of the cornea it falls upon, or what part of the retina receives the picture of an object which is placed in it.

II. When the two optic axes are directed to a small object not very distant, they may be conceived to form two sides of a triangle, the base of which is the interval between the points of the corneas, where the axes enter the eyes; but if the object be

be very distant, then they may be supposed to be two sides of a parallelogram, whose base is the same interval. To avoid circumlocution, I shall call this interval the *visual base*.

III. If there be drawn a line from the middle of the visual base, through the point of intersection of the optic axes, or parallel to them, if they be parallel to each other, I name it the *common axis*.^{*} This term I believe was invented by Alhazen ; but with him it signified a line drawn from the centre of the junction of the optic nerves, through the middle of the interval between the centres of the retinas.

Such

^{*} It may be said, perhaps, that as I do not define the points of the corneas, upon which the optic axes fall, I cannot, with propriety, desire the line which connects them to be divided. To this I answer, that it is not necessary for the purpose I have mentioned, that they should be defined ; if it be granted to me, and I think it cannot be refused, that upon whatever point of the right cornea the right axis falls, the left axis will fall upon a similarly situated point of the left cornea ; that is, if this point of the right cornea be at any given distance from its middle, and upon the inside of it, the corresponding point of the left cornea will be at the same distance from the middle of this, and also upon its inside. Whatever extent, therefore, the line connecting these places of the corneas may have, its middle point will be the same.

Such a line was consequently immoveable. As the term, however, is not in modern use, no mistake can arise from confounding the two meanings, and the reason will soon be seen, why I employ it in the sense I have mentioned. Those who are acquainted with the writings of the older opticians will perceive, that I give it nearly the same signification as they did to their *common radius*.

PROPOSITION I.

Objects situated in the Optic Axis, do not appear to be in that Line, but in the Common Axis.

EVERY person knows, that, if an object be viewed through two small holes, one applied to each eye, the two holes appear but as one. The theories hitherto invented afford two explanations of this fact. According to Aguilonius, Dechales, Dr. Porterfield and Dr. Smith, the two holes,
or

or rather their borders, will be seen in the same place as the object viewed through them, and will consequently appear united, for the same reason, that the object itself is seen single. But whoever makes the experiment will distinctly perceive, that the united hole is much nearer to him than the object ; not to mention, that any fallacy on this head might be corrected by the information from the sense of touch, that the card, or other substance, in which the holes have been made, is within an inch or less of our face. The other explanation is that furnished by the theory of Dr. Reid. According to it, the centres of the retinas, which in this experiment receive the pictures of the holes, will, by an original property, represent but one. This theory, however, though it makes the two holes to appear one, does not determine where this one is to be seen. It cannot be seen in only one of the perpendiculars to the images upon the retinas, for no reason can be given why this law of visible direction, which Dr. Reid thinks established beyond dispute, if it ope-

rates at all, should not operate upon both eyes at the same time ; and if it be seen by both eyes in such lines, it must appear where those lines cross each other, that is, in the same place with the object viewed through the holes, which, as I have already mentioned, is contrary to experience. Nor is it seen in any direction, the consequence of a law affecting both eyes considered as one organ, but suspended when each eye is used separately. For when the two holes appear one, if we pay attention to its situation, and then close one eye, the truly single hole will be seen by the eye remaining open, in exactly the same direction as the apparently single hole was by both eyes.

Hitherto I have supposed the holes almost touching the face. But they have the same unity of appearance, in whatever parts of the optic axes they are placed ; whether both be at the same distance from the eyes, or one be close to the eye in the axis of which it is, and the other almost contiguous to the object seen through them. If a line, therefore, be drawn from the object to one of the eyes, it will represent all
the

the real or tangible positions of the hole, which allow the object to be seen by that eye, and the whole of it will coincide with the optic axis. Let a similar line be drawn to the other eye, and the two must appear but as one line; for if they do not, the two holes in the optic axes will not, at every distance, appear one, whereas experiments prove that they do. This united line will, therefore, represent the visible direction of every object situated in either of the optic axes. But the end of it, which is toward the face, is seen by the right eye to the left, and by the left eye as much to the right. It must be seen then in the middle between the two, and, consequently, in the *common axis*. And as its other extremity coincides with the point where the optic axes intersect each other, the whole of it must lie in the common axis. Hence the truth of the proposition is evident, that *objects, situated in the optic axis, do not appear to be in that line, but in the common*

Many other experiments might be mentioned which demonstrate the same thing. If, for example, the head of a pin, or of a needle, be interposed between each eye, and any small object to which both the optic axes are directed, the heads of the two pins or needles will constantly appear as one in the common axis. When the heads, however, are near to the eyes, this experiment is not so satisfactory as the former, since, in these positions, they seem as broad transparent shadows, for reasons known to every person a little conversant in optics ; whereas the holes appear well defined, though almost touching us. Again ; if we hold two thin rulers in such a manner, that their sharp edges shall be in the optic axes, one in each, or rather a little below them, the two edges will be seen united in the common axis, and this apparent edge will seem of the same length with that of either of the real edges, when seen alone by the eye in the axis of which it is placed. If instead of two rulers we employ two strings of different colours, as red and green, the like unity of appearance will

will be observed. But in this experiment it frequently happens, that, contrary to what we might naturally expect, only one of the strings is seen at a time. When, however, only one is seen, its apparent situation is exactly the same as that of the string, compounded, if I may so express myself, of the two when seen together ; and hence we have a convincing proof, if any were wanted, that the single appearances of objects must depend upon some law of visible direction affecting each eye, when employed by itself, in the same manner as when it is used conjointly with the other.*

PRO-

* Du Tour expected, that if two objects of different colours were seen in the same place by both eyes, which however he says, he was never able to observe, the colour of the apparently united object would be compounded of those of the two really single objects. *Memoires des Savans Etrangers*, Tom. iv. p. 500. And Dr. Reid mentions expressly that it is so compounded. *Inquiry*, p. 293. But in all my experiments upon this subject I have remarked, that, when the two objects appeared united, each was seen, notwithstanding, in its proper colour ; the red, for example, appearing as it were through a transparent green, and the green, in the same experiment, as through a transparent red. Nor is there any thing in this inconsistent with the received doctrine of the composition of colours. For in every instance of the production of a new colour, from rays of different colours being at
same

PROPOSITION II.

Subjects, situated in the Common Axis, do not appear to be in that Line, but in the Axis of the Eye, by which they are not seen.

THE facts which demonstrate the truth of this proposition, are both numerous and common.

same time sent to the eye, these rays fall upon the same sentient extremities of the same nerve. But, in the case before us, the differently-coloured rays fall upon the sentient extremities of two different nerves, which have no communication with each other, except through the medium of the brain. We have greater reason, therefore, for expecting, that the colours impressed upon the two eyes, should be perceived uncompoundd, than there is for two colours being perceived separately, which are impressed upon two different parts of the same eye.

From the fact of the two colours being thus perceived distinct from each other, I would infer, by analogy, a mode of argument indeed often fallacious, that if it were possible for us to hear any one sound with one ear only, and another sound with the other ear only, such sounds would in no case coalesce either wholly or in part, as two sounds frequently do, when heard at the same time by one ear; that consequently, if the sounds of one musical instrument were to be heard by one ear only, and those of another, by the other ear only, we could have little or no perception of harmony from such sounds; and that, if in any succession of sounds emitted by one instrument, we were to hear the 1st, 3d, 5th, and so on, by one ear only, and the 2d, 4th, 6th, and so on, by the other ear only, we would be deprived, in a considerable degree, of the melody of such sounds, as this

common. If a piece of wire, or any other substance, representing a physical line, be placed in the common axis, with one of its extremities near to the visual base, and if both the optic axes be directed to its farther or distant extremity, instead of one, two wires will be seen, meeting each other at their farther ends, and gradually diverging as they approach the face, till they apparently terminate at the eyes. If the right eye be closed, the wire which seemed to terminate at the left eye, disappears; and if the left eye be closed, then the other wire disappears; whose termination was at the right eye. The real wire, therefore, in the common axis, appears to the right eye to be situated in the axis of the left, and to the left eye to be situated in the axis of the right, agreeably to what the proposition asserts.

The following experiments will illustrate and confirm both this and the preceding proposition.

seems to depend in a great measure upon a new impression being made upon the auditory nerve by one sound, before the impression of the sound immediately preceding has passed away,

proposition. Through a piece of card, or pasteboard, let two small holes be made, the interval between which is such, that while a very remote object is seen through one of them by the right eye, the same object may be seen through the other by the left eye. Make afterward another hole in the card, or pasteboard, exactly in the middle between the two former; and let the object be viewed through them as before. These, or the outer holes, will now appear one, precisely where the sense of feeling indicates the middle hole to be; while the middle hole will appear as two, which seemingly occupy the places of the real outer ones. The two appearances of the middle hole, which is placed by construction in the common axis, are therefore seen in the optic axes; and as the left is not seen when the right eye is shut, nor the right when the left eye is shut, each appearance is observed in the axis of the eye, by which it is not seen. As I have supposed the distance between the outer holes to be adapted to the interval of the

together or successively, as was formerly mentioned, and at right angles to the visual base, or edge of the table; in short, exactly in the situation, which the yellow string in reality possesses; and the yellow string, instead of appearing single in the common axis, and perpendicular to the visual base, will now be seen as two, each inclined to the base; that seen by the right eye, apparently occupying the place in reality possessed by the green string, and that seen by the left eye, the place of the red string.

PROPOSITION III.

Objects, situated in any Line drawn through the mutual Intersection of the Optic Axes to the Visual Base, do not appear to be in that Line, but in another, drawn through the same Intersection, to a Point in the Visual Base distant half this Base from the similar Extremity of the former Line, towards the left, if the Objects be seen by the Right Eye, but towards the right, if seen by the Left Eye.

TWO cases of this proposition have already been proved. For it has been shown
by

by the first proposition, that objects, placed in the axis of either eye, appear to it to be situated in the common axis. But the common axis is a line drawn thro' the mutual intersection of the optic axes to the visual base, and its termination there is distant, by construction, half that base, from the similar terminations of the axes of both eyes, to the left of the right axis, and to the right of the left. Again, it has been shown by the second proposition, that objects, placed in the common axis, appear to each eye to be situated in the axis of the other; and the terminations of both optic axes, at the visual base, are distant half this base, from the similar termination of the common axis, the left being to its right, and the right to its left.

Let it now be supposed that two objects, one placed in the axis of either eye, the right, for instance, and the other in the common axis, be viewed at the same time by that eye, it is evident that the visible directions of both will be equally removed to the left, from their real positions. But

such an alteration of visible direction, from real position, cannot be imagined to happen, with respect to objects placed in the optic and common axes, unless a similar effect be, at the same time, produced upon such as are situated any where between those lines, or in their vicinity. Facts confirm this: If a line, for example, be drawn through the intersection of the optic axes to a point in the visual base, exactly in the middle between the terminations there of the right and common axes, its apparent situation, to the right eye, will be found to have the same relation to the apparent situations of lines placed in the right and common axes, as its real situation has to the real situations of such lines. And the like will be found, by observation, to be true of every other line, which may be drawn through the point of intersection of the optic axes to the visual base.

The whole of what has here been said may be illustrated and confirmed, by having again recourse to the experiments with strings of different colours. In formerly
describing

describing those experiments, I did not mention all the appearances which occurred upon making them, but only such, as had immediate reference to the points then under consideration. When, for instance, a red string was placed in the axis of the right eye, and a green one in that of the left, I said that they both appeared in the common axis. But this is not the only phenomenon to be observed with respect to their apparent number in this experiment. For as the red string is also seen by the left eye, and the green by the right, two other strings become visible, beside that in the common axis, the apparent positions of both of which will be found to be the same with those, which ought to follow from the present proposition. Should now a yellow string be placed between the two former, as in the proof of the second proposition, its appearance to the right eye will bisect the space between the appearances of the red and green strings to that eye; and the like will be true with respect to the appearances of the three strings to the left

5

eye,

eye, agreeably to what the same proposition teaches us to expect.

I believe I need scarcely remark, that, although in most of the proofs and illustrations of these propositions, I have confined myself to the visible appearances of lines *between* the intersection of the optic axes and the visual base, the same things, however, must be equally true of those lines, when they are produced beyond the intersection, with this difference only, that, while the portions within, seem, to the right eye, to be farther situated to the left than they really are, but to the left eye farther to the right, the portions beyond the intersection will seem to the right eye to the right of their real positions, but to the left eye to the left of them. For it is manifest, that, if a line be seen by one eye in a certain direction, a prolongation of it must be seen in the same direction; and that, if a line be made to turn upon any point in itself, the two extremities must move contrary ways.

Should the optic axes be parallel to each other, the same proofs and illustrations will
still

still apply, since we may here suppose them to meet at an infinite distance from the visual base. In this case, the visible appearances of lines, drawn from this supposed point of intersection to the visual base, will be parallel to the real lines, and distant half this base from them, through their whole extent.

AS I have thus, I think, sufficiently proved, that the apparent directions of objects are governed by a law, different from any which has hitherto been thought to exist, I shall now proceed to state, in a few words, in what manner the phenomena of single and double vision with two eyes are dependant upon it.

I formerly mentioned, that, since an object is never seen double, merely from its being seen at different distances by the two eyes, the only difficulty in explaining its single appearance consists in showing how its two visible directions may coincide, consistently with the attending phenomena. But we are enabled to do this, with the utmost

most ease, by the theory I have endeavoured to establish. For, if the question be concerning an object at the concurrence of the optic axes, I say it is seen single, because its two similar appearances, in regard to size, shape, and colour, are seen by both eyes in one and the same direction, or, if you will, in two directions, which coincide with each other through the whole of their extent. It therefore matters not, whether the distance be truly or falsely estimated; whether the object be thought to touch our eyes, or to be infinitely remote. And hence we have a reason, which no other theory of visible direction affords, why objects appeared single to the young gentleman mentioned by Mr. Cheselden, immediately after his being couched, and before he could have learned to judge of distance by sight.

When two similar objects are placed in the optic axes, one in each, at equal distances from the eyes, they will appear in the same place, and therefore one, for the same reason that a truly single object, in the concurrence of the optic axes, is seen single.

Here

Here again, as the two visible directions coincide in every point, it is not necessary that the united appearance should be judged to be at any particular distance; that it should be referred, for instance, to the concurrence of the optic axes, where the two other theories of visible direction are obliged to place it, in opposition to the plainest observations.

Objects, any where in the horopter, will be seen single, because their apparent directions to the two eyes will then completely coincide. And for a contrary reason, those placed in any other part of the plane of the optic axes will appear double. To make these things evident, let a line pass through the point of intersection of the optic axes, and any given object, to the visual base, which is to be produced, if necessary; and let it be called the line of the object's real position. Take afterward, in the visual base, or its production, two points, one on each side of the line of real position, and both distant from its termination there, half the visual base. Lines drawn from these

I

points

points, through the point of intersection of the optic axes, must consequently contain the two visible positions of the object. But when this is situated in the horopter, the line of real position will coincide with the horopter, and will not therefore reach the visual base, unless at an infinite distance from the eyes. For which reason, the two lines, containing the visible positions of the object, must fall upon the visual base at a like distance, and must consequently be regarded as coinciding with each other. When the object is not in the horopter, the two lines of visible direction will be found, by the same means, not to coincide.

That I might simplify a matter, which under my management, must, I fear, still be of difficult apprehension, I have, in expressing the law of vision, so frequently mentioned, purposely confined it to objects situated in the plane of the optic axes. But in persons who do not squint, or whose eyes are not distorted by external violence, the two appearances of an object, seen double, are always, either in that plane,
or

or in some one parallel to it; so that, if the visual base be parallel to the horizon, a line joining the two appearances will, in every case, be also parallel to the horizon. Whoever then is able to explain, why objects in the plane of the optic axes appear either single or double, may readily give a reason for the like appearances of such as are placed any where else. Not to spend much time, therefore, upon this part of the subject, I shall shortly observe, that if planes be supposed to pass through the two optic and common axes, perpendicular to that in which they all lie, and if two lines be drawn from any point of the common intersection of the former planes to the visual base, one along each of the perpendicular planes which pass through the optic axes, these two lines will appear as one, in the perpendicular plane of the common axis; the single visible line, however, possessing the same elevation, in regard to the horizon, as the two real lines: And again, that, if a line^{be} drawn from any point of the same intersection to the visual

base, along the perpendicular plane of the common axis, it will appear as two, one in each of the planes which pass through the optic axes; the two visible lines having the same inclination to the horizon in their progress to the visual base, as the real single one. In this manner, every thing may be shown to be true, with respect to the single and double appearances of objects without the plane of the optic axes, which has already been done with regard to those placed in it. But farther; since any point, taken at pleasure, in the common intersection of the three perpendicular planes, appears single, the whole of the line of intersection must appear so, and likewise every point of a plane made to pass through it, parallel to the visual base. Such a plane necessarily includes the horopter, and is the same as that, which is called by Aguilonius the plane of the horopter.

To exemplify the principal property of this plane, I shall mention an experiment, which at first I did not understand, though
the

the result was a direct consequence of my own principles. I suspended a fine chord at right angles to the horizon, and retreating a step or two, I looked steadily at a point in it, which was upon a level with my eyes. The chord, in these circumstances, appeared single; but whenever I directed my eyes to any other point of it, either above or below the former, two chords would appear, crossing each other at the part, to which the eyes were directed. In the first case, the whole chord was in the plane of the horopter, but in every other, only that point of it to which both eyes happened to be turned. A conclusion from this experiment is, that no object, which is truly perpendicular to the horizon, will appear to be so, while our bodies are erect, unless we direct our eyes to a point in it exactly upon a level with themselves.

It was once my intention to subjoin here several instances, from the most approved authors, of inaccurate descriptions of the single and double appearances of objects; in order to show, that the theory of
visible

visible direction, which I have advanced, is not only consistent with the universally received facts, but that it also discovers to us, some minute errors, which unguided sense has committed upon this subject; it being, perhaps, one of the surest tests of the soundness, as well as one of the chief uses, of theories in philosophy, that they lead to the knowledge of what, otherwise, might have remained for ever hidden. But fearing I have already proved tiresome, I give up this design, and hasten to the consideration of some consequences from my theory, which seem to me both curious and important, and which, when first mentioned, may appear to carry with them their own refutation.

END OF PART II.

AN
E S S A Y
UPON
SINGLE VISION WITH TWO EYES.

P A R T III.

*Of some Consequences from the foregoing Theory of Objects
being seen single with two Eyes, together with the
Explanation of several other Phenomena
of Vision.*

IT has hitherto, I believe, been thought by opticians, that, if the position of the eye be unchanged, the visible direction of an object will be the same, as long as its picture occupies any one point of the retina; and that, in every different position of the eye, a picture, which continues to occupy the same point of the retina, will represent its
5 object

object in a different direction. But if the theory be just, which I have advanced in the preceding part of this Essay, neither of those opinions can be universally true. For it follows, from what was there mentioned, that if one of the optic axes be kept fixed, and the other be at different times variously bent toward it, objects, though situated in the fixed axis, will nevertheless change their visible directions, with every variation of the moveable axis; since they must always appear in the common axis, which alters its position with every change of the moveable axis: And again, that, if the two optic axes should vary their inclinations to each other in such a manner, that the common axis, may, notwithstanding, remain fixed, an object placed in either optic axis, and following it in every motion, will possess but one visible direction, in all this variety of real positions. That these conclusions from my theory, or rather parts of it, are true in fact, I can assert upon the authority of observations, and I shall now attempt to
trace

trace them both to a common principle, by means of some experiments, which were instituted with a very different view.

When we have looked steadily for some time at the flame of a candle, or any other luminous body, a coloured spot will appear upon every object, to which we shortly after direct our eyes, accompanying them in all their motions, and exactly covering the point, which we desire to see the most accurately. Whatever therefore can be proved concerning the apparent direction of such a spot, in any given position of the eyes, must likewise be true in the same position of the eyes, with regard to the apparent direction of an object, situated at the concurrence of the optic axes; as its pictures must occupy, in this case, the very parts of the retinas, upon the affections of which the illusion of the spot depends. This being premised, I shall now relate one or two observations, respecting the apparent directions of the spot, and consequently upon those of external objects,

K

which

which, as far as I know, have not been mentioned by any other person.

1. The spot is always seen single, whether the surface, upon which it is projected, be touching the face, or at the greatest distance from us; and the reason is plain. For the parts of the retinas, by whose affections from the luminous body it is occasioned, are those likewise which receive the pictures of objects, placed at the intersection of the optic axes; and as such objects always appear single, so must also the spot. The fact indeed is so open to observation, and its cause so easily shown, that I should scarcely have thought of mentioning it, had not Dr. Darwin* lately told us, that the spot is seen double, as often as the eyes are directed to an object more or less distant than the luminous body which gave rise to it. With respect to our different assertions upon this point, I shall only say, that I have made the experiment, I believe, upward of an hundred times, uniformly
with

* Philosoph. Transact. for 1786, p. 318. Dr. Darwin indeed, says, p. 341, that Buffon had observed the same fact; *but* it is evident he has mistaken that author's meaning.

with the same result; and that, if the spot ever appears double, this must be from some cause very wide of a change in the mutual inclination of the optic axes, to which he attributes it.*

2. The spot not only appears single in every ordinary position of the optic axes, but cannot even be made to appear double, by any means whatsoever. If it be projected, for example, upon a piece of white paper, whoever makes the trial will find, that, although, on pressing one eye upward or downward, or to either side, the paper will be seen double, yet the spot

* The only way, in which I think it possible for the spot to appear double, consistently with the universally acknowledged fact, that an object at the intersection of the optic axes is always seen single, is this, that, when the intersection is near to the face, an object placed in it shall not send its pictures to the same points of the two retinas, as it does, when the intersection is more remote. And such I once hoped to find to be the case; for I had formed, upon the supposition of its truth, a more plausible account of the manner in which the eyes are fitted to receive, successively, pictures equally distinct from objects at different distances, than any I had met with. But, after many experiments to ascertain the matter, I was obliged to return to the common opinion, that the picture of an object in the optic axis, whatever be its distance from the eye, is always received upon the same point of the retina.

will always appear single, and to possess its former place on the paper, as seen by the eye, which is not disturbed. Before I knew the result of this experiment, I had imagined, that, the position of one eye being forcibly altered, the external situation of the spot, which was suggested by the affection of that eye, would likewise be altered, and the spot by consequence be seen double. As the event, however, was contrary to my expectation, I began to suspect some cause of fallacy had been overlooked, which at length I thought might be this, that the spot had been seen by that eye only whose position was not disturbed, the violence, suffered by the other, interrupting the due exercise of its functions. To determine, therefore, whether my conjecture was well founded or not, I made another experiment, which is mentioned in the following article :

3. Having looked steadily for some time at the flame of a candle, with *one* eye only, I directed afterward, with both eyes open, my attention to the middle
of

of a sheet of paper, a few feet distant ; the consequence of which was, that a spot appeared upon it in the same manner, as if I had viewed the flame with both eyes, though somewhat fainter. My attention remaining fixed upon the sheet, I now pushed the eye, by which the spot was seen, successively upward and downward, to the right and to the left, and in every oblique direction ; the spot however never altered its position, but kept constantly upon the middle of the appearance of the paper, perceived by the undistorted eye, though the appearance of the paper to the distorted eye, was always separate from the former, and the sheet consequently seen double. My conjecture, therefore, was proved to be ill grounded, and all suspicion of fallacy in the former experiment ~~was~~ ceased.

Now it is evident, from these two last experiments, that the situation of the spot does not depend upon the bare position of the eyes, or else, in the former of them, it would have appeared double, and in the latter,

latter, it would have been moved from the middle of the paper, when the only eye by which it was seen was pushed from its place. Neither can it depend upon the bare position of the muscles of the eye, as these were also moved in the same experiments; nor upon any affection whatever of the optic nerve. For since this last substance is altogether passive, even in those motions of the eyes which do occasion a change of the spot's situation, every alteration, induced upon the nerve by those motions, must be ultimately ascribed to a change of its position; and we have seen, that similar changes of its position have been produced by external violence, without any alteration of the spot's situation. The apparent situation of the spot being, therefore, dependant upon none of these circumstances, and being at the same time affected by the *voluntary* motions of the eye, it must, I think, be necessarily owing to the *action* of the muscles, by which these motions are performed. Assuming then as true, that the apparent direction of an object, which sends its picture
to

to any given point of the retina; depends upon the state of action existing at the same time in the muscles of the eye, and consequently that it cannot be altered, except by a change in the state of that action, I shall proceed to trace to this principle, several phenomena of vision, particularly the uniform singleness of the spot already described, and the two facts respecting the visible directions of objects in the optic axis, which were mentioned in the beginning of this part of my Essay.

The thing itself is universally acknowledged, though a dispute has arisen whether custom or an original property be the cause, that every voluntary motion of one eye, in persons who do not squint, is attended with a corresponding motion in the other. Now as all voluntary motions are produced by muscular action, it follows, that every state of action, in the muscles of one eye, has its corresponding state in those of the other, and that the two are constantly conjoined. When, therefore, the spot appears single to both eyes in their

free positions, the states of action in the muscles must be such, that the direction, in which it is seen by one eye, coincides with that in which it is seen by the other. But, if we push one eye from its place, no change is hereby made in the action of its muscles; for the state of action in those of the free eye is confessedly the same as it was; and it will be attended with a corresponding state in those of the distorted eye; in proof of which it may be observed, that, whenever the pressure is removed, the distorted eye immediately returns to its former position, without the aid of any new muscular effort. The conclusion then is, that, since there has been no alteration in the action of its muscles, neither ought there to be any in the direction of the spot seen by it, which is the fact to be explained.

Hence also is to be derived the true reason, why objects appear double, when one eye is pushed from its place. For as their pictures must fall upon points of the retina in this eye, different from what they
formerly

formerly possessed ; and as no change is made, by the distortion, upon the visible direction, suggested by any part of the retina, the objects will be seen by the pressed eye, exactly in the same directions as they would have been, *before* it was pressed, had the pictures *then* fallen upon the points of the retina, which they *now* occupy. They must therefore be *now* seen in different directions by the two eyes, and consequently double. An experiment with a contrary event will confirm this explanation, and likewise show more clearly, in what I differ from those who have endeavoured to account for the same fact. Both eyes being open, let one of them be pushed from its situation, and let two similar objects, such as two pieces of money of the same metal and stamp, be afterward so placed, that one shall lie in each optic axis ; these two objects will now appear to be one, and the object so compounded will be seen in the place, to which the undisturbed eye refers the truly single object lying in its axis.

L

Another

Another inference from this doctrine is, that, if the eyes are in any very unusual position with respect to each other from the action of their own muscles, as in persons who squint, two objects placed in the optic axes, one in each, will not appear as one object; for each will be seen in the direction, which is determined by the state of action in the muscles of the eye, upon whose retina its picture falls; and as this state, in one eye, does not correspond with that in the other, the directions cannot coincide. This conclusion is verified by the result of an experiment of Dr. Reid upon a person, affected with strabismus, and by that of another, made by myself, both of which have been already related.

To explain, therefore, why an object in the optic axis appears at different times in different directions, though the axis be kept fixed, it is only necessary to show, that, whenever this happens, a change, notwithstanding, occurs in the actions of the muscles which move the eye. With this view, I observe, that the motions of that
organ

(75)
organ may be divided into two sets; the first, consisting of those, by which one eye is carried along with the other, upward and downward, to the right and to the left, and in every oblique direction, the interval between the pupils remaining constantly the same; the second, of the motions of the pupils, or the anterior parts of the eyes, to and from each other. Supposing now, that both the optic axes are perpendicular to the visual base; should the left axis be afterward inclined to the right side, the natural tendency of the right axis is to incline equally to the same side, so as to preserve its former parallelism to the left. This tendency, however, in the right axis to follow the left, may be counteracted by an effort of the muscles, which regulate the interval of the pupils, until the two axes intersect each other within two or three inches of the face. But it is evident, that the same degree of muscular force will be required to retain the right eye in its original position, as is necessary to give to the left eye its motion toward the right; and

hence, that, in every different inclination of the left axis to the right, an object placed in the latter, though its real position be unchanged, will, nevertheless, appear in a different direction, in consequence of the different state of action in the muscles of the right eye, which accompanies every new degree of inclination of the axes to each other. As the object must always appear in the common axis, the alteration, in this example, of its visible direction, from an *increase* of the mutual inclinations of the optic axes, will be from left to right; but when the inclination decreases, from right to left. If the right axis be the one which is moved, and the left fixed, the alterations of visible direction in an object placed in the latter, from similar changes in their inclinations, will be contrary to those which have just been mentioned.

The reason also can now be made to appear, why an object, preserving constantly its place in the optic axis, may, in a considerable variety of its real positions, possess but one visible direction. For, in such cases,

cases, the change of its visible direction, which might be expected to accompany the motion of the eye in the axis of which it is situated, is prevented from occurring, by a tendency to a change of its visible direction the contrary way, produced by the muscular actions which regulate the mutual distance of the pupils. To know how this happens, suppose the two optic axes to be parallel to each other, and perpendicular to the visual base ; and let a physical line be placed in either of them, so as entirely to coincide with it. This line will, therefore, not only be in reality perpendicular to the visual base, but will, in the present state of things, likewise appear so.—Incline afterward both the axes equally to the left side, and it is manifest that the line coinciding, say, with the right axis, must appear equally inclined. Let now the right axis be kept fixed, and the left be carried back again, and its motion continued, until it be as much inclined toward the right side, as itself was just before, and as the right axis is still to the left side ; the consequence

quence will be, that the line in the right axis must again be seen perpendicular to the visual base ; for such is the present position of the common axis. Here then we have had two opposite causes of change of apparent direction acting in succession. The muscular actions, producing the joint motions of the eyes, first bent the visible position of a line, in the right optic axis, from a perpendicular to the visual base toward the left ; and the muscular actions, which regulate the mutual distances of the pupils, by *increasing* the inclinations of the axes to each other, moved it afterward, from the left to the right, back again to a perpendicular to the visual base. Let these two causes act together, and it is plain, that no observable effect will be produced by either, as long as they are thus proportioned. When they are not so, only the difference of their forces will be exhibited by the phenomena.

But farther ; to show the extent of this theory of visible direction being dependant upon the actions of the muscles of the eyes,

I shall

I shall now apply it to the explanation of an instance of apparent motion, which at first may be thought to furnish an argument against it. Look with one eye, the other being closed, at any remote object through a small hole in a card. If you should afterward suddenly attempt to view the hole itself accurately, with the same eye, you will observe both it and the distant object, particularly the latter, to move from left to right, if the right eye be used ; but if the left eye be the one employed, then from right to left. Shift now your attention as suddenly back from the hole to the object seen through it, and both will return to the places they formerly occupied. In this experiment, no real change can be supposed to have occurred in the position of the distant object ; and had any happened, with respect to either the eye or the hole, the object would not have been seen through the latter. No other fallacy, therefore, exists here, than that things, which are truly at rest, appear, notwithstanding, to be in motion,

The

The argument, which I have mentioned may hence be derived against my theory, is this : The visible directions of objects, in the optic axis which remained fixed, were formerly said to be altered, because a new state of muscular exertion was required to keep it so, in every different degree of the inclination to it of the moveable axis. But in the last experiment, there seems no good reason for supposing any change in the inclination of the moveable axis to the other ; for, as one eye is closed, the obvious intention of directing the two axes to the same object, which is, that we may see it single, no longer exists. If then an apparent lateral motion be, in one instance, observed in objects truly at rest, without any change of the interval of the pupils, may not every other motion of the like kind be also independent of the muscular actions, which regulate that interval ?

It is evident, that this argument rests altogether upon the supposition, that in the experiment just mentioned, no alteration occurs in the interval of the pupils. Now,

we may be easily convinced, that some alteration does occur, by applying a finger to the closed eye, which will, by this means, be felt to move toward the nose, when we endeavour to view the hole accurately, and from the nose, when we carry our attention back again to the remote object. Were, indeed, the opinion of Aguilonius * just, that the mind perceives only those objects distinctly, which are situated at the concurrence of the optic axes, whether they are seen with one or with two eyes, both the necessity and the degree of the alteration would be clearly ascertained. But as this opinion is not just, which I mean to prove from experiments in a succeeding part of this volume, I shall proceed to give another reason, and I think the true one, why the interval of the pupils should be as much altered, when we look with one eye at objects successively, which are placed at different distances, as if we were to view them with both.

It is a fact, for which I have the authority of experiments almost without number,

M

though

* Optica, p. 84.

though I do not recollect to have seen it mentioned by any author beside Dr. Porterfield, that every change of the mutual positions of the optic axes is conjoined, in persons who do not squint, with a change of the power, in both eyes, to refract the rays of light which fall upon them.—

When the axes are parallel to each other, the eyes are in their lowest refracting state ; but in their highest, when the axes are mutually intersected within two or three inches of the face ; every intermediate inclination being also conjoined with an intermediate degree of refracting power. Now, since those objects are seen most distinctly, the rarious pencils from which are accurately brought to points in the retina, it follows, that, although we employ one eye only, the same reason exists for adjusting its refractive power to their distances, as if we saw with both. When, therefore, we view a remote object with one eye, we use it in its lowest refracting state, which, I have observed, is conjoined with the widest interval of the pupils. Should we afterward attempt to

see accurately a very near object, the eye will assume its highest refractive state, and the interval of the pupils be lessened; the consequence of which must be, that both the objects lying in the optic axes will appear to move in the manner already related.

To finish this part of my subject, it seems only necessary to determine, whether the dependance of visible direction upon the actions of the muscles of the eyes be established by nature, or by custom. But facts are here wanting. As far as they go, however, they serve to prove, that it arises from an original principle of our constitution. For Mr. Cheselden's patient saw objects single, and consequently in the same directions with both eyes, immediately after he was couched; and persons affected with squinting from their earliest infancy, see objects in the same directions with the eye they have never been *accustomed* to employ, as they do with the other they have constantly used.

Having thus shown in what directions external bodies are seen, when their situa-

(64)

tion with respect to the eye is given, and upon what circumstance the various directions depend, in which a picture upon any one place of the retina can exhibit the object producing it; I should render the theory of visible direction complete, were I now to point out the relative positions of the two lines of direction, in which any two different parts of the retina represent their objects. To ascertain this, the first step must be, to find the place of the retina which receives the picture of an object, whose situation with respect to the external eye is known; and if two such points of the retina were determined, I think the chief difficulty in this matter would then be overcome. But as it appears to me, that the structure of the eye has not yet been sufficiently explained, to enable any person to take this first step, I forbear saying any thing more upon the subject.

END OF THE ESSAY UPON SINGLE VISION WITH TWO EYES,

EXPE-

EXPERIMENTS *and* OBSERVATIONS

ON

SEVERAL SUBJECTS IN OPTICS.

ARTICLE I.

On Visible Position, and Visible Motion,

IN the estimates we make by sight of the situation of external objects, we have always some secret reference to the position of our own bodies, with respect to the plane of the horizon; and from this cause, we often judge such to be at rest, whose relative places to us are continually changing; and others to be in motion, though they may constantly preserve, in regard to us, the same distance and direction. To give

an instance, let us suppose our eyes first directed to a star near to the horizon; should we afterward, by a mere motion of the head, point them to another, some degrees above the former, this second star will appear higher than the first did. Were we now, while the eyes are kept fixed in relation to the head, and the head in relation to the shoulders, to incline the trunk of the body backward, until we bring the optic axes to a third star, this will appear still higher than the second was perceived to be. If instead of directing the eyes successively to different objects, the same object be suffered to remain at the concurrence of the optic axes in all these different positions of the body, it is evident, that it must be seen to move, during the change from one position to another.

The facts I have mentioned are so obvious, that I should not have spoken of them, had I not intended they should introduce the following question: What is there within us, to indicate these positions of the body? To me it appears evident, that, since they are occasioned and preserved by combinations

combinations of the actions of various voluntary muscles, some feeling must attend every such combination, which suggests, from experience perhaps, the particular position produced by it. But in almost all the positions of the body, the chief part of our muscular efforts is directed toward sustaining it against the influence of its own gravity. Each position, therefore, in which this takes place, must be attended with a feeling, which serves to indicate its relation to the horizontal plane of the earth; and consequently, if our bodies possessed no gravity, or, if the thing were possible, had we been created unembodied spirits, but with the same faculties of perception as we enjoy at present, we could no more have judged one line to be perpendicular, and another to be parallel to the horizon, than we can at present determine, without some external aid, which is the eastern, and which the western point of the heavens. I shall now draw from these principles, the explanation of a fact, which was first mentioned by one of the most ingenious authors

that

that have written upon vision, but left by him still to be justly accounted for.

“ I have frequently (says Mr. Melvill)*
“ observed, when at sea, that, though I
“ pressed my body and head firmly to a cor-
“ ner of the cabin, so as to be at rest in
“ respect to every object about me, the diffe-
“ rent irregular motions of the ship, in roll-
“ ing and pitching, were still discernible by
“ sight. How is this fact to be recon-
“ ciled to optical principles? Shall
“ we conclude that the eye, by the
“ sudden motions of the vessel, is rolled
“ out of its due position? Or, if it retains a
“ fixed situation in the head, is the percep-
“ tion of the ship’s motion, owing to a ver-
“ tigo in the brain, a deception of the ima-
“ gination, or to what other cause?”

I need not, I believe, offer to show, that the fact here spoken of, is not owing to any of the causes Mr. Melvill has specified. I shall therefore, in a few words, point out its dependance upon the principles which have just been mentioned.

It is generally known, I suppose, that when a vessel at sea, in the language of sailors,
is

* Edinburgh Physical Essays, vol. ii. p. 89.

is said to pitch, its two extremities turn upon its shorter axis, and that the term of rolling is confined by them to its motions upon the longer axis. In both pitching and rolling then, the relative position of a vessel to a horizontal plane is necessarily changed. Consequently, though, in the above-mentioned experiment, Mr. Melvill's body and head were at rest with respect to every object about him, still a different degree of muscular effort was required to keep them so, in every such different position of the vessel. But each degree of muscular effort, to sustain his body against the operation of its gravity, would suggest to him its concomitant position with regard to the plane of the horizon ; each deviation, therefore, of the vessel from its former situation, relatively to the same plane, would be perceived, and the vessel itself be seen to move. In short, nothing more takes place in this, than in the following experiment : Let a pole be placed upon firm ground, at right angles to the horizon. If, while we are standing erect, it be inclined upon its

N

lower

lower extremity, successively backward and forward, to the right and to the left, these motions must, without contradiction, be perceived. Suppose now, our bodies to be similarly inclined with the pole, during its different positions, so as to be constantly parallel to it ; it is evident, that its motions will be as readily perceived in this case, as they were, when our bodies were erect ; and this is all that happens in the experiment of Mr. Melvill.

Should the necessity of supporting the body against its gravity, by the actions of our voluntary muscles, be suspended in whole, or in part, our judgments of the situation of objects, with respect to the horizon, must become irregular and uncertain, notwithstanding any general habit we may have acquired from experience. An instance of this, I think, I have observed ; for I have frequently remarked during a sea voyage, that, when the wind blew so strongly, and in such a direction, as to occasion the vessel to heel, or lean much to one side, chords freely suspended from the

roof of the cabin, and kept stretched by heavy bodies attached to them, have appeared to me, as long as I lay in bed, though they were necessarily perpendicular to the horizon, to decline considerably from that position; while the sides of the cabin seemed, if not perpendicular, at least much less inclined to the horizon than they were in reality. My body being here supported by the bed, I was consequently without those feelings, which indicate its position with respect to the horizon. Objects therefore appeared to me in those situations, in which I had been accustomed to see them. In confirmation of which I may mention, that, when I got up, and stood upon the floor of the cabin, the chords seemed perpendicular, or nearly so, and the sides of the cabin inclined; for I was now obliged to exert a proper degree of muscular force, to keep myself upright. What I ^{here} ~~say~~ say, however, is from the recollection of things observed some years ago, when I had no thought of making the use of them I now do; for which reason,

I may possibly have committed some trifling error in stating them ; but none, I believe, sufficient to affect the theory they are brought to support.

It being my intention to treat, in the present article, of several facts relative to visible position and motion, which seem to me to need explanation, without regarding whether or not they depend upon any common cause ; I pass to the consideration of the apparent rotation of objects, when we have become giddy, by turning ourselves quickly and frequently round.

Some of the older writers upon optics imagined the visive spirits to be contained in the head, as water is in a vessel, which therefore, when once put in motion by the rotation of our bodies, must continue in it for some time after this has ceased ; and to this real circular movement of the visive spirits, while the body is at rest, they attributed the apparent motions of objects in giddiness. Dechaies* saw the weakness of this hypothesis, and conjectured, that

* *Curfus Mathematic. Tom. ii. p. 422.*

that the phenomenon might be owing to a real movement of the eyes, but produced no fact in proof of his opinion. Dr. Porterfield,* on the contrary, supposed the difficulty of explaining it to consist in showing, why objects at rest appear in motion to an eye which is also at rest. The solution he offered of this representation of the phenomenon, is not only extremely ingenious, but is, I believe, the only probable one which can be given. It does not apply, however, to the fact which truly exists; for I shall immediately show, that the eye is not at rest, as he imagined. The last author, I know of, who has touched upon this subject, is Dr. Darwin.† His words are, “When any one turns round rapidly on one foot till he becomes dizzy, and falls upon the ground, the spectra of the ambient objects continue to present themselves in rotation, or appear to librate, and he seems to behold them for some time in motion.” I do not indeed pretend

* Treatise on the eye. Vol. ii. p. 426.

† Philosoph. Transact. Vol. lxxvi. p. 315.

pretend to understand his opinion fully; but this much seems clear, that, if such an apparent motion of the surrounding objects depends, in any way, upon their spectra, or the illusive representations of those objects, occasioned by their former impressions upon the retinas, no similar motion would be observed, were we to turn ourselves round with our eyes shut, and not to open them till we became giddy; for in this case, as the surrounding objects could not send their pictures to the retinas, there would, consequently, be no spectra to present themselves afterward in rotation. But whoever will make the experiment, will find, that objects about him appear to be equally in motion, when he has become giddy by turning himself round, whether this has been done with his eyes open or shut. I shall now venture to propose my own opinion upon this subject.

If the eye be at rest, we judge an object to be in motion when its picture falls in succeeding times upon different parts of the retina; and if the eye be in motion, we
judge

judge an object to be at rest, as long as the change in the place of its picture upon the retina, holds a certain correspondence with the change of the eye's position. Let us now suppose the eye to be in motion, while, from some disorder in the system of sensation, we are either without those feelings, which indicate the various positions of the eye, or are not able to attend to them. It is evident, that, in such a state of things, an object at rest must appear to be in motion, since it sends in succeeding times its picture to different parts of the retina. And this seems to be what happens in giddiness. I was first led to think so from observing, that, during a slight fit of giddiness I was accidentally seized with, a coloured spot, occasioned by looking steadily at a luminous body, and upon which I happened at that moment to be making an experiment, was moved in a manner altogether independant of the positions I conceived my eyes to possess. To determine this point, I again produced the spot, by looking some time at the flame of a candle; then

then turning myself round till I became giddy, I suddenly discontinued this motion, and directed my eyes to the middle of a sheet of paper, fixed upon the wall of my chamber. The spot now appeared upon the paper, but only for a moment ; for it immediately after seemed to move to one side, and the paper to the other, notwithstanding I conceived the position of my eyes to be in the mean while unchanged. To go on with the experiment, when the paper and spot had proceeded to a certain distance from each other, they suddenly came together again ; and this separation and conjunction were alternately repeated a number of times ; the limits of the separation gradually becoming less, till, at length, the paper and spot both appeared to be at rest, and the latter to be projected upon the middle of the former. I found also, upon repeating and varying the experiment a little, that when I had turned myself from left to right, the paper moved from right to left, and the spot consequently the contrary way ; but that when I had turned from right to left, the paper would then move from left to right.

These

These were the appearances observed while I stood erect. When I inclined, however, my head in such a manner, as to bring the side of my face parallel to the horizon, the spot and paper would then move from each other, one upward and the other downward. But all these phenomena demonstrate, that there was a real motion in my eyes at the time I imagined them to be at rest ; for the apparent situation of the spot, with respect to the paper, could not possibly have been altered, without a real change of the position of those organs. To have the same thing proved in another way, I desired a person to turn quickly round, till he became very giddy ; then to stop himself and look steadfastly at me. He did so, and I could plainly see, that, although he thought his eyes were fixed, they were in reality moving in their sockets, first toward one side, and then toward the other.

The last instance of visible motion I shall notice, is one which has been mentioned by Mr. Le Cat, in the following words :*

* *Traité des Sens*, p. 419.

“ Place a lighted candle at a moderate dif-
 “ tance from a polished body of confiderable
 “ convexity, fo that the image of the flame,
 “ which is feen by reflection from it, may
 “ appear as a fmall luminous point. The
 “ experiment will fucceed better, if the di-
 “ rect rays of the flame be intercepted
 “ from the fight. Clofe, after this, one
 “ eye, and view the luminous point in a
 “ carelefs way, (*en revant*) that is to fay,
 “ with the eye in a relaxed or dilated ftate.
 “ The point will then be feen enlarged and
 “ radiated. If you bring now your fin-
 “ ger to the right of the eye which is open,
 “ and gradually move it toward the left,
 “ in order to conceal the luminous point
 “ from this eye, you will diftinctly perceive
 “ the fhadow of your finger to proceed from
 “ left to right, and to pafs over the point
 “ in a direction, contrary to that which
 “ you gave it. Should you, afterward,
 “ move your finger back from right to left,
 “ and in like manner, if your finger be
 “ moved from above downward, or from
 “ below upward, the fhadow will always
 proceed

“ proceed the contrary way. It is there-
“ fore manifest, that the soul must here
“ see objects inverted, as their images
“ in the eye truly are; and that it refers
“ impressions to those parts of the eye
“ where it feels them, and not to the places
“ from which the rays are emitted, as
“ it does when it possesses the means
“ of rectifying its judgment. Whence
“ does this happen? Doubtless, because
“ the luminous point has neither a high
“ nor a low, neither a right nor left side,
“ nor any well-enlightened object in its
“ vicinity, to awaken the attention of the
“ soul; in short, nothing which can de-
“ termine its judgment.”

I should scarcely have mentioned this experiment, from any respect for the authority of its author in optics; but as Haller* seems to assent to the conclusion he draws from it, that the soul sometimes sees objects inverted; and as the Abbot Derochon,† a member of that learned body, the Aca-

* *Elementa Physiologiæ*, Tom. v. p. 479.

† *Memoires de Physique*, p. 66.

demy of Sciences of Paris, has lately, but in my opinion unsuccessfully, attempted to reconcile it to the commonly-received principles of vision, I think it worth while to show, in a few words, that it is a direct consequence of the very doctrine Mr. Le Cat means to overthrow by its means.

It would be proper, indeed, to mention before hand, the opinion of the Abbot Derochon; but this I must, notwithstanding, omit doing, as it could not be understood without the figure by which he has illustrated it. I shall observe, however, respecting it, *first*, that it requires the side of the finger next to the eye, to be without the least illumination; whereas the experiment will succeed, whether it be illuminated or not: *secondly*, that, according to it, the experiment ought to succeed equally well, whether the image of the flame in the mirror be seen as a point, or as a surface; though, in truth, it never does succeed, except in the latter case: *thirdly*, that the apparent shadow of the
finger

finger is always much larger than it ought to be, were it seen by reflection, as the Abbot thinks: *fourthly*, that, while the eye, mirror, flame, and finger, remain in the same positions, the shadow seems at one time larger than at another, owing to the different degrees of relaxation in the eye; but that this, for the reason just mentioned, ought never to happen, according to his theory: *fifthly*, that, agreeably to his own reasoning, the shadow ought to move in the same direction with the finger, which is the very reverse of the fact to be explained. But as arguments against error may be infinitely extended, and as only one solution of a phenomenon can be true, the readiest way of exposing the insufficiency of others, is to exhibit that which is just.

This, in the present case, seems to lie upon the very surface of optical knowledge, and has already been given by others, of various forms of the same fact. When the image of the flame is seen in the mirror as a point, its rays must be accurately

rately collected to a focus in the retina ; but when seen as a surface, this must necessarily be attributed to their focus being either before or behind it ; in either of which cases, they will occupy a place upon that membrane of some assignable dimensions. In the present instance, their diffusion over a part of the retina, depends on the focus being behind it ; for the eye is now, from a condition of the experiment, in a more relaxed state than it was just before, when the rays of the same object were brought there accurately to a point. The rays, therefore, which go to the right side of the enlightened surface of the retina, or picture as I shall call it, are those which enter the eye at the right side of the pupil, and its left side is formed of the rays entering at the left side of the pupil ; and the like must be true of its upper and lower parts. Should we then begin to move a finger from right to left across the eye, the rays forming the right side of the picture must be first intercepted. But from the known fact, that the points of an external object are always in an inverted position,

position, with respect to the parts of the retina, by the affections of which they are suggested, when the *right* side of the picture there is effaced, the *left* side of the external object it suggests must disappear. And for the same reason, if the motion of the finger be continued from right to left across the eye, the other parts of the luminous surface in the mirror will successively vanish from left to right, and thereby furnish the appearance of a shadow passing over it in that direction.—In like manner, it may be shown, that if the finger proceeds from left to right, from above downward, or from below upward, the shadow must move the opposite way.

That this is the true explanation of Mr. Le Cat's experiment, is, I think, plain, both from its intrinsic evidence, and the following considerations:—If the mirror be brought within four or five inches of the eye, and the candle be so placed, that the image of the flame must, from the laws of reflection, be regarded as a mere point ;

5 though

though we should now view it with the utmost care, and though there should be in its neighbourhood some well-enlightened object to awaken the attention of the soul, as Mr. Le Cat expresses it, still the seeming shadow will move in a direction contrary to the finger. For the image is now so near to the eye, that no exertion we can make is sufficient to bring its rays to a point upon the retina; the picture, therefore, upon that membrane will be formed of rays passing to a focus behind it, which is the only condition necessary for the success of the experiment. Again, if a short-sighted person should place the mirror at the distance of some feet from him, complying in other respects with Mr. Le Cat's instructions, he will constantly observe the shadow to move in the same direction with the finger. For, in his eye, the rays of the image, when at such a distance, must meet before they fall upon the retina. The right side, therefore, of the picture upon that membrane, must be composed, in this case, of rays which

enter

enter the eye at the left side of the pupil. Consequently, when these are cut off, the left side of the apparent luminous surface must disappear, and the shadow be seen to move the same way as the finger, when this successively intercepts the rays proceeding from the image to the eye.*

* Scheiner observed a fact of the like kind (*Fundamentum Opticum*, p. 33) namely, that, if a small hole, made in any substance, be held near to the eye, and an opaque body be passed between them, from right to left, the left side of the hole will first disappear. Mr. Grey afterward took notice (*Philosoph. Transact.* Vol. xix. p. 286) that a needle he employed in this experiment was seen inverted; from which he supposed that the hole, or something in it, produced the effect of a concave speculum. Mr. Harris, however, says (*Treatise of Optics*, p. 141) that it is not the needle, but its shadow on the other side, which is seen, and is the cause of the inverted appearance. But the truth is, that the hole is to be regarded as a luminous point, the rays of which fall upon the retina before they are collected to a focus; and hence that the same appearances must be here observed as in the experiment of Mr. Le Cat. In proof of this it may be mentioned, that if the hole be placed at such a distance, that the eye may refract its rays accurately to a point on the retina, no shadow or image of the needle will be seen; that if the hole be still farther removed, and the eye be adapted to a less distance, the shadow or image will again appear, but its position will now be upright, and its motion the same way as that of the needle itself; and lastly, that, at one given distance of the hole, either no shadow will appear, or it will be seen upright, or it will be seen inverted, according as the eye may be made to assume different states with respect to its power of refraction.

ARTICLE II.

On a supposed Consequence of the Duration of Impressions upon the Retina; and the Effects of accurate Vision being confined to a single Point of that Membrane:

FEW things, at first, appear more incredible to a person, not conversant in optics, than that he does not, at any one time, see distinctly a surface larger than the head of a pin. After he is convinced, by proper trials, of the truth of this, he naturally asks, Whence comes it then, that, in ordinary vision, I seem to view distinctly so many objects at once? I go into a crowded street, and I fancy I have an accurate perception by sight, of men, houses, carriages, and many other things, all at the same time; whence proceeds this illusion?

Only one answer, as far as I know, has been given to this question. The impressions made upon the retina by external objects, do not, it is said, immediately cease.

cease, along with the reception of the rays which flow from them ; and, as in the ordinary mode of vision, the eye is continually passing from object to object, the impression left by a former one may be still vivid, though the eye be directed to another ; and hence we may imagine we see both of them distinctly, though the picture of only one occupies that place of the retina, which alone furnishes us with accurate vision.

There are, however, objections to this answer, which seem to me insurmountable. For, *in the first place*, as the duration of impressions on the retina must be greater or less, according to the vivacity of the pictures which occasion them, it follows, that, were this answer just, the apparent field of our distinct vision ought to be in proportion to the quantity of light admitted by the eye ; that it should be contracted, therefore, by every cloud which passes over us, and be enlarged by every burst of sunshine ; that, at mid-day, it should possess its greatest extent, and

ought from that time gradually to decrease till the evening, when its limits should be nearly the same with those of the real field of accurate vision. *Secondly*, since the coloured spot, which is produced by looking steadily for some time at a luminous body, appears projected upon every object to which we direct our eyes, during its continuance, and as such a spot is necessarily the sign and effect of the duration of an impression upon the retina; every other visible appearance from the same cause ought, in like manner, to have its situation determined by the position of the eye, as far as this may be occasioned by the action of its muscles. No object, therefore, ought to appear separate and distinct from others, if the answer were true which I am combatting; but, on the contrary, all those to which we successively direct our eyes during the limits of the duration of an impression upon the retina, should seem crowded into one place; and, consequently, none of them should be perceived with any tolerable accuracy.—

Such

Such are the conclusions from the truth of this answer. I need scarcely mention, that they are contradicted by experience.

There is another form of the same fact, to which, it may be thought, an explanation taken from the duration of impressions on the retina will better apply; I mean the appearance of a fiery circle, when any red-hot body is moved quickly round. But it seems, to me, that such an explanation cannot even here be admitted. For, if the circle depended upon the cause I have mentioned, it could only be observed as long as the impressions upon the retina were also disposed in the form of a circle. Were this broken upon, which it must be by every movement of the eye, the appearance suggested by the last impression would no longer be so arranged, with respect to the appearance suggested by the present impression, as to lie with it in the circumference of a circle; and hence some very different figure would be observed. Every person, however, may easily convince himself, that the circular form of
the

(115)
the fiery appearance is equally perceived, whether the eye be at rest, or be moved in the most irregular manner.

If these arguments be thought sufficient for the purpose I had in view, it must also follow from them, since the fact still remains to be explained, why we apparently see so many objects with equal distinctness at once, that past impressions upon the retina are perceived as present, by means of some higher faculty than that of sight. This faculty cannot, with propriety, be named *memory*, as it is essential to a thing's being remembered, that it be perceived as past. Nor can it be called *imagination*, since we believe in the present existence of what it perceives. In one point of view it may seem rather a defect in our natures, that we should not be able to distinguish between things past and present. However this may be, I am inclined to be of opinion, that many other phenomena, both of thought and external sense, are partly to be resolved into the same general fact. From the present instance of it,

it, we learn, that several muscular actions may be performed, in succession, during the least perceptible portion of time.

The question I have just treated, naturally gives rise to another: Would it have been more to our advantage, if accurate vision, instead of being confined to one point of the retina, had been possessed by every part of that membrane? I answer, I think not, for the following reasons.

First; The diffusion of such a property over the whole retina would be of little use, unless our power of attention was also increased. For we would otherwise be still unable to perceive more than one visible object at once, with distinctness, since, by our present constitution, we are capable of attending accurately to only one thing at a time. The only benefit, indeed, I can see to arise from such a condition of the retina, is this, that our attention might be shifted more quickly from picture to picture on that membrane, than our eyes can be turned from one external object to another.

another. This advantage, however, would be far out-weighed by an inconvenience accompanying it. For it is a well-known fact, with respect to perception, that we are capable of attending, more or less accurately, to any particular impression upon the senses, in proportion to the force of the other impressions, which are at the same time received. But in the supposed state of the retina, there would be, almost always, several impressions of the same strength as the one to which we might desire particularly to attend; whereas, in its present state, the vivacity of the impression from the object, to which we turn the optic axis, most commonly surpasses, considerably, that of every other upon the same membrane; by which means our attention is rendered less liable to interruption.

Secondly ; The extension of accurate vision, to every part of the retina, would deprive us, in great measure, of the help, which we obtain, at present, from the eye, in learning the thoughts of other men.

As

As far as I have been able to observe, the changes produced by our internal feelings, upon the state of the eye itself, are very few, and relate only to the quantity of moisture, which is diffused over its surface, and the degree of fulness in the blood-vessels, which are spread upon its white and glistening part. Both of these circumstances, however, are similarly altered by opposite passions, and, consequently, neither of them can be regarded as the appropriate expression of any. The whole variety, then, of the expressions of feeling, which are justly attributed to the eye, must, I think, depend upon its motions. Some of these are the immediate effects of certain passions; the eye, for instance, being moved differently in anger and in grief; and such may be esteemed as directly expressive of the passions by which they are produced. But the far greater number of them do little more, than merely point out the external cause, or object of the sentiment, which the changes of other parts of the countenance declare to

Q

exist

exist within us ; or distinguish certain external appearances depending upon a mental cause, from similar appearances arising from a different source. Thus, blushing is often distinguished from an accidental flush of the cheek, by the eye being turned away from the person who occasions it.

That many of the expressions, which we attribute to the eye, do in fact depend on changes in other parts of the countenance, is evident from the alterations we think induced upon it, by the eye-lashes falling off from disease, by a slight inflammation of the edges of the eye-lids, without its being communicated to the eye itself, by artificially colouring the eye-brows, and by many other similar circumstances. And how essential to the right understanding of the expressions of the other features, are the motions of the eyes, when conducted with design, and properly directed, must be known to every one, who has attended in discourse to the countenances of very short-sighted people, and more especially to those of persons afflicted with

blindness

blindness from a *gutta serena*, in which the eye, with respect to its external condition, seems without fault. But whatever is the assistance the motions of the eye afford, in expressing our internal feelings, the whole of it must ultimately be referred to the circumstance of accurate vision being confined to one point of the retina ; since the intent of those motions is, to bring the pictures of external objects upon the most sensible part of that membrane. Their necessity, therefore, would no longer exist, if the same property were extended, and the advantages we at present enjoy from them would, consequently, cease.

ARTICLE III.

*On the Connection between the different refractive States of
the Eyes, and the different Inclinations of the
Optic Axes to each other.*

I HAVE mentioned, in my Essay upon Single Vision with Two Eyes,* that I had been convinced, by experiments almost without number, that every different degree of the mutual inclination of the optic axes, is attended by a different state of the refracting power of each eye. The experiments I there alluded to were chiefly of this sort. I placed a luminous point, most commonly the reflected image of the flame of a candle from the bulb of a small thermometer, at such a distance, that when both my eyes were accurately directed to it, its visible appearance to one of them was likewise that of a point. Keeping then the axis of this eye fixed, and making the other to cross it, sometimes before and sometimes behind the luminous point, I found that in
both

both cases it appeared as a surface to the eye, in the axis of which it was situated; and that the more remote from it was the concurrence of the axes, the larger was the luminous surface. Now when the axes met before the point, the apparent surface must have been occasioned by the rays coming to a focus, previously to their incidence upon the retina; because, when I passed my finger across the eye by which it was seen, its parts disappeared, in an order corresponding to the direction in which the finger moved. The disappearance of the parts was in an order, contrary to the motion of the finger, when my optic axes intersected each other beyond the point; which is an equal proof, that the rays, in that case, tended to a focus behind the retina.

One application of this fact has already been shown,* and I shall now proceed to mention several other phenomena in vision, which it may serve either in whole, or in part, to explain.

I. It

* Essay upon Single Vision, p. 83.

1. It accounts for the following beautiful observation made by Aguilonius,† that if we close one eye, and look with the other at an object placed in its own axis, we will not be able to see this object distinctly, unless we also direct to it the axis of the closed eye. For in persons, who are neither presbytic nor myopic, the refractive states of the eyes are so adapted to the mutual inclinations of the optic axes, that pencils of rays flowing from bodies at moderate distances are more accurately collected upon the retina, when they are situated at the intersection of those lines, than if their position was, in any considerable degree, either nearer or more remote. The reason given by Aguilonius himself, is, that the mind perceives only those objects distinctly, which are placed at the concurrence of the optic axes. But the following experiment proves that the solution is true no farther, than as it coincides with the one I have advanced. Hold, in the axis of either eye, a concave lens, at
such

† Aguilonii Optica, p. 24.

such a distance, that the letters of a book, placed a little farther off, may appear through it very indistinct to that eye, when both axes are directed to any particular word. View afterward the lens itself with both eyes, and the letters will immediately become more distinct. In this experiment then, an object is more accurately perceived when distant from the concurrence of the optic axes, than when situated exactly in it.

It may be said, perhaps, that the distinctness of the letters is here to be attributed to the contraction of the pupil, which is occasioned by the eyes being directed to a nearer object than they were formerly. But that this is not the case, may be made evident by another experiment: Place a convex lens in such a manner before one eye, that the flame of a candle, at the distance of two or three feet from the face, may appear indistinctly terminated to that eye, when both axes are pointed to it. The same eye being kept fixed, let the two axes afterward meet
beyond

beyond the flame, and it will now be seen much better defined, though the pupil is at the same time become larger. The insufficiency of the explanation of Aguilonius, is also proved, by a circumstance frequently noticed in persons who are very short-sighted; for such are observed, when they desire to view an object with much attention, to hold it close to one eye, and to turn the other aside; in this way occasioning the two axes to meet very remotely from the object.

2. The reason commonly given, why short-sighted people view an object with one eye only in the manner above-mentioned, is, that by this means they avoid the uneasy straining of the muscles, which must be employed to direct both axes to the same point. But it is evident they must derive from the practice this farther advantage, that, as their optic axes are now parallel to each other, or nearly so, they, consequently, see the object in the least refractive state of their eyes. Pencils, therefore, will now have their focuses in the retina,
the

the rays of which would have crossed each other, before they fell upon it, had both the axes been directed to the object.

3. Spectacles were long employed, before the manner in which they assisted sight was known. About the year 1601, this was proposed as the subject of a question to Kepler,* by his principal patron at that time, Ludovic L. B. a Dietrickstein, a learned nobleman of Austria. The first answer he gave was, that convex glasses were of use, by occasioning objects to appear larger. But his patron observed, that if objects were rendered by them more distinct, because larger, no person would be benefited by concave glasses, since these diminish objects. It was not till three years after, that, in consequence of finding out in what manner vision is performed, he was able to give a just solution of this problem, though his attention had been directed to it during the whole of that interval. According to the discovery he then made, convex glasses were said by him to assist the sight of pres-

R

bytic

* Paralipomena in Vitellionem, p. 206.

bytic persons, by so altering the directions of rays diverging from a near object, that they shall afterward fall upon the eye, as if they had proceeded from a more remote one; and concave glasses to benefit the myopic, by producing a contrary effect upon rays which diverge from a distant object. Now it is manifest, that by this theory, to which I believe no addition has been made by any succeeding writer, precisely the same effects are attributed to lenses, whether they be employed singly, or in the form of spectacles. I am inclined, however, to think, that a difference, sometimes at least, exists here, which has hitherto escaped notice. For in regard to such spectacles as I have tried upon myself, I have always found, that, when I looked with them at objects placed at moderate distances directly before me, my optic axes passed through the glasses, more inwardly than their centres. With respect, therefore, to spectacles for long-sighted people, as the inner halves of their glasses may be regarded as two prisms, whose refracting angles face each

other,

other, to have allowed both my eyes to receive through them pencils of rays from the same point of an object, the intervals of my pupils must have been less than was necessary for that purpose in naked vision. The consequence of which would be, an increase of the refractive power of my eyes. Again; as the like parts of glasses in spectacles for short-sighted persons, may be esteemed to be two prisms, the refracting angles of which are turned from each other, the interval of the pupils must have been increased, and the refracting power of my eyes by this means diminished, when I looked at an object through them, which was directly before me. And effects similar to what I have mentioned, must have followed my viewing objects placed obliquely, through glasses of both kinds. Here then is one advantage, which persons, who see with both eyes, either do or may enjoy from spectacles, but which they cannot derive from using single glasses. For if they are presbytic, they can see an object by the means of them with a higher refractive state

of the eyes, than if the optic axes met there, as in naked vision ; and if myopic, with a lens. It is also worthy of remark, that this advantage does not ultimately tend to increase the evil, which first gives occasion for spectacles. On the contrary, if what every writer upon vision asserts be true, that we are apt to become short or long-sighted, according as we are much accustomed to view near or distant objects, it must serve to diminish that evil. In support of this opinion, I shall mention a fact, with which I have been made acquainted by Mr. George Adams,* of this place, who is not only well skilled in the theory of vision, but, from his situation, as an artist, has better opportunities, than most persons, of learning such matters. The fact is this, that he does not know a short-sighted person, who has had occasion to increase the depth of his glasses, if he began to use them in the form of spectacles ; whereas he can recollect several instances, where those have been obliged to change their

concave

* Mathematical Instrument Maker to the King.

concave glasses repeatedly, for others of higher powers, who had been accustomed to apply them to one eye only. This indeed may have happened by accident; but, at any rate, the fact is worthy of farther attention and inquiry.

It would seem, however, that the long-sighted derive more benefit from the alteration in the mutual inclinations of the optic axes, which is produced by spectacles, than the short-sighted. For, as the inner halves of the convex glasses are to be regarded as prisms, with their refracting angles continually increasing as we approach their edges, if two objects, situated at different distances, be viewed successively through them, the inclination of the optic axes to each other, when the nearer object is seen, must bear a higher proportion to their inclination, when we look at the one more remote, than the different inclinations of the optic axes do to each other, when they are successively directed to the same objects, without the intervention of such glasses. Hence the nearer the object
is,

is, the greater will be the effect of the variation in the inclination of the axes produced by spectacles with convex glasses; which is the order of things, the best adapted to the wants of those who use them. But with respect to short-sighted persons, since the refracting angles of their glasses, considered as prisms, *decrease*, in proportion as the objects seen through them become more remote; they must, consequently, derive the least benefit from an alteration in the mutual inclinations of the optic axes occasioned by their spectacles, at the time they most require it.

If it were asked, then, what is the real foundation of the common reproach against spectacles for long-sighted people? I should answer, a very different one from that, which is, for the most part, assigned.—For the change, in the conformation of the eyes, which renders them useful, seems to be one of those which nature has destined to take place at a particular age, and to which there is no gradual approach through the preceding course of life. A person, for instance,

instance, at forty, sees an object distinctly, at the same distance that he did at twenty. When he draws near to fifty, the change I have spoken of commonly comes on, and obliges him in a short time to wear spectacles. As it proceeds, he is under the necessity of using others with a higher power. But, instead of supposing that his sight is thus gradually becoming worse, from a natural process, he attributes the increase of the defect in it to his too early and frequent use of glasses. Upon the whole, I should draw this inference from what has been said, that no person, whose sight begins to grow long, ought to be, in the least, prevented from enjoying the immediate advantage which spectacles will afford him, by the fear that they will ultimately injure his eyes; not that I think the convexity of each glass, considered by itself, can do no harm, but that I believe the benefit, arising from the combination of the two, to be at least sufficient to compensate it. Whether those, who have a tendency to short-sight, should be also early in their employment of spectacles,

spectacles, I shall not pretend to say; as there is not the same ground, from theory, for supposing, that the benefit arising from the combination of the two glasses is able to over-balance the injury, produced by the concavity of each considered separately.

All that I have said, however, upon the subject of spectacles, proceeds upon the supposition, that, when objects, placed directly before us, at moderate distances, are viewed through them, the optic axes penetrate the glasses more inwardly than their centres. But I can be by no means sure, that the interval of the pupils of other persons bears the same proportion, to the interval of the centres of the lenses in spectacles, as that of mine does. It concerns those, therefore, who are choosing them, to have attention to this circumstance. To me it appears proper, that the glasses in spectacles, both for long and short-sighted people, should be so far asunder, that, when we look at a very remote object directly before us, our optic axes may pass exactly through their centres. For if the
centres

centres of convex glasses be nearer to each other, very remote objects will appear double ; and if they are more distant, though the object viewed be infinitely far from us, the optic axes will, however, be inclined to one another, and the refractive power of the eyes increased, when this may be of disservice ; since there are few eyes which are not able, even without the aid of the convexity of a glass, to bring parallel rays to a focus upon the retina. If the centres of lenses in spectacles, for the short-sighted, be less distant than what I have mentioned, the optic axes must be bent toward each other, when very remote objects are seen, and the refractive state of the eye, therefore, heightened, which is the very reverse of what is here to be desired. Should the interval of the centres of those lenses be greater, objects at very considerable distances will be seen double.

There are two other observations relative to glasses for the sight, which I wish to add to what I have already said upon this subject. The first is, that the single con-

(130)

convex glasses with which some persons read, must be very injurious, if they be sufficiently large, to admit the same object to be seen with both eyes. For as both axes will then pass through them, one on each side of the centre, the interval of the pupils will be widened, and the refracting power of the eyes, be diminished; so that here a disadvantage is to be added to the prejudice of the convexity of the glass, not a benefit to be placed against it, as in the case of common spectacles for the long-sighted. If, indeed, the defect in sight does not arise from the conformation of the eye, but from a want of transparency in its cornea or humours, then such glasses, by magnifying objects, will be useful, for the same reason, that, in a very faint light, we can read a book of a large print, with more ease than one of a smaller. The second observation is, that if flat-sided prisms were fixed in spectacle-frames, with their refracting angles toward each other, they would assist the long-sighted somewhat, without producing the evil which is said to arise

arise from the convexity of lenses ; and spectacles of this kind might, with more propriety, I think, than any others, be called *preservers*. A like combination of such prisms, but with their angles turned the other way, might, when the object was moderately distant, be of service to the short-sighted. But objects, very remote, would be made by them to appear double.

A R T I C L E IV.

On the Limits of perfect or distinct Vision.

DR. Jurin,* I believe, was the first who distinguished between *perfect* and *distinct* vision ; confining the former term to those cases, where the rays of a single pencil are collected to a single point of the retina ; and marking, by the latter, the perception we have of visible objects, when the rays of the pencils, diverging from them, though not collected to single points of

* Essay on distinct and indistinct Vision.

432

the retina, yet occupy so small portions of it, as to allow the objects to be distinctly seen. But as few authors have adopted this division, I shall, in the present article, use both terms in the sense, which he has appropriated to the first. Neither of them is indeed free from objection, since bodies to be distinctly or perfectly seen, not only require, that their pictures should be accurately formed upon the retina, but that they should fall upon a particular part of it.

Although it has long been a subject of inquiry, within what limits of distance objects are distinctly perceived by sight, yet the only experiments I have met with in books, which have been made, with any tolerable show of accuracy, to determine this matter, are those of Dr. Porterfield. I shall not here say what they were, as his Treatise is in every body's hands, but shall only mention, that the principal conclusions which he drew from them were, *first*, that objects could be distinctly seen by him, that is, the pencils of rays
which

which came from them could be accurately collected to points upon the retina, when their distances from his eye did not exceed twenty-seven inches, and were not less than seven; and *secondly*, that, as often as the axes of both eyes were directed to any one point, situated within those distances, the rays proceeding from it had their focus in each retina.

As the results of some experiments, which I have made upon the same subject, differ from these conclusions, of Dr. Porterfield, I have read over what he has written upon the matter with more than ordinary attention, and I think I can thence show reason, why they should not be received without caution. For, *in the first place*, his experiments are related so circumstantially, and with such an appearance of accuracy in the making of them, that you would scarcely suppose he left the least possible room for error. And yet, after finishing his account of them, he tells us, that he would have repeated them *with more care and exactness*,* had he not been

* Treatise on the Eye, Vol. I. p. 423.

been interrupted. *Secondly*, his experiments were made upon one eye only, though his conclusions apply to both eyes; an inaccuracy which gives occasion to suspect others. *Lastly*, he says, that he could not see an object distinctly at the distance of seven inches, unless both axes were pointed to another object, at only half that distance. Had he then directed both axes to an object seven inches distant, which he does not mention he ever did, it must consequently have been seen *indistinctly*; and yet one of his conclusions states, that objects, distant from about seven, to about twenty-seven inches, were always *distinctly* seen, when the axes of both eyes were directed to them. Such are the reasons which lead me to think, that the whole of the difference, between the results of the experiments of Dr. Porterfield and myself, is not to be attributed to a difference in the structure of our eyes.

The experiments, which I made upon this subject, were with luminous points. They proved to ~~me~~, *first*, that, when both
 optic

optic axes are directed to any object, placed at a less distance from my eyes than about seventeen inches, my vision of it by the left eye is indistinct, from the rays of light tending to focuses behind the retina ; *secondly*, that my vision by the same eye is perfect, if the object seen, and to which both axes are turned, be from about seventeen to about nineteen inches distant ; *thirdly*, that the vision of my left eye becomes again imperfect, if the object be moved to a greater distance than that of nineteen inches, the rays being now collected to focuses, previously to their falling upon the retina ; and *fourthly*, that I have, by my right eye, imperfect vision of all objects, to which I direct both axes, unless their distances be so great, that the rays of each pencil, proceeding from them, may be regarded as parallel.

A conclusion is furnished by these experiments, similar to one, which was drawn by Mr. Delahire, * from some made by himself ; namely, that each eye sees objects dif-

* Memoires de Mathematique et de Physique, 4to. p. 298.

distinctly only at one distance ; as I take for granted, that, in every case of ordinary vision, both axes are directed to the object which is viewed. But Mr. Delahire drew a second conclusion from his experiments, which he seems to have regarded only as another expression of the first, but which, in truth, includes a very different fact. It was, that the refractive state of the eye is always the same, whether we look at a very near or a very distant object. The following observations, however, will prove the contrary, at the same time that they show, in what I farther differ from Dr. Porterfield.

1. Though an object, to which both axes are pointed, does not appear distinct to my left eye, unless it be from about seventeen to about nineteen inches distant ; nor to my right eye, unless it be at a very considerable distance ; yet I find, that when the axes are made to meet at a point, about two inches distant from a line connecting the two pupils, which however cannot be effected without much straining, my left eye

eye

eye will now see an object distinctly, which is only about seven inches from it, and my right eye will at the same time see an object distinctly, the distance of which is about ten inches. I find also, that my left eye is made to see an object distinctly, though placed more than nineteen inches from it, if I direct both axes to a point still more remote.

2. I formerly mentioned, that every degree of the mutual inclination of the optic axes is attended, by a particular state of the refracting power of each eye. But I must now remark, that these states are sometimes subject to slight variations, while the inclinations of the optic axes to each other remain the same. For I find, that, when a luminous point, to which both axes are turned, is distinctly seen by my left eye, I can, by certain efforts not easily to be described, but without changing the position of either axis, make it afterward appear as a surface, and this too, at one time, from the rays coming to a focus too soon, and at another, too late, for perfect

T

vision

vision.* One instance of these variations deserves to be minutely described, as it proves, that the refractive power of the eyes is subject to greater changes, than what are shown by any experiments I have met with in authors. When I look attentively at a bright star, with the optic axes parallel to each other, it appears to my left eye a surface of some extent, and to my right eye, though not a point, yet a surface of very small extent, as small as the sphericity of the cornea and crystalline, the various refrangibility of the different kinds of light, and the width of the pupil at night, can be supposed to allow; for I find, that, if I now pass a needle across the axis of the right eye, its shadow will not be seen. But should I,
after

* The variations, however, seem produced in such a manner, that the middle of the set belonging to one degree of the mutual inclination of the optic axes, is always different from the middle of the set belonging to another degree of their inclination; and that, when no other effort is made, than to direct both axes to the same object, the eyes always assume the middle state of the refractive power, which accompanies that particular inclination of the axes. No argument, therefore, can hence be derived against the applications I formerly made of the general fact, respecting the connection of the refractive states of the eyes with the mutual inclinations of the optic axes,

after this, withdraw my accurate attention from the star, and view it in the state of sight we have, when we are said to be in a *reverie*, in which, though our eyes are open, we are yet scarcely conscious of seeing surrounding objects, the appearance to the right eye expands itself, and if a needle be again passed before this eye, its shadow will be observed to move over the star, in a direction contrary to that of the needle itself; a sure indication that the rays of light now tend to a focus behind the retina. In the same state of things, the appearance of the star to the left eye contracts, and if a needle be held before the eye, no shadow is seen; a sign that the rays are collected to a focus on the retina; whereas they had formerly crossed one another before they reached that membrane.

Upon the whole then it is manifest, from the experiments I have related, that my left eye can collect to focuses in the retina, rays which proceed from objects at every distance whatsoever, not less than seven inches; that my right eye can collect to fo-

focuses in the retina, rays which proceed from objects at every distance whatsoever, not less than ten inches, and even such as are somewhat convergent, since it can make those, which are parallel, to meet before they fall upon the retina; and lastly, that, while both the optic axes are directed to a point within the limits of distinct vision, the rays proceeding from it are never accurately collected to focuses in both retinas, and scarcely ever to a focus in either retina. These are likewise the principal circumstances, in which my experiments differ in their results from those of Dr. Porterfield.

In making such experiments with luminous points, one or other of two appearances very constantly occurs, neither of which, as far as I know, has been spoken of by any preceding author. The most proper way of mentioning what they are, is, perhaps, to show what ought to happen in those situations, in which they are observed.

When a beam of white light passes, obliquely, from one medium into another of different refractive power, its variously coloured rays must begin to diverge from each other, at the point of the beam's incidence upon the latter medium. In achromatic telescopes, the mutual separation of these rays is checked, and its farther increase prevented, before it becomes perceptible to sense, by the contrary refractions which they undergo, from passing, successively, through the different parts of the object-glass. Hence, some have imagined, that, since objects, in ordinary vision, are seen without colour, as far as this depends on the refractions of the eye, nature has furnished us with an instrument, constituted upon principles similar to those of the object-glass of an achromatic telescope. But every one, the least acquainted with the structure of the eye, must know, that this cannot be the case, as the refractions in it are all made one way.* And there are experimental proofs

* There are indeed some exceptions to this, but not of sufficient consequence to affect the present argument.

proofs, that compounded light is always separated into its parts, by passing through the eye. For if we interpose any opaque substance between us and a luminous body, so that only a very small portion of this may remain visible, it will appear to consist of three differently coloured parts, red, yellow, and blue. The reason, therefore, of objects being, for the most part, seen colourless, must be elsewhere sought.*

Now let us suppose, that a luminous point is the only object which is seen at any one time ; should the focus of its mean refrangible rays be anterior to the retina, the middle of its picture upon that membrane must be chiefly composed of the less refrangible rays ; and this must be the reason, that, when I look attentively - at a bright star with my left eye, the centre of it always appears of a light orange colour. As the beams, however, from the luminous point, which enter the eye near to its axis, suffer
but ..

* Dr. Maskelyne has very learnedly treated this subject in the Philosophical Transactions, Vol. lxxix. part 2.

but little refraction, the brightness of their white light, will, in great measure, overpower the colour given to the middle of the picture upon the retina, by the less refrangible rays of those, which enter the eye at a distance from its axis. Were you then to intercept the former beams, the effect I have mentioned of the latter, must be more observable : and hence it is, that when I place a pin or needle between my eye and a luminous point, the rays of which come to a focus before they fall upon the retina, the shadow, instead of appearing black, is always of a red or deep orange colour ; which is one of the phenomena respecting luminous points, to which I have alluded.

On the other hand, should the focus of the mean refrangible rays of a luminous point lie behind the retina, the middle of the picture there will be principally formed of the more refrangible rays ; and if the beams, which enter the eye near to its axis, be also in the present case intercepted, the effect of the latter rays, in giving colour

to the middle of the picture, will consequently be rendered more evident. Hence it is, that, when a luminous point is not sufficiently [remote for distinct vision, the seeming shadow upon it, occasioned by any small opaque object held before my eye, is always blue; and this is the second of the appearances, which I said are frequently to be observed, in experiments upon luminous points.

THE END.

